

Nash County Emergency Medical Services



Clinical Standards and Practice Guidelines

“Making a Positive Difference”

**Nash County EMS System Policies,
Procedures, Protocols and associated
References**

Revised 1/2022

Nash EMS 2022 Treatment Protocol Index

Protocol Introduction PI (Black)

PI-1. Introduction

PI-2. Key to Protocol Utilization

Universal Protocols UP (Light Green)

UP-1. Universal Patient Care

UP-2. Triage

UP-3. Abdominal Pain / Vomiting and Diarrhea

UP-4. Altered Mental Status

UP-5. Back Pain

UP-6. IV or IO Access

UP-7. Dental

UP-8. Emergencies Involving Indwelling Central Lines

UP-9. Epistaxis

UP-10. Fever / Infection Control

UP-11. Pain Control

UP-12. Police Custody

UP-13. Seizure

UP-14. Suspected Stroke

UP-15. Suspected Sepsis

UP-16. Syncope

UP-17. Behavioral Health Crisis

UP-18. Behavioral Agitation / Sedation Guide

UP-19. Behavioral Excited Delirium Syndrome / Violent

Airway Respiratory Section AR (Light Blue)

AR-1. Adult Airway

AR-2. Adult, Failed Airway

AR-4. COPD / Asthma

AR-5. Pediatric Airway

AR-6. Pediatric Failed Airway

AR-7. Pediatric Respiratory Distress

AR-8 Post-intubation / BIAD Management

AR-9. Ventilator Emergencies

AR-10. Tracheostomy Tube Emergencies

Adult Cardiac Section AC (Dark Blue)

AC-1. Asystole / Pulseless Electrical Activity

AC-2. Bradycardia; Pulse Present

AC-3. Cardiac Arrest; Adult

AC-4. Chest Pain: Cardiac and STEMI

AC-5. CHF / Pulmonary Edema

AC-6. Adult Tachycardia Narrow Complex (≤ 0.11 sec)

AC-7. Adult Monomorphic Tachycardia Wide Complex (≥ 0.12 sec)

AC-8. Adult Polymorphic Tachycardia WIDE (≥ 0.12 sec) Torsades de pointes

AC-9. Ventricular Fibrillation Pulseless Ventricular Tachycardia

AC-10. Post Resuscitation

AC-11. Team Focused CPR (Optional)

AC-12. Onscene Resuscitation / Termination of CPR (Optional)

AC-14. Mechanical Circulatory Support LVAD, RVAD, Bi-VAD

Adult Medical Section AM (Olive Green)

AM-1. Allergic Reaction / Anaphylaxis

AM-2. Diabetic; Adult

AM-3. Dialysis / Renal Failure

AM-4. Hypertension

AM-5. Hypotension / Shock

Adult Obstetrical Section AO (Dark Purple)

AO-1. Childbirth / Labor

AO-2. Newly Born

AO-3. Obstetrical Emergency

Trauma and Burn Section TB (Red)

TB-1. Blast Injury / Incident

TB-2. Chemical and Electrical Burn

TB-3. Crush Syndrome Trauma

TB-4. Extremity Trauma

TB-5. Head Trauma

TB-6. Multiple Trauma

TB-7. Radiation Incident

TB-8. Spinal Motion Restriction

TB-9. Thermal Burn

TB-10. Traumatic Arrest (Optional)

Pediatric Cardiac Section PC (Light Purple)

- PC-1. Pediatric Asystole / PEA
- PC-2. Pediatric Bradycardia
- PC-3. Pediatric CHF / Pulmonary Edema
- PC-4. Pediatric Cardiac Arrest
- PC-5. Pediatric Tachycardia Narrow Complex
- PC-6. Pediatric Ventricular Wide Complex
- PC-7. Pediatric Ventricular Fibrillation / Pulseless VT
- PC-8. Pediatric Post Resuscitation

Pediatric Medical Section PM (Grey Blue)

- PM-1. Pediatric Allergic Reaction
- PM-2. Pediatric Diabetic
- PM-3. Pediatric Hypotension / Shock

Toxin-Environmental Section TE (Gold)

- TE-1. Bites and Envenomations
- TE-2. Carbon Monoxide / Cyanide
- TE-3. Drowning
- TE-4. Hyperthermia
- TE-5. Hypothermia / Frostbite
- TE-6. Marine Envenomation / Injury
- TE-7. Overdose / Toxic Ingestion
- TE-8. WMD – Nerve Agent Protocol

Special Circumstances Section SC (Black)

- SC-1. Suspected Viral Hemorrhagic Fever - Ebola
- SC-2. High Consequence Pathogens
- SC-3. Hospice or Palliative Care Patient
- SC-4. Mass Vaccination/Immunization Medication Distribution

Special Operations Section SO (Gold)

- SO-1. Scene Rehabilitation: General (Optional)
- SO-2. Scene Rehabilitation: Responder (Optional)

Introduction

The following medical treatment protocols are developed for North Carolina EMS agencies. The process has evolved since 2007 and continues with input from Medical Directors, EMS Administration, North Carolina Chapter of Emergency Physicians Protocol Committee, North Carolina Office of EMS, EMS field personnel and the public at large through on-line surveys, public meetings across North Carolina and direct communication with stakeholders. The 2017 update expands on the 2012 and 2009 version and continues to incorporate evidence-based guidelines, expert opinion and historically proven practices meant to ensure that citizens and visitors of North Carolina will continue to be provided the highest quality pre-hospital patient care available. The North Carolina Chapter of Emergency Physicians develops and provides final approval.

The purpose of the protocol section is to provide treatment protocols outlining permissible and appropriate assessment, delivery of care, reassessment and procedures which may be rendered by pre-hospital providers. The protocols also outline which medical situations require direct voice communication with medical control. In general treatment protocols are specific orders which may and should be initiated prior to contact with Medical Control.

Please note the medical protocols are divided into three (3) to four (4) sections. The upper section includes three (3) boxes (History, Signs and Symptoms and Differential) which serves as a guide to assist in obtaining pertinent patient information and exam findings as well as considering multiple potential causes of the patients complaint. It is not expected that every historical element or sign / symptom be recorded for every patient. It is expected that those elements pertinent to your patient encounter will be included in the patient evaluation.

The algorithm section describes the essentials of patient care. Virtually every patient should receive the care outlined in this section, usually in the order described. However each medical emergency must be dealt with individually and appropriate care determined accordingly. Professional judgment is mandatory in determining treatment modalities within the parameters of these protocols. Circumstances will arise where treatment may move ahead in the algorithm, move outside to another protocol and then re-enter later. While protocols are written based on body systems and primary complaints the patient should be treated as a whole and therefore the protocols should be considered as a whole in providing care.

Professional judgment hierarchy:

The pre-hospital provider may determine that no specific treatment is needed;

Or

The pre-hospital provider may follow the appropriate treatment protocols and then consult Medical Control;

Or

The pre-hospital provider may consult Medical Control before initiating any specific treatment.

Some protocols will encompass two (2) pages. Protocols which exist in a single page format may have page 2 added by the local medical director. The PEARLS section will either be located at the bottom of page 1 (single page protocol) or page 2 (double page protocol). The PEARLS section provides points regarding the main protocol based on evidence to date, common medical knowledge and expert medical opinion.

Information boxes highlighted in purple. These areas are editable at the local level. They will mainly involve specific medications and dosages utilized by the local EMS agency. Page 2 will have a large section highlighted in purple where the local Medical Director may edit as they see fit to provide expanded points and treatment not otherwise specified in the algorithm. If the box is not to be utilized – add “***This Space Left Blank Intentionally.***”

Finally these medical treatment protocols are established to ensure safe, efficient and effective interventions to relieve pain and suffering and improve patient outcomes without inflicting harm. They also serve to ensure a structure of accountability for Medical Directors, EMS agencies, pre-hospital providers and facilities to provide continual performance improvement. A recent report of the Institute of Medicine calls for the development of standardized, evidence-based pre-hospital care protocols for the triage, treatment and transport of patients. These protocols establish expectations of pre-hospital care in North Carolina.

Introduction

Authors:

R. Darrell Nelson, MD

Medical Director Davie and Stokes County
Assistant Medical Director Forsyth County
Assistant Professor
Wake Forest University
Department of Emergency Medicine
Co-Chairman NCCEP EMS Committee
Chairman NCCEP Protocol Committee

Doug Swanson, MD

Medical Director Mecklenburg County EMS Agency
Medical Director MedCenter Air CCT
Associate Professor
Carolinas Medical Center / UNC-Chapel Hill
Department of Emergency Medicine
Co-Chairman NCCEP EMS Committee

Roberto (Bobby) Portela, MD

Medical Director Pitt County EMS
Clinical Assistant Professor
East Carolina University
Department of Emergency Medicine

Eric Hawkins, MD

Medical Director Union County EMS
Carolinas Medical Center

Juan March, MD

Professor and Chief, Division of EMS
East Carolina University
Department of Emergency Medicine

Jason Stopyra, MD

Medical Director Randolph County EMS
Medical Director Surry County EMS
Assistant Professor
Wake Forest University
Department of Emergency Medicine

Jeff Williams, MD

Deputy Medical Director Wake County EMS
Assistant Professor
UNC-Chapel Hill
Department of Emergency Medicine

Bryan Kitch, MD

Medical Director Hyde County
Assistant Professor
East Carolina University
Department of Emergency Medicine

Seth C. Hawkins, MD

Medical Director Burke County EMS
Assistant Professor
Wake Forest University
Department of Emergency Medicine

Matthew Harmody, MD

Medical Director, FirstHealth EMS and CCT
Medical Director, Moore Regional Hospital
Medical Director Moore Regional Hospital-Hoke Campus EDs

Mark Quale, MD

Medical Director Alamance County EMS
Clinical Affiliate
Wake Forest University
Department of Emergency Medicine

Henderson McGinnis, MD

Medical Director Wilkes County EMS
Medical Director Wake Forest Baptist AirCare CCT
Associate Professor
Wake Forest University
Department of Emergency Medicine

Jose Cabanos, MD

Director Wake County EMS
Medical Director Wake County EMS
Adjunct Associate Professor
UNC-Chapel Hill
Department of Emergency Medicine

Jane Brice, MD

Professor and Chair
UNC-Chapel Hill
Department of Emergency Medicine

Key to Protocol Utilization

History

- Important history items
- Circumstances of event
- SAMPLE
- Time of onset
- Duration

Signs and Symptoms

- Important Signs and Symptoms specific to each protocol

Differential

- A list of other disease or injury which should be considered

Black Box

Highlights Important Information



Universal Patient Care Protocol
Assumed all protocols utilize and will not appear on individual protocols

Red Box

Highlights Critical Information

May direct to another protocol



Signals protocol within a protocol

Information box

Indicates Entry / Exit from / to another protocol(s)



Decision Point
 Darker outline to highlight



Highlights medication after Contact Medical Control
 May be added by Local Medical Director

Purple Shading of Information Box

Indicates items changeable at local agency level, including medications / dosages on NCMB formulary
 Local Medical Director may add / change at his / her discretion
 Local medical director may add page 2 to any protocol where none exists for additional comments

Algorithm Legend


	Emergency Medical Responder
B	Emergency Medical Technician
A	Advanced Emergency Medical Technician
P	Paramedic
	Notify Destination or Contact Medical Control

Pearls

- Important information specific to each protocol will appear here.
- Will usually appear on page.
- Important exam items listed here specific to protocol.



Universal Patient Care



Universal Protocol Section



Universal Patient Care

- **Pearls**

- **Recommended Exam: Minimal exam if not noted on the specific protocol is vital signs, mental status with GCS, and location of injury or complaint.**

- **Any patient contact which does not result in an EMS transport must have a completed disposition form.**
- **Vital signs should be obtained before, 10 minutes after, and at patient hand off with all pain medications.**
- **2 complete vital sign acquisitions should occur at a minimum with a patient encounter.**

- **Patient Refusal**

Patient refusal is a high risk situation. Encourage patient to accept transport to medical facility.

Encourage patient to allow an assessment, including vital signs. Documentation of the event is very important including a mental status assessment describing the patient's capacity to refuse care.

Guide to Assessing capacity:

C: Patient should be able to communicate a clear choice: This should remain stable over time. Inability to communicate a choice or an inability to express the choice consistently demonstrates incapacity.

R: Relevant information is understood: Patient should be able to display a factual understanding of the illness, the options and risks and benefits.

A: Appreciation of the situation: Ability to communicate an understanding of the facts of the situation. They should be able to recognize the significance of the outcome potentially from their decision.

M: Manipulation of information in a rational manner: Demonstrate a rational process to come to a decision. Should be able to describe the logic they are using to come to the decision, though you may not agree with decision.

- **Pediatric Patient General Considerations:**

A pediatric patient is defined by fitting a Length-based Resuscitation Tape, Age ≤ 15 , weight ≤ 49 kg.

Patients off the Broselow-Luten tape should have weight based medications until age ≥ 16 or weight ≥ 50 kg.


Special needs children may require continued use of Pediatric based protocols regardless of age and weight. Initial assessment should utilize the Pediatric Assessment Triangle which encompasses Appearance, Work of Breathing and Circulation to skin.

The order of assessment may require alteration dependent on the developmental state of the pediatric patient. Generally the child or infant should not be separated from the caregiver unless absolutely necessary during assessment and treatment.

- Timing of transport should be based on patient's clinical condition and the transport policy.
- Never hesitate to contact medical control for patient who refuses transport.
- Blood Pressure is defined as a Systolic / Diastolic reading. A palpated Systolic reading may be necessary at times.
- SAMPLE: Signs / Symptoms; Allergies; Medications; PMH; Last oral intake; Events leading to illness / injury



Triage





Triage

Pearls

- **When approaching a multiple casualty incident where resources are limited:**
Triage decisions must be made rapidly with less time to gather information
Emphasis shifts from ensuring the best possible outcome for an individual patient to ensuring the best possible outcome for the greatest number of patients.
- **Scene Size Up:**
 1. **Conduct a scene size up. Assure well being of responders. Determine or ensure scene safety before entering. If there are several patients with the same complaints consider HazMat, WMC or CO poisoning.**
 2. **Take Triage system kit.**
 3. **Determine number of patients. Communicate the number of patients and nature of the incident, establish command and establish a medical officer and triage officer if personnel available**
- **Triage is a continual process and should recur in each section as resources allow.**
- **Step 1: Global sorting:**
Call out to those involved in the incident to walk to a designated area and assess third.
For those who cannot walk, have them wave / indicate a purposeful movement and assess them second.
Those involved who are not moving or have an obvious life threat, assess first.
- **Step 2: Individual assessments:**
Control major hemorrhage
Open airway and if child, give 2 rescue breaths
Perform Needle Chest Decompression Procedure if indicated.
Administer injector antidotes if indicated
- **Assess the first patient you encounter using the three objective criteria which can be remembered by RPM.**
R: Respiratory
P: Perfusion
M: Mental Status
- If your patient falls into the RED TAG category, stop, place RED TAG and move on to next patient. Attempt only to correct airway problems, treat uncontrolled bleeding, or administer an antidote before moving to next patient.
- **Treatment:**
Once casualties are triaged focus on treatment can begin. You may need to move patients to treatment areas.
RED TAGs are moved / treated first followed by YELLOW TAGs. BLACK TAGs should remain in place.
You may also indicate deceased patients by pulling their shirt / clothing over their head.
As more help arrives then the triage / treatment process may proceed simultaneously.
- Capillary refill can be altered by many factors including skin temperature. Age-appropriate heart rate may also be used in triage decisions.
- SMART triage tag system is utilized in NC.



Abdominal Pain Vomiting and Diarrhea

History


- Age
- Time of last meal
- Last bowel movement/emesis
- Improvement or worsening with food or activity
- Duration of problem
- Other sick contacts
- Past medical history
- Past surgical history
- Medications
- Menstrual history (pregnancy)
- Travel history
- Bloody emesis / diarrhea

Signs and Symptoms

- Pain
 - Character of pain (constant, intermittent, sharp, dull, etc.)
 - Distention
 - Constipation
 - Diarrhea
 - Anorexia
 - Radiation
- Associated symptoms:**
Fever, headache, blurred vision, weakness, malaise, myalgias, cough, headache, dysuria, mental status changes, rash

Differential

- CNS (increased pressure, headache, stroke, CNS lesions, trauma or hemorrhage, vestibular)
- Myocardial infarction
- Drugs (NSAID's, antibiotics, narcotics, chemotherapy)
- GI or Renal disorders
- Diabetic ketoacidosis
- OB-Gyn disease (ovarian cyst, PID, Pregnancy)
- Infections (pneumonia, influenza)
- Electrolyte abnormalities
- Food or toxin induced
- Medication or Substance abuse
- Psychological



Universal Protocol Section



Abdominal Pain Vomiting and Diarrhea

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Age specific blood pressure 0 – 28 days > 60 mmHg, 1 month - 1 year > 70 mmHg, 1 - 10 years > 70 + (2 x age) mmHg and 11 years and older > 90 mmHg.**
- **Abdominal / back pain in women of childbearing age should be treated as pregnancy related until proven otherwise.**
- **The diagnosis of abdominal aneurysm should be considered with abdominal pain, with or without back and / or lower extremity pain or diminished pulses, especially in patients over 50 and / or patients with shock/ poor perfusion. Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients > 50, diabetics and / or women especially with upper abdominal complaints.**
- **Repeat vital signs after each fluid bolus.**
- **Heart Rate: One of the first clinical signs of dehydration, almost always increased heart rate, tachycardia increases as dehydration becomes more severe, very unlikely to be significantly dehydrated if heart rate is close to normal.**
- **Promethazine (Phenergan) may cause sedative effects in pediatric patients and ages ≥ 60 and the debilitated, etc.) When giving promethazine IV dilute with 10 mL of normal saline and administer slowly as it can also harm the veins.**
- Beware of vomiting only in children. Pyloric stenosis, bowel obstruction, and CNS processes (bleeding, tumors, or increased CSF pressures) all often present with vomiting.
- Document the mental status and vital signs prior to administration of Promethazine (Phenergan).
- Isolated vomiting may be caused by pyloric stenosis, bowel obstruction, and CNS processes (bleeding, tumors, or increased CSF pressures).
- Vomiting and diarrhea are common symptoms, but can be the symptoms of uncommon and serious pathology such as stroke, carbon monoxide poisoning, acute MI, new onset diabetes, diabetic ketoacidosis (DKA), and organophosphate poisoning. Maintain a high index of suspicion.



Altered Mental Status

History

- Known diabetic, medic alert tag
- Drugs, drug paraphernalia
- Report of illicit drug use or toxic ingestion
- Past medical history
- Medications
- History of trauma
- Change in condition
- Changes in feeding or sleep habits


Signs and Symptoms

- Decreased mental status or lethargy
- Change in baseline mental status
- Bizarre behavior
- Hypoglycemia (cool, diaphoretic skin)
- Hyperglycemia (warm, dry skin; fruity breath; Kussmaul respirations; signs of dehydration)
- Irritability

Differential

- Head trauma
- CNS (stroke, tumor, seizure, infection)
- Cardiac (MI, CHF)
- Hypothermia
- Infection (CNS and other)
- Thyroid (hyper / hypo)
- Shock (septic, metabolic, traumatic)
- Diabetes (hyper / hypoglycemia)
- Toxicological or Ingestion
- Acidosis / Alkalosis
- Environmental exposure
- Pulmonary (Hypoxia)
- Electrolyte abnormality
- Psychiatric disorder

Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 <i>if indicated</i>	
	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure
A	IV / IO Procedure
Age Appropriate Diabetic Protocol(s) AM 2 / PM 2 <i>if indicated</i>	



Universal Protocol Section



Altered Mental Status

Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.**
- **AMS may present as a sign of an environmental toxin or Haz-Mat exposure - protect personal safety.**
- **General:**
 - The patient with AMS poses one of the most significant challenges.**
 - A careful assessment of the patient, the scene and the circumstances should be undertaken.**
 - Assume the patient has a life threatening cause of their AMS until proven otherwise.**
 - Pay careful attention to the head exam for signs of bruising or other injury.**
 - Information found at the scene must be communicated to the receiving facility.**
- **Substance misuse:**
 - Patients ingesting substances can pose a great challenge.
 - DO NOT assume recreational drug use and / or alcohol are the sole reasons for AMS.
 - Misuse of alcohol may lead to hypoglycemia.
 - More serious underlying medical and trauma conditions may be the cause.
- **Behavioral health:**
 - The behavioral health patient may present a great challenge in forming a differential.
 - DO NOT assume AMS is the result solely of an underlying psychiatric etiology.
 - Often an underlying medical or trauma condition precipitates a deterioration of a patient's underlying disease.
- **Spinal Motion Restriction / Trauma:**
 - Only utilize spinal immobilization if the situation warrants.
 - The patient with AMS may worsen with increased agitation when immobilized.
- **It is safer to assume hypoglycemia than hyperglycemia if doubt exists. Recheck blood glucose after Dextrose or Glucagon**
- Consider Restraints if necessary for patient's and/or personnel's protection per the restraint procedure.



Back Pain

History


- Age
- Past medical history
- Past surgical history
- Medications
- Onset of pain / injury
- Previous back injury
- Traumatic mechanism
- Location of pain
- Fever
- Improvement or worsening with activity

Signs and Symptoms

- Pain (paraspinous, spinous process)
- Swelling
- Pain with range of motion
- Extremity weakness
- Extremity numbness
- Shooting pain into an extremity
- Bowel / bladder dysfunction

Differential

- Muscle spasm / strain
- Herniated disc with nerve compression
- Sciatica
- Spine fracture
- Kidney stone
- Pyelonephritis
- Aneurysm
- Pneumonia
- Spinal Epidural Abscess
- Metastatic Cancer
- AAA





Back Pain

Pearls

- **Recommended Exam: Mental Status, Heart, Lungs, Abdomen, Neuro, Lower extremity perfusion**
- **Back pain is one of the most common complaints in medicine and affects more than 90 % of adults at some point in their life. Back pain is also common in the pediatric population. Most often it is a benign process but in some circumstances can be life or limb threatening.**
- **Consider pregnancy or ectopic pregnancy with abdominal or back pain in women of childbearing age.**
- **Consider abdominal aortic aneurysm with abdominal pain especially in patients over 50 and/or patients with shock/ poor perfusion. Patients may have abdominal pain and / or lower extremity pain with diminished pulses, . Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients > 50, diabetics and / or women especially with upper abdominal complaints.**
- **Red Flags which may signal more serious process associated with back pain:**
 - Age > 50 or < 18
 - Neurological deficit (leg weakness, urinary retention, or bowel incontinence)
 - IV Drug use
 - Fever
 - History of cancer, either current or remote
 - Night time pain in pediatric patients
- **Cauda equina syndrome is where the terminal nerves of spinal cord are being compressed (Symptoms include):**
 - Saddle anesthesia
 - Recent onset of bladder and bowel dysfunction. (Urine retention and bowel incontinence)
 - Severe or progressive neurological deficit in the lower extremity.
 - Motor weakness of thigh muscles or foot drop
- **Back pain associated with infection:**
 - Fever / chills.
 - IV Drug user (consider spinal epidural abscess)
 - Recent bacterial infection like pneumonia.
 - Immune suppression such as HIV or patients on chronic steroids like prednisone.
 - Meningitis.
- **Spinal motion restriction in patients with underlying spinal deformity should be maintained in their functional position.**
- **Kidney stones typically present with an acute onset of flank pain which radiates around to the groin area.**



IV or IO Access

History


- Chronic medical conditions requiring recurrent need for IV access for medication, hydration, or blood sampling.
- Medical condition requiring administration of IV medications at home.
- End-stage renal disease requiring hemodialysis.
- Chronic medical condition requiring IV nutrition.

Signs and Symptoms

- Fever
- Bleeding
- Hypotension
- Redness, swelling, and/or pain at IV catheter site
- Shortness of breath
- Chest pain
- IV catheter patency

Differential

- Infection or sepsis
- Infection of catheter
- Clotted IV catheter
- Air embolism
- Pneumothorax
- Overdose of home medication
- Shock







Universal Protocol Section

Pearls

- **Frequent encounter of patients with IV access devices and confusion as to which device can be accessed and used by EMS providers.**
- **If unclear about device use, always ask "Is this device used for dialysis?"**
- **When accessing central catheter, always ensure sterility of catheter connection point by cleaning port with alcohol, or similar disinfectant, 2 – 3 times prior to access.**
- **Central line catheters placed for administration of chemotherapy, medications, electrolytes, antibiotics, and blood are available to EMS providers for access and administration of fluids, medications, antibiotics, and blood products.**
- **Central line catheters placed for hemodialysis are NOT available for access by EMS providers unless the patient is in cardiac arrest.**
- Long term IV access is frequently needed for a variety of indications:
 - Medication administration such as antibiotics, pain relief, or chemotherapy
 - Administration of IV nutrition or feeding
 - Need for multiple IV line access or recurrent blood sampling
 - Poor vasculature requiring repeated attempts at IV access
 - End-stage renal disease requiring hemodialysis
- Common complications of central access devices:
 - Infection
 - Damage to vasculature
 - Air embolism
 - Loss of patency due to clogging or clotting
 - Pneumothorax

IV or IO Access





Dental Problems

History


- Age
- Past medical history
- Medications
- Onset of pain / injury
- Trauma with "knocked out" tooth
- Location of tooth
- Whole vs. partial tooth injury

Signs and Symptoms

- Bleeding
- Pain
- Fever
- Swelling
- Tooth missing or fractured

Differential

- Decay
- Infection
- Fracture
- Avulsion
- Abscess
- Facial cellulitis
- Impacted tooth (wisdom)
- TMJ syndrome
- Myocardial infarction



Pearls

- **Recommended Exam: Mental Status, HEENT, Neck, Chest, Lungs, Neuro**
- Significant soft tissue swelling to the face or oral cavity can represent a cellulitis or abscess.
- Scene and transport times should be minimized in complete tooth avulsions. Reimplantation is possible within 4 hours if the tooth is properly cared for.
- Occasionally cardiac chest pain can radiate to the jaw.
- All pain associated with teeth should be associated with a tooth which is tender to tapping or touch (or sensitivity to cold or hot).



Emergencies Involving Indwelling Central Lines

History


- Central Venous Catheter Type
Tunneled Catheter (Broviac / Hickman)
- PICC (peripherally inserted central catheter)
- Implanted catheter (Mediport / Hickman)
- Occlusion of line
- Complete or partial dislodge
- Complete or partial disruption

Signs and Symptoms

- External catheter dislodgement
- Complete catheter dislodgement
- Damaged catheter
- Bleeding at catheter site
- Internal bleeding
- Blood clot
- Air embolus
- Erythema, warmth or drainage about catheter site indicating infection

Differential

- Fever
- Hemorrhage
- Reactions from home nutrient or medication
- Respiratory distress
- Shock



Universal Protocol Section

Pearls

- **Always talk to family / caregivers as they have specific knowledge and skills.**
- **Use strict sterile technique when accessing / manipulating an indwelling catheter.**
- **Cardiac arrest: May access central catheter and utilize if functioning properly.**
- Do not attempt to force catheter open if occlusion evident.
- Some infusions may be detrimental to stop. Ask family or caregiver if it is appropriate to stop or change infusion.
- Hyperalimentation infusions (IV nutrition): If stopped for any reason monitor for hypoglycemia.



Epistaxis

History


- Age
- Past medical history
- Medications (HTN, anticoagulants, aspirin, NSAIDs)
- Previous episodes of epistaxis
- Trauma
- Duration of bleeding
- Quantity of bleeding

Signs and Symptoms

- Bleeding from nasal passage
- Pain
- Nausea
- Vomiting

Differential

- Trauma
- Infection (viral URI or Sinusitis)
- Allergic rhinitis
- Lesions (polyps, ulcers)
- Hypertension



Universal Protocol Section

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Neuro**
- **Age specific hypotension: 0 – 28 days < 60 mmHg, 1 month – 1 year < 70 mmHg, 1 year – 10 years < 70 + (2 x age)mmHg, 11 years and greater < 90 mmHg.**
- It is very difficult to quantify the amount of blood loss with epistaxis.
- Bleeding may also be occurring posteriorly. Evaluate for posterior blood loss by examining the posterior pharynx.
- Anticoagulants include warfarin (Coumadin), Apixaban (Eliquis), heparin, enoxaparin (Lovenox), dabigatran (Pradaxa), rivaroxaban (Xarelto), and many over the counter headache relief powders.
- Anti-platelet agents like aspirin, clopidogrel (Plavix), aspirin/dipyridamole (Aggrenox), and ticlopidine (Ticlid) can contribute to bleeding.



Fever / Infection Control

History

- Age
- Duration of fever
- Severity of fever
- Past medical history
- Medications
- Immunocompromised (transplant, HIV, diabetes, cancer)
- Environmental exposure
- Last acetaminophen or ibuprofen

Signs and Symptoms


- Warm
- Flushed
- Sweaty
- Chills/Rigors

Associated Symptoms (Helpful to localize source)

- myalgias, cough, chest pain, headache, dysuria, abdominal pain, mental status changes, rash

Differential

- Infections / Sepsis
- Cancer / Tumors / Lymphomas
- Medication or drug reaction
- Connective tissue disease
 - Arthritis
 - Vasculitis
- Hyperthyroidism
- Heat Stroke
- Meningitis



Universal Protocol Section

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- Febrile seizures are more likely in children with a history of febrile seizures and with a rapid elevation in temperature.
- **Droplet precautions:**
 - Include standard PPE plus a standard surgical mask for providers who accompany patients in the back of the ambulance and a surgical mask or NRB O2 mask for the patient.
 - This level of precaution should be utilized when influenza, meningitis, mumps, streptococcal pharyngitis, and other illnesses spread via large particle droplets are suspected.
 - A patient with a potentially infectious rash should be treated with droplet precautions.
- **Airborne precautions:**
 - Include standard PPE plus utilization of a gown, change of gloves after every patient contact, and strict hand washing precautions.
 - This level of precaution is utilized when multi-drug resistant organisms (e.g. MRSA), scabies, or zoster (shingles), or other illnesses spread by contact are suspected.
- **All-hazards precautions:**
 - Include standard PPE plus airborne precautions plus contact precautions.
 - This level of precaution is utilized during the initial phases of an outbreak when the etiology of the infection is unknown or when the causative agent is found to be highly contagious (e.g. SARS).
- All patients should have drug allergies documented prior to administering pain medications.
- Allergies to NSAIDs (non-steroidal anti-inflammatory medications) are a contraindication to Ibuprofen. Do not give to patients who have renal disease or renal transplant.
- NSAIDs should not be used in the setting of environmental heat emergencies.
- **Do not** give aspirin to a child, age ≤ 15 years.



Pain Control

History

- Age
- Location
- Duration
- Severity (1 - 10)
- If child use Wong-Baker faces scale
- Past medical history
- Medications
- Drug allergies


Signs and Symptoms

- Severity (pain scale)
- Quality (sharp, dull, etc.)
- Radiation
- Relation to movement, respiration
- Increased with palpation of area

Differential

- Per the specific protocol
- Musculoskeletal
- Visceral (abdominal)
- Cardiac
- Pleural / Respiratory
- Neurogenic
- Renal (colic)

Specific Complaint Protocol



Universal Protocol Section



Pain Control

Pearls

- **Recommended Exam: Mental Status, Area of Pain, Neuro**
- **Pain severity (0-10) is a vital sign to be recorded before and after PO, IV, IO or IM medication delivery and at patient hand off. Monitor BP closely as sedative and pain control agents may cause hypotension.**
- **Ketamine:**
Ketamine may be used in patients who are outside a Pediatric Medication/Skill Resuscitation System product. Ketamine may be used in patients who fit within a Pediatric Medication/Skill Resuscitation System product only with DIRECT ONLINE MEDICAL ORDER, by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR.
- **Ketamine: appropriate indications for pain control:**
Patients who have developed opioid-tolerance. Sickle cell crisis patients with opioid-tolerance.
Patients who have obstructive sleep apnea.
May use in combination with opioids to limit total amount of opioid administration.
- **Ketamine: caution when using for pain control:**
Slow infusion or IV push over 10 minutes is associated with less side effects. Do not administer by rapid IV push.
Avoid in patients who have cardiac disease or uncontrolled hypertension.
Avoid in patients with increased intraocular pressure such as glaucoma.
Avoid use in combination with benzodiazepines due to decreased respiratory effort.
- **Both arms of the treatment may be used in concert. For patients in Moderate pain for instance, you may use the combination of an oral medication and parenteral if no contraindications are present.**
- **Pediatrics:**
For children use Wong-Baker faces scale or the FLACC score (see Assessment Pain Procedure)
Use Numeric (> 9 yrs), Wong-Baker faces (4-16yrs) or FLACC scale (0-7 yrs) as needed to assess pain
- **Vital signs should be obtained before, 10 minutes after, and at patient hand off with all pain medications.**
- All patients who receive IM or IV medications must be observed 15 minutes for drug reaction in the event no transport occurs.
- Do not administer **Acetaminophen** to patients with a history of liver disease.
- Burn patients may required higher than usual opioid doses to titrate adequate pain control.
- Consider agency-specific anti-emetic(s) for nausea and/or vomiting.



Police Custody

History


- Traumatic Injury
- Drug Abuse
- Cardiac History
- History of Asthma
- Psychiatric History

Signs and Symptoms

- External signs of trauma
- Palpitations
- Shortness of breath
- Wheezing
- Altered Mental Status
- Intoxication/Substance Abuse

Differential

- Agitated Delirium Secondary to Psychiatric Illness
- Agitated Delirium Secondary to Substance Abuse
- Traumatic Injury
- Closed Head Injury
- Asthma Exacerbation
- Cardiac Dysrhythmia





Police Custody

Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Neurologic status**
- **Patient does not have to be in police custody or under arrest to utilize this protocol.**
- **EMS agency should formulate a policy with local law enforcement agencies concerning patients requiring EMS and Law Enforcement involvement simultaneously.**
- **Agencies should work together to formulate a disposition in the best interest of the patient.**
- **Law Enforcement:**
 - **Any patient who is handcuffed or restrained by Law Enforcement and transported by EMS, must be accompanied by law enforcement during transport capable of removing the devices.**
 - **Patient should not be transported with upper extremities hand-cuffed behind back as this prevents proper assessment and could lead to injury.**
 - **Consider multidisciplinary coordination with law enforcement to approach verbal de-escalation, restraint, and/or take-down restraint procedure.**
- **Maintain high-index of suspicion for underlying medical or traumatic disorder causing or contributing to behavioral disturbance. Medical causes more likely in ages < 12 or > 40.**
- **Medications are not to be used solely to aid in placing an individual into police custody. Physical and/or chemical restraints are reserved for a medical emergency in order to prevent imminent injury to a patient and/or providers.**
- **Restraints:**
 - **All patients who receive either physical or chemical restraint must be continuously observed by ALS personnel on scene or immediately upon their arrival.**
 - **Do not position or transport any restrained patient in such a way that could impact the patient's respiratory or circulatory status.**
 - **However, when EMS providers have utilized physical restraints in accordance with Restraint Procedure USP 5, the law enforcement agent may follow behind the ambulance during transport.**
- The responsibility for patient care rests with the highest authorized medical provider on scene per North Carolina law.
- If an asthmatic patient is exposed to pepper spray and released to law enforcement, all parties should be advised to immediately contact EMS if wheezing/difficulty breathing occurs.
- Patients exposed to chemical spray, with or without history of respiratory disease, may develop respiratory complaints up to 20 minutes post exposure.
- All patients with decision-making capacity in police custody retain the right to participate in decision making regarding their care and may request care or refuse care of EMS.
- If extremity / chemical / law enforcement restraints are applied, follow Restraint Procedure.
- **Excited Delirium Syndrome and Violent:**
 - **Medical emergency: Combination of delirium, psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent / bizarre behavior, insensitivity to pain, hyperthermia and increased strength.**
 - **Potentially life-threatening and associated with use of physical control measures, including physical restraints.**
 - **Most commonly seen in male subjects with a history of serious mental illness and/or acute or chronic drug abuse, particularly stimulant drugs such as cocaine, crack cocaine, methamphetamine, amphetamines or similar agents.**
 - **Alcohol or substance withdrawal as well as head trauma may also contribute to the condition.**
 - **If patient suspected of EDS suffers cardiac arrest, consider a fluid bolus and sodium bicarbonate early.**



Seizure

History

- Reported / witnessed seizure activity
- Previous seizure history
- Medical alert tag information
- Seizure medications
- History of trauma
- History of diabetes
- History of pregnancy
- Time of seizure onset
- Document number of seizures
- Alcohol use, abuse or abrupt cessation
- Fever

Signs and Symptoms

- Decreased mental status
- Sleepiness
- Incontinence
- Observed seizure activity
- Evidence of trauma
- Unconscious

Differential

- CNS (Head) trauma
- Tumor
- Metabolic, Hepatic, or Renal failure
- Hypoxia
- Electrolyte abnormality (Na, Ca, Mg)
- Drugs, Medications, Non-compliance
- Infection / Fever
- Alcohol withdrawal
- Eclampsia
- Stroke
- Hyperthermia
- Hypoglycemia

Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 <i>as indicated</i>
Altered Mental Status Protocol UP 4 <i>if indicated</i>
Childbirth/Labor Protocol AO 1 Obstetrical Emergency Protocol AO 3 <i>if indicated</i>
Behavioral Protocol UP 17, 18, 19 <i>if indicated</i>
Loosen any constrictive clothing Protect patient

Active Seizure Activity

IV / IO Access

P	
---	--

P	
---	--

	Blood Glucose Analysis Procedure
A	IV / IO Procedure <i>if indicated</i>
P	Cardiac Monitor <i>if indicated</i>
	Monitor and Reassess

	Notify Destination or Contact Medical Control	
--	--	--

Universal Protocol Section



Seizure

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Extremities, Neuro**
- **Items in Red Text are key performance measures used to evaluate protocol compliance and care.**
- **Brief seizure-like activity can be seen following ventricular fibrillation or ventricular tachycardia associated cardiac arrest.**
- **Status epilepticus is defined by seizure activity lasting > 5 minutes or multiple seizures without return to baseline.**
- **Most seizure activity is brief, lasting only 1 – 2 minutes, and is associated with transient hypoventilation.**
- **Be prepared for airway problems and continued seizures.**
- **Seizure activity may be a marker of closed head injury, especially in the very young, examine for trauma.**
- **Adult:**
 - **Midazolam 10 mg IM is effective in termination of seizures.**
 - **Do not delay IM administration with difficult IV or IO access. IM Preferred over IO.**
- **Pediatrics:**
 - **Midazolam 0.2 mg/kg (Maximum 5 mg) IM is effective in termination of seizures.**
 - **Do not delay IM administration with difficult IV or IO access. IM Preferred over IO.**
- **Do not delay administration of anti-epileptic drugs to check for blood glucose.**
- **Grand mal seizures (generalized)** are associated with loss of consciousness, incontinence, and tongue trauma.
- **Focal seizures** affect only a part of the body and are not usually associated with a loss of consciousness, but can propagate to generalized seizures with loss of consciousness.
- **Be prepared to assist ventilations especially if diazepam or midazolam is used.**
- **For any seizure in a pregnant patient, follow the OB Emergencies Protocol.**
- **Diazepam (Valium) is not effective when administered IM. Give IV.**
- **Optimal conditions for patients refusing transport following a seizure:**

Known history of seizures/epilepsy	Seizure not associated with drugs or alcohol
Full recovery to baseline mental status	Only 1 seizure episode in the past hour
No injuries requiring treatment or evaluation	Seizure not associated with pregnancy
Adequate supervision	

Suspected Stroke

History


- Previous CVA, TIA's
- Previous cardiac / vascular surgery
- Associated diseases: diabetes, hypertension, CAD
- Atrial fibrillation
- Medications (blood thinners)
- History of trauma
- Sickle Cell Disease
- Immune disorders
- Congenital heart defects
- Maternal infection / hypertension

Signs and Symptoms

- Altered mental status
- Weakness / Paralysis
- Blindness or other sensory loss
- Aphasia / Dysarthria
- Syncope
- Vertigo / Dizziness
- Vomiting
- Headache
- Seizures
- Respiratory pattern change
- Hypertension / hypotension

Differential

- See Altered Mental Status
- TIA (Transient ischemic attack)
- Seizure
- Todd's Paralysis
- Hypoglycemia
- Stroke
 - Thrombotic or Embolic (~85%)
 - Hemorrhagic (~15%)
- Tumor
- Trauma
- Dialysis / Renal Failure





Suspected Stroke

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Neuro**
- **Items in Red Text are key performance measures used in the EMS Acute Stroke Care Toolkit.**
- **Acute Stroke care is evolving rapidly. Time of onset / last seen normal may be changed at any time depending on the capabilities and resources of your hospital based on Stroke: EMS Triage and Destination Plan.**
- **Time of Onset or Last Seen Normal:**
 - **One of the most important items the pre-hospital provider can obtain, of which all treatment decisions are based.**
 - **Be very precise in gathering data to establish the time of onset and report as an actual time (i.e. 13:47 NOT “about 45 minutes ago.”)**
 - **Without this information patient may not be able to receive thrombolytics at facility.**
 - **Wake up stroke: Time starts when patient last awake or symptom free.**
- **You are often in the best position to determine the actual Time of Onset while you have family, friends or caretakers available. Often these sources of information may arrive well after you have delivered the patient to the hospital. Delays in decisions due to lack of information may prevent an eligible patient from receiving thrombolytics.**
- **The Reperfusion Checklist should be completed for any suspected stroke patient. With a duration of symptoms of less than , scene times should be limited to ≤ 10 minutes, early notification / activation of receiving facility should be performed and transport times should be minimized.**
- **If possible place 2 IV sites.**
- **Blood Draw:**
 - **Many systems utilize EMS venous blood samples. Follow your local policy and procedures.**
- **The differential listed on the Altered Mental Status Protocol should also be considered.**
- **Be alert for airway problems (swallowing difficulty, vomiting/aspiration).**
- **Hypoglycemia can present as a localized neurologic deficit, especially in the elderly.**
- **Document the Stroke Screen results in the PCR.**
- **Agencies may use validated pre-hospital stroke screen of choice.**
- **Pediatrics:**
 - **Strokes do occur in children, they are slightly more common in ages < 2 , in boys, and in African-Americans. Newborn and infant symptoms consist of seizures, extreme sleepiness, and using only one side of the body. Children and teenagers symptoms may consist of severe headaches, vomiting, sleepiness, dizziness, and/or loss of balance or coordination.**



Suspected Sepsis

History

- Duration and severity of fever
- Past medical history
- Medications / Recent antibiotics
- Immunocompromised (transplant, HIV, diabetes, cancer)
- Indwelling medical device
- Last acetaminophen or ibuprofen
- Recent Hospital / healthcare facility
- Bedridden or immobile
- Elderly and very young – at risk
- Prosthetic device / indwelling device

Signs and Symptoms

- Warm
- Flushed
- Sweaty
- Chills / Rigors
- Delayed cap refill
- Mental status changes

Associated Symptoms (Helpful to localize source)

- myalgias, cough, chest pain, headache, dysuria, abdominal pain, rash

Differential

- Infections: UTI, Pneumonia, skin/ wound
- Cancer / Tumors / Lymphomas
- Medication or drug reaction
- Connective tissue disease: Arthritis, Vasculitis
- Hyperthyroidism
- Heat Stroke
- Meningitis
- Hypoglycemia/hypothermia
- MI / CVA

Consider: Contact, Droplet, and Airborne Precautions	
Temperature Measurement Procedure <i>if available</i>	
	Fever / Infection Control Protocol UP 10 <i>if needed</i>
	Altered Mental Status Protocol UP 4 <i>if needed</i>
B	12 Lead ECG Procedure
A	IV / IO Procedure <i>if indicated</i>
P	Cardiac Monitor

Exit to
Age Appropriate
Condition Appropriate
Protocol(s)

Sepsis Screen
Positive


SEPSIS ALERT
Notify Receiving Facility
Immediately

Venous Access Blood Draw <i>if applicable</i>	
A	
P	
	Age Appropriate Hypotension / Shock Protocol AM 5 / PM 3

MAP
(Mean Arterial Pressure)

$SBP + 2(DBP)$
3

Monitor usually calculates this
value on screen



Notify Destination or
Contact Medical Control

Universal Protocol Section



Suspected Sepsis

• Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Recommended Exam Pediatrics: In childhood, physical assessment reveals important clues for sepsis. Look for mental status abnormalities such as anxiety, restlessness, agitation, irritability, confusion, or lethargy. Cardiovascular findings to look for include cool extremities, capillary refill >3 seconds, or mottled skin.**
- **Sepsis is a life threatening condition where the body's immune response to infection injures its own tissues and organs.**
- **Severe sepsis is a suspected infection and 2 or more SIRS criteria (or qSOFA) with organ dysfunction such as AMS or hypotension.**
- **Septic shock is severe sepsis and poor perfusion unimproved after fluid bolus.**
- **Agencies administering antibiotics should inquire about drug allergies specific to antibiotics or family of antibiotics.**
- **Following each fluid bolus, assess for pulmonary edema. Consider administration of agency specific vasopressor.**
- **Supplemental oxygen should be given and titrated to oxygenation saturation $\geq 94\%$.**
- **EKG should be obtained with suspected sepsis, but should not delay care in order to obtain.**
- **Abnormally low temperatures increase mortality and found often in geriatric patients.**
- Quantitative waveform capnography can be a reliable surrogate for lactate monitoring in detecting metabolic distress in sepsis patients. EtCO₂ < 25 mm Hg are associated with serum lactate levels > 4 mmol/L.
- Patients with a history of liver failure should not receive acetaminophen.
- **Droplet precautions:**
 - Include standard PPE plus a standard surgical mask for providers who accompany patients in the back of the ambulance and a surgical mask or NRB O2 mask for the patient.
 - This level of precaution should be utilized when influenza, meningitis, mumps, streptococcal pharyngitis, and other illnesses spread via large particle droplets are suspected.
 - A patient with a potentially infectious rash should be treated with droplet precautions.
- **Airborne precautions:**
 - Include standard PPE plus utilization of a gown, change of gloves after every patient contact, and strict hand washing precautions.
 - This level of precaution is utilized when multi-drug resistant organisms (e.g. MRSA), scabies, or zoster (shingles), or other illnesses spread by contact are suspected.
- **All-hazards precautions:**
 - Include standard PPE plus airborne precautions plus contact precautions.
 - This level of precaution is utilized during the initial phases of an outbreak when the etiology of the infection is unknown or when the causative agent is found to be highly contagious (e.g. SARS).
- All patients should have drug allergies documented prior to administering pain medications.
- Allergies to NSAIDs (non-steroidal anti-inflammatory medications) are a contraindication to Ibuprofen.
- Agency Medical Director may require contact of medical control prior to EMT / MR administering any medication.
- **Sepsis Screen:**
 - Agencies may use Adult / Pediatric Systemic Inflammatory Response Syndrome (SIRS) criteria or quickSOFA (qSOFA) criteria.
 - Receiving facility should be involved in determining Sepsis Screen utilized by EMS.



Syncope

History

- Cardiac history, stroke, seizure
- Occult blood loss (GI, ectopic)
- Females: LMP, vaginal bleeding
- Fluid loss: nausea, vomiting, diarrhea
- Past medical history
- Medications

Signs and Symptoms

- Loss of consciousness with recovery
- Lightheadedness, dizziness
- Palpitations, slow or rapid pulse
- Pulse irregularity
- Decreased blood pressure

Differential

- Vasovagal
- Orthostatic hypotension
- Cardiac syncope
- Micturition / Defecation syncope
- Psychiatric
- Stroke
- Hypoglycemia
- Seizure
- Shock (see Shock Protocol)
- Toxicological (Alcohol)
- Medication effect (hypertension)
- PE
- AAA

Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 if indicated	
	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure
I	IV / IO Procedure
P	Cardiac Monitor
Altered Mental Status Protocol UP 4 if indicated	
Age Appropriate Cardiac Protocol(s) if indicated	
Age Appropriate Hypotension / Shock Protocol AM 5 / PM 3 if indicated	
Multiple Trauma Protocol TB 6 Spinal Motion Restriction Procedure / Protocol TB 8 if indicated	

Serious Signs / Symptoms
Hypotension, poor
perfusion, shock

YES

NO



**Notify Destination or
Contact Medical Control**



A	IV / IO Procedure Consider 2 Large Bore sites

Exit to
Age Appropriate
Condition Appropriate
Protocol(s)



Syncope

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Syncope is both loss of consciousness and loss of postural tone. Symptoms preceding the event are important in determining etiology.**
- **Syncope often is due to a benign process but can be an indication of serious underlying disease in both the adult and pediatric patient.**
- **Often patients with syncope are found normal on EMS evaluation. In general patients experiencing syncope require cardiac monitoring and emergency department evaluation.**
- **Differential should remain wide and include:**

Cardiac arrhythmia	Neurological problem	Choking	Pulmonary embolism
Hemorrhage	Stroke	Respiratory	Hypo or Hyperglycemia
GI Hemorrhage	Seizure	Sepsis	
- **High-risk patients:**

Age \geq 60	Syncope with exertion
History of CHF	Syncope with chest pain
Abnormal ECG	Syncope with dyspnea
- **Age specific blood pressure 0 – 28 days $>$ 60 mmHg, 1 month - 1 year $>$ 70 mmHg, 1 - 10 years $>$ 70 + (2 x age) mmHg and 11 years and older $>$ 90 mmHg.**
- **Abdominal / back pain in women of childbearing age should be treated as pregnancy related until proven otherwise.**
- **The diagnosis of abdominal aneurysm should be considered with abdominal pain, with or without back and / or lower extremity pain or diminished pulses, especially in patients over 50 and / or patients with shock/ poor perfusion. Notify receiving facility early with suspected abdominal aneurysm.**
- **Consider cardiac etiology in patients $>$ 50, diabetics and / or women especially with upper abdominal complaints.**
- **Heart Rate: One of the first clinical signs of dehydration, almost always increased heart rate, tachycardia increases as dehydration becomes more severe, very unlikely to be significantly dehydrated if heart rate is close to normal.**
- **Syncope with no preceding symptoms or event may be associated with arrhythmia.**
- **Assess for signs and symptoms of trauma if associated or questionable fall with syncope.**
- **Consider dysrhythmias, GI bleed, ectopic pregnancy, and seizure as possible causes of syncope.**
- **These patients should be transported. Patients who experience syncope associated with headache, neck pain, chest pain, abdominal pain, back pain, dyspnea, or dyspnea on exertion need prompt medical evaluation.**
- **More than 25% of geriatric syncope is cardiac dysrhythmia based.**



Behavioral Health Crisis

History


- Situational crisis
- Psychiatric illness/medications
- Injury to self or threats to others
- Medic alert tag
- Substance abuse / overdose
- Diabetes

Signs and Symptoms

- Anxiety, agitation, confusion
- Affect change, hallucinations
- Delusional thoughts, bizarre behavior
- Combative violent
- Expression of suicidal / homicidal thoughts

Differential

- Altered Mental Status
- Alcohol Intoxication
- Toxin / Substance abuse
- Medication effect / overdose / withdrawal
- Depression
- Bipolar (manic-depressive)
- Schizophrenia
- Anxiety disorders





Behavioral Health Crisis


Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Neurologic status**
- **Crew / responders safety is the main priority. Call for assistance, stage, or withdraw from scene if necessary.**
- **Law Enforcement:**
 - **Any patient who is handcuffed or restrained by Law Enforcement and transported by EMS, must be accompanied by law enforcement during transport.**
 - **Patient should not be transported with upper extremities hand-cuffed behind back as this prevents proper assessment and could lead to injury.**
 - **Consider multidisciplinary coordination with law enforcement to approach verbal de-escalation, restraint, and/or take-down restraint procedure.**
- **Maintain high-index of suspicion for underlying medical or traumatic disorder causing or contributing to behavioral disturbance. Medical causes more likely in ages < 12 or > 40.**
- **General communications techniques**
 - **Ask Open-ended questions (questions that cannot be answered with a yes/no)**
"Tell me how we can help you?" "What caused you to call 911 today?"
 - **Active listening (stay engaged, be able to summarize patient's story, use your body language to convey listening)**
Eye contact, nodding your head, periodically repeating back part of patient's story
 - **Encouraging (remain positive, convey interest in patient's crisis)**
"Tell me more about that..."
 - **Clarifying questions (ask patient to rephrase or repeat if you don't understand)**
"I'm not sure I understand, can you...?"
 - **Emotional labeling (naming emotions patient is demonstrating, validating emotions)**
"You look upset." "You seem angry."
 - **Conversational pause (okay to allow a period of silence for patient to process information)**
- **Behavioral health disturbance incidents are increasing and commonly involve the following:**

Substance misuse	Psychosis
Depression / Anxiety / Stress Reactions / Bipolar	Schizophrenia or schizophrenia-like illness
- **Restraints:**
 - **All patients who receive either physical or chemical restraint must be continuously observed by ALS personnel on scene or immediately upon their arrival.**
 - **Do not position or transport any restrained patient in such a way that could impact the patient's respiratory or circulatory status.**
- **Maintain high-index of suspicion for medical, trauma, abuse, or neglect causes:**
 - Hypoglycemia, hyperglycemia, overdose, substance abuse, hypoxia, head injury, shock, sepsis, stroke, etc.
 - Domestic violence, child or geriatric abuse/neglect.
- **Extrapyramidal reactions:**
 - Condition causing involuntary muscle movements or spasms typically of the face, neck and upper extremities. May present with contorted neck and trunk with difficult motor movements. Typically an adverse reaction to antipsychotic drugs like Haloperidol and may occur with your administration. When recognized give **Diphenhydramine 50 mg IV / IO / IM / PO** in adults or **1 mg/kg IV / IO / IM / PO** in pediatrics.
- **May add page 3 to protocol for specific for local mental health and / or substance misuse resources or destinations.**



Behavioral CIT Paramedic (Optional)




Universal Protocol Section

Alternative Destinations / Crisis Providers For Centerpoint

County		
Resource Agency	Resource Agency	Resource Agency
Hours of Operation	Hours of Operation	Hours of Operation



Behavioral Agitation / Sedation Guide





Behavioral Excited Delirium Syndrome / Violent

Call for help
Call for additional resources
Stage prior to arrival
or
Withdraw from scene until safe

Behavioral Activity Rating Scale (BARS)

Behavioral Activity Rating Scale
BARS scoring: 7

P	
	IV / IO Protocol UP 6 Preferably 2 large bore
A	
	Restraint Physical Procedure USP 5 Monitor per restraint procedure <i>if indicated</i>
	Restraint Therapeutic Take Down Procedure USP 6 <i>if indicated</i>
	Hyperthermia Protocol
P	Cardiac Monitor

Notify Destination or Contact Medical Control

- General Guidance**
- Use of protocol is indicated when a medical emergency requires a combination of both physical restraint and chemical restraint in order to prevent imminent injury to a patient and/or providers.
 - Use of this protocol requires medical judgement and consultation with medical control where indicated.
 - Non-medical personnel requests or opinions should not be used as a factor when implementing this protocol.




Behavioral Excited Delirium Syndrome / Violent

Pearls

- **Ketamine:**
Ketamine may be used in patients who are outside a Pediatric Medication/Skill Resuscitation System product. Ketamine may be used in patients who fit within a Pediatric Medication/Skill Resuscitation System product only with DIRECT ONLINE MEDICAL ORDER, by the system MEDICAL DIRECTOR or ASSISTANT MEDICAL DIRECTOR.
- **Excited Delirium Syndrome and Violent:**
Medical emergency: Combination of delirium, psychomotor agitation, anxiety, hallucinations, speech disturbances, disorientation, violent / bizarre behavior, insensitivity to pain, hyperthermia and increased strength. Potentially life-threatening and associated with use of physical control measures, including physical restraints. Most commonly seen in male subjects with a history of serious mental illness and/or acute or chronic drug abuse, particularly stimulant drugs such as cocaine, crack cocaine, methamphetamine, amphetamines or similar agents.
Alcohol or substance withdrawal as well as head trauma may also contribute to the condition.
- **Restraint use:**
Physical restraints are not contraindicated in agitated or excited delirium, but you must use caution. Once sedated, prevent patient from continued struggle which can worsen metabolic condition. Prevent patient from assuming a prone position for prolonged period, move to supine position as quickly as possible.
Team approach for sedation and Restraint Therapeutic Take Down Procedure USP-6:
1 provider for each limb. 1 provider to lead restraint, maintain airway and control head. 1 Provider to administer medication.
Do not position prone with restraints as this can impede respiration and ventilation
- Hyperthermia: Assess for and treat hyperthermia.



Adult Airway



Protocols AR 1, 2, and 3 should be utilized together (even if agency is not using Drug Assisted Airway Protocol) as they contain useful information for airway management.

- Capnography Monitoring**
- End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.
 - EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.

Unable to Ventilate and Oxygenate ≥ 90% during or after one (1) or more unsuccessful intubation attempts . and/or Anatomy inconsistent with continued attempts. and/or Three (3) unsuccessful attempts by most experienced Paramedic/ AEMT.

Exit to Adult Failed Airway Protocol AR 2



Adult Airway

Pearls

- See Pearls section of protocols AR 2 and 3.
- For the purposes of this protocol a secure airway is when the patient is receiving appropriate oxygenation and ventilation.
- If an effective airway is being maintained by BVM with continuous pulse oximetry values of $\geq 90\%$, it is acceptable to continue with basic airway measures.
- Ventilation rate should be 8-10 per minute to maintain a EtCO₂ of 35-45. Avoid hyperventilation.
- **Anticipating the Difficult Airway and Airway Assessment:**
 - **Difficult BVM Ventilation (MOANS):** Mask seal difficulty (hair, secretions, trauma); Obese, obstruction, OB – 2d and 3d trimesters; Age ≥ 55 ; No teeth; Stiff lungs or neck
 - **Difficult Laryngoscopy (LEON):** Look externally for anatomical problems; Evaluate 3-3-2 (Mouth opening should equal 3 of patient's finger's width, mental area to neck should equal 3 of patient's finger's width, base of chin to thyroid prominence should equal 2 of patient's finger's width); Obese, obstruction, OB – 2d and 3d trimesters; Neck mobility limited.
 - **Difficulty BIAD (RODS):** Restricted mouth opening; Obese, obstruction, OB – 2d and 3d trimesters; Distorted or disrupted airway; Stiff lungs or neck
 - **Difficulty Cricothyrotomy / Surgical Airway (SMART):** Surgery scars; Mass or hematoma, Access or anatomical problems; Radiation treatment to face, neck, or chest; Tumor.
- **Capnography Monitoring (EtCO₂):**
 - Continuous Waveform or Quantitative Capnography and Pulse Oximetry are required for intubation verification and ongoing patient monitoring (Not validated and may prove impossible in the neonatal population - verification by two (2) other means is recommended in this population.)
 - Capnography verification and monitoring is required for BIAD verification and monitoring once available on scene.
- **Nasotracheal intubation:**
 - Procedure requires spontaneous breathing and may require considerable time, exposing patient to critical desaturation. Contraindicated in combative, anatomically disrupted or distorted airways, increased ICP, severe facial trauma, basal skull fracture, and head injury. Orotracheal route is preferred.
- Intubation attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.
- If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment)
- AEMT and Paramedics should consider using a BIAD if oral-tracheal intubation is unsuccessful.
- During intubation attempts use External Laryngeal Manipulation to improve view of glottis.
- Gastric tube placement should be considered in all intubated patients if available or time allows.
- It is important to secure the endotracheal tube well to better maintain ETT placement. Manual stabilization of endotracheal tube should be used during all patient moves / transfers.
- **DOPE:** Displaced tracheostomy tube / ETT. Obstructed tracheostomy tube / ETT. Pneumothorax and Equipment failure.



Adult, Failed Airway

Protocols AR 1, 2, and 3 should be utilized together (even if agency is not using Drug Assisted Airway as they contain useful information for airway management.

Unable to Ventilate and Oxygenate $\geq 90\%$ during or after one (1) or more unsuccessful intubation attempts.
and/or
Anatomy inconsistent with continued attempts.
and/or
Three (3) unsuccessful attempts by most experienced Paramedic/AEMT.
Each attempt should include change in approach or equipment

NO MORE THAN THREE (3) ATTEMPTS TOTAL

- Capnography Monitoring**
- End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.
 - EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.

Failed Airway

Call for additional resources if available

BVM
Adjunctive Airway NP / OP
Maintains
Oxygen Saturation $\geq 90\%$
Preferably $\geq 94\%$

Continue BVM
Supplemental Oxygen

Exit to
Appropriate Protocol(s)

NO

B	Attempt Airway Blind Insertion Airway Device Procedure
A	Airway Video Laryngoscopy Device Procedure <i>if available</i> Optional
P	Airway Cricothyrotomy Surgical Procedure
	Supplemental oxygen BVM with Airway Adjuncts Maintain Oxygen Saturation $\geq 90\%$ Preferably $\geq 94\%$
	Post-intubation BIAD Management Protocol AR 8

 **Notify Destination or Contact Medical Control** 



Adult, Failed Airway

Pearls

- **For the purposes of this protocol a secure airway is when the patient is receiving appropriate oxygenation and ventilation.**
- **If an effective airway is being maintained by BVM with continuous pulse oximetry values of $\geq 90\%$, it is acceptable to continue with basic airway measures.**
- **Ventilation rate should be 8-10 per minute to maintain a EtCO₂ of 35-45. Avoid hyperventilation.**
- **Anticipating the Difficult Airway and Airway Assessment:**
 - **Difficult BVM Ventilation (MOANS):** Mask seal difficulty (hair, secretions, trauma); Obese, obstruction, OB – 2d and 3d trimesters; Age ≥ 55 ; No teeth; Stiff lungs or neck
 - **Difficult Laryngoscopy (LEON):** Look externally for anatomical problems; Evaluate 3-3-2 (Mouth opening should equal 3 of patient's finger's width, mental area to neck should equal 3 of patient's finger's width, base of chin to thyroid prominence should equal 2 of patient's finger's width); Obese, obstruction, OB – 2d and 3d trimesters; Neck mobility limited.
 - **Difficulty BIAD (RODS):** Restricted mouth opening; Obese, obstruction, OB – 2d and 3d trimesters; Distorted or disrupted airway; Stiff lungs or neck
 - **Difficulty Cricothyrotomy / Surgical Airway (SMART):** Surgery scars; Mass or hematoma, Access or anatomical problems; Radiation treatment to face, neck, or chest; Tumor.
- **Capnography Monitoring (EtCO₂):**
 - **Continuous Waveform or Quantitative Capnography and Pulse Oximetry are required for intubation verification and ongoing patient monitoring (Not validated and may prove impossible in the neonatal population - verification by two (2) other means is recommended in this population.)**
 - **Capnography verification and monitoring is required for BIAD verification and monitoring once available on scene.**
- **Nasotracheal intubation:**
 - **Procedure requires spontaneous breathing and may require considerable time, exposing patient to critical desaturation. Contraindicated in combative, anatomically disrupted or distorted airways, increased ICP, severe facial trauma, basal skull fracture, and head injury. Orotracheal route is preferred.**
 - **Intubation attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.**
 - **If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment)**
 - **AEMT and Paramedics should consider using a BIAD if oral-tracheal intubation is unsuccessful.**
 - **During intubation attempts use External Laryngeal Manipulation to improve view of glottis.**
 - **Gastric tube placement should be considered in all intubated patients if available or time allows.**
 - **It is important to secure the endotracheal tube well to better maintain ETT placement. Manual stabilization of endotracheal tube should be used during all patient moves / transfers.**
 - **DOPE: Displaced tracheostomy tube / ETT, Obstructed tracheostomy tube / ETT, Pneumothorax and Equipment failure.**

Adult COPD / Asthma Respiratory Distress

History


- Asthma; COPD -- chronic bronchitis, emphysema, congestive heart failure
- Home treatment (oxygen, nebulizer)
- Medications (theophylline, steroids, inhalers)
- Toxic exposure, smoke inhalation

Signs and Symptoms

- Shortness of breath
- Pursed lip breathing
- Decreased ability to speak
- Increased respiratory rate and effort
- Wheezing, rhonchi
- Use of accessory muscles
- Fever, cough
- Tachycardia

Differential

- Asthma
- Anaphylaxis
- Aspiration
- COPD (Emphysema, Bronchitis)
- Pleural effusion
- Pneumonia
- Pulmonary embolus
- Pneumothorax
- Cardiac (MI or CHF)
- Pericardial tamponade
- Hyperventilation
- Inhaled toxin (Carbon monoxide, etc.)



Adult COPD / Asthma Respiratory Distress

May substitute Albuterol 2.5 mg with Levalbuterol 1.25 mg as needed

COVID-19 has precipitated respiratory emergencies that require treatment via nebulized medications. The CDC has advised that these procedures are "HIGH RISK" and require full PPE. Because nebulized therapy with bronchodilators for presumptive or confirmed COVID-19 patients may not be safe due to generation of aerosols, which increase the risk that respiratory droplets will remain in the air and spread the virus, delivery of those drugs via Metered-Dose Inhaler (MDI) is preferred.

Best practices for management of the COVID-19 are dynamic due to the rapidly evolving situation and changing scientific knowledge. In the interest of provider safety, infection control, and high-quality patient care, the use of MDI's are allowed "ONLY" in the presumptive or confirmed COVID-19 patient. MDI's are a single patient use. MDI's are NOT to replace nebulized treatment on the non COVID-19 patients.

The patient that is presenting with respiratory distress AND COVID-19 signs and symptoms:

Adult patient: Albuterol 2 puffs (90mcg each puff), repeat 1 time (2 puffs) after 10 minutes if no improvement

Pediatrics patient: (>4 years): Albuterol 1 puff (90mcg each puff), repeat 1 time (1 puff) after 10 minutes if no improvement.


If respiratory distress persists, or if the patient is < 4 years, Contact Medical Control to discuss further treatment options

Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Neck, Heart, Lungs, Abdomen, Extremities, Neuro**
- **Items in Red Text are key performance measures used to evaluate protocol compliance and care.**
- **This protocol includes all patients with respiratory distress, COPD, Asthma, Reactive Airway Disease, or Bronchospasm. Patients may also have wheezing and respiratory distress with viral upper respiratory tract infections and pneumonia.**
- **Combination nebulizers containing albuterol and ipratropium:**
 - Patients may receive more than 3 nebulizer treatments, treatments should continue until improvement. Following 3 combination nebulizers, it is acceptable to continue albuterol solely with subsequent treatments as there is no proven benefit to continual use of ipratropium.
- **Epinephrine:**
 - If allergic reaction or anaphylaxis is suspected, give immediately and repeat until improvement.
 - If allergic reaction is not suspected, administer with impending respiratory failure and no improvement.
 - Consider Magnesium Sulfate with impending respiratory failure and no improvement.
 - Pulse oximetry should be monitored continuously and consider End-tidal CO₂ monitoring if available.
- **CPAP or Non-Invasive Positive Pressure Ventilation:**
 - May be used with COPD, Asthma, Allergic reactions, and CHF.
 - Consider early in treatment course.
 - Consider removal if SBP remains < 100 mmHg and not responding to other treatments.
- A silent chest in respiratory distress is a pre-respiratory arrest sign.
- **EMT may administer Albuterol if patient already prescribed and may administer from EMS supply.** Agency Medical Director may require contact of medical control prior to EMT / EMR administering any medication.



Pediatric Airway



Capnography Monitoring

- End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.
- EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.

Unable to Ventilate and Oxygenate ≥ 90% during or after one (1) or more unsuccessful intubation attempts . and/or Anatomy inconsistent with continued attempts. and/or Three (3) unsuccessful attempts by most experienced Paramedic/AEMT.
Exit to Pediatric Failed Airway Protocol AR 6



Pediatric Airway

Pearls

- **This protocol is for use in patients who FIT within a Pediatric Medication/Skill Resuscitation System Product.**
- For the purposes of this protocol a secure airway is when the patient is receiving appropriate oxygenation and ventilation.
- If an effective airway is being maintained by BVM with continuous pulse oximetry values of $\geq 90\%$, it is acceptable to continue with basic airway measures.
- **Ventilation rate:**
30 for Neonates, 25 for Toddlers, 20 for School Age, and for Adolescents the normal Adult rate of 8 - 10 per minute.
Maintain a EtCO₂ between 35 and 45 and avoid hyperventilation.
- **Ketamine:**
May be used during airway management of patients who FIT within a Pediatric Medication/Skill Resuscitation System product with a DIRECT, ONLINE MEDICAL ORDER, by the system MEDICAL DIRECTOR OR ASSISTANT MEDICAL DIRECTOR ONLY.
Systems using Ketamine in the pediatric population must also be using in their adult population.
- **Agencies utilizing Ketamine must submit a local systems plan to State Medical Director detailing how the drug is used in your program.**
Ketamine may be used within this protocol only WITHOUT a paralytic agent in conjunction with either a OP, NP, BIAD or endotracheal tube.
Ketamine may be used during the resuscitation of hypoxia or hypotension in conjunction with airway management.
Ketamine may be used in the dangerously combative patient requiring airway management IM. IV / IO should be established as soon as possible.
Ketamine may NOT be used for purposes of sedation only – it must be used only during airway management procedures.
- **Capnography Monitoring (EtCO₂):**
Continuous Waveform or Quantitative Capnography and Pulse Oximetry are required for intubation verification and ongoing patient monitoring (Not validated and may prove impossible in the neonatal population - verification by two (2) other means is recommended in this population.)
Capnography verification and monitoring is required for BIAD verification and monitoring once available on scene.
- Intubation attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.
- If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment)
- AEMT and Paramedics should consider using a BIAD if oral-tracheal intubation is unsuccessful.
- During intubation attempts use External Laryngeal Manipulation to improve view of glottis.
- It is important to secure the endotracheal tube well and consider c-collar (even in absence of trauma) to better maintain ETT placement. Manual stabilization of endotracheal tube should be used during all patient moves / transfers.
- **DOPE:** Displaced tracheostomy tube / ETT, Obstructed tracheostomy tube / ETT, Pneumothorax and Equipment failure.



Pediatric Failed Airway

Unable to Ventilate and Oxygenate $\geq 90\%$ during or after one (1) or more unsuccessful intubation attempts.
and/or
Anatomy inconsistent with continued attempts.
and/or
Three (3) unsuccessful attempts by most experienced Paramedic / AEMT.
Each attempt should include change in approach or equipment

NO MORE THAN THREE (3) ATTEMPTS TOTAL

Call for additional resources if available

Failed Airway

BVM
Adjunctive Airway NP / OP
Maintains
Oxygen Saturation $\geq 90\%$
Preferably $\geq 94\%$

Continue BVM
Supplemental Oxygen

Exit to Appropriate Protocol(s)

A Airway Video Laryngoscopy Device Procedure
if available
Optional

B Attempt Airway Blind Insertion Airway Device Procedure

P

BIAD / Cricothyrotomy
Successful
Or
Oxygenation / Ventilation Adequate

Exit to Post-intubation / BIAD Management Protocol AR 8

Capnography Monitoring

- End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.
- EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.

Supplemental oxygen
BVM with Airway Adjuncts
Maintain Oxygen Saturation $\geq 90\%$
Preferably $\geq 94\%$

Notify Destination or Contact Medical Control

Airway Respiratory Protocol Section



Pediatric Failed Airway

Pearls

- **This protocol is for use in patients who FIT within a Pediatric Medication/Skill Resuscitation System Product.**
- For the purposes of this protocol a secure airway is when the patient is receiving appropriate oxygenation and ventilation.
- If an effective airway is being maintained by BVM with continuous pulse oximetry values of $\geq 90\%$, it is acceptable to continue with basic airway measures.
- **Ventilation rate:**
30 for Neonates, 25 for Toddlers, 20 for School Age, and for Adolescents the normal Adult rate of 8 - 10 per minute.
Maintain a EtCO₂ between 35 and 45 and avoid hyperventilation.
- **Ketamine:**
May be used during airway management of patients who FIT within a Pediatric Medication/Skill Resuscitation System product with a DIRECT, ONLINE MEDICAL ORDER, by the system MEDICAL DIRECTOR OR ASSISTANT MEDICAL DIRECTOR ONLY.
Systems using Ketamine in the pediatric population must also be using in their adult population.
- **Agencies utilizing Ketamine must submit a local systems plan to State Medical Director detailing how the drug is used in your program.**
Ketamine may be used within this protocol only WITHOUT a paralytic agent in conjunction with either a OP, NP, BIAD or endotracheal tube.
Ketamine may be used during the resuscitation of hypoxia or hypotension in conjunction with airway management.
Ketamine may be used in the dangerously combative patient requiring airway management IM. IV / IO should be established as soon as possible.
Ketamine may NOT be used for purposes of sedation only – it must be used only during airway management procedures.
- **Capnography Monitoring (EtCO₂):**
Continuous Waveform or Quantitative Capnography and Pulse Oximetry are required for intubation verification and ongoing patient monitoring (Not validated and may prove impossible in the neonatal population - verification by two (2) other means is recommended in this population.)
Capnography verification and monitoring is required for BIAD verification and monitoring once available on scene.
- Intubation attempt defined as laryngoscope blade passing the teeth or endotracheal tube passed into the nostril.
- If First intubation attempt fails, make an adjustment and try again: (Consider change of provider in addition to equipment)
- AEMT and Paramedics should consider using a BIAD if oral-tracheal intubation is unsuccessful.
- During intubation attempts use External Laryngeal Manipulation to improve view of glottis.
- It is important to secure the endotracheal tube well and consider c-collar (even in absence of trauma) to better maintain ETT placement.
Manual stabilization of endotracheal tube should be used during all patient moves / transfer
- **DOPE:** Displaced tracheostomy tube / ETT, Obstructed tracheostomy tube / ETT, Pneumothorax and Equipment failure.

Pediatric Asthma Respiratory Distress

History


- Time of onset
- Possibility of foreign body
- Past Medical History
- Medications
- Fever / Illness
- Sick Contacts
- History of trauma
- History / possibility of choking
- Ingestion / OD
- Congenital heart disease

Signs and Symptoms

- Wheezing / Stridor / Crackles / Rales
- Nasal Flaring / Retractions / Grunting
- Increased Heart Rate
- AMS
- Anxiety
- Attentiveness / Distractability
- Cyanosis
- Poor feeding
- JVD / Frothy Sputum
- Hypotension

Differential

- Asthma / Reactive Airway Disease
- Aspiration
- Foreign body
- Upper or lower airway infection
- Congenital heart disease
- OD / Toxic ingestion / CHF
- Anaphylaxis
- Trauma



Pediatric Asthma Respiratory Distress

May substitute Albuterol 2.5 mg with Xopenox 1.25 mg as needed

COVID-19 has precipitated respiratory emergencies that require treatment via nebulized medications. The CDC has advised that these procedures are "HIGH RISK" and require full PPE. Because nebulized therapy with bronchodilators for presumptive or confirmed COVID-19 patients may not be safe due to generation of aerosols, which increase the risk that respiratory droplets will remain in the air and spread the virus, delivery of those drugs via Metered-Dose Inhaler (MDI) is preferred.

Best practices for management of the COVID-19 are dynamic due to the rapidly evolving situation and changing scientific knowledge. In the interest of provider safety, infection control, and high-quality patient care, the use of MDI's are allowed "ONLY" in the presumptive or confirmed COVID-19 patient. MDI's are a single patient use. MDI's are NOT to replace nebulized treatment on the non COVID-19 patients.

The patient that is presenting with respiratory distress AND COVID-19 signs and symptoms:

Adult patient: Albuterol 2 puffs (90mcg each puff), repeat 1 time (2 puffs) after 10 minutes if no improvement

Pediatrics patient: (>4 years): Albuterol 1 puff (90mcg each puff), repeat 1 time (1 puff) after 10 minutes if no improvement.
If respiratory distress persists, or if the patient is < 4 years, Contact Medical Control to discuss further treatment options

Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Neck, Heart, Lungs, Abdomen, Extremities, Neuro**
- **Items in Red Text are key performance measures used to evaluate protocol compliance and care.**
- **Pulse oximetry should be monitored continuously in the patient with respiratory distress.**
- **This protocol includes all patients with respiratory distress, Asthma, Reactive Airway Disease, croup, or Bronchospasm. Patients may also have wheezing and respiratory distress with viral upper respiratory tract infections and pneumonia.**
- **Combination nebulizers containing albuterol and ipratropium:**
 - Patients may receive more than 3 nebulizer treatments, treatments should continue until improvement. Following 3 combination nebulizers, it is acceptable to continue albuterol solely with subsequent treatments as there is no proven benefit to continual use of ipratropium.
- **Epinephrine:**
 - If allergic reaction or anaphylaxis is suspected, give immediately and repeat until improvement.
 - If allergic reaction is not suspected, administer with impending respiratory failure and no improvement.
- **Consider Magnesium Sulfate with impending respiratory failure and no improvement.**
- **Albuterol dosing: ≤ 1 year of age 1.25 mg; 1 – 6 years 1.25 – 2.5 mg; 6 – 14 years 2.5 mg; ≥ 15 years 2.5 – 5 mg.**
- **Consider IV access when Pulse oximetry remains ≤ 92 % after first beta agonist treatment.**
- **Do not force a child into a position, allow them to assume position of comfort. They will protect their airway by their body position.**
- Bronchiolitis is a viral infection typically affecting infants which results in wheezing which may not respond to beta-agonists. Consider Epinephrine nebulizer if patient < 18 months and not responding to initial beta-agonist treatment.
- Croup typically affects children < 2 years of age. It is viral, possible fever, gradual onset, no drooling is noted.
- Epiglottitis typically affects children > 2 years of age. It is bacterial, with fever, rapid onset, possible stridor, patient wants to sit up to keep airway open, drooling is common. Airway manipulation may worsen the condition.
- In patients using levalbuterol (Xopenex) you may use Albuterol for the first treatment then use the patient's supply for repeat nebulizers or agency's supply.
- **EMT may administer Albuterol if patient already prescribed and may administer from EMS supply.** Agency medical director may require Contact of Medical Control prior to administration.

Revised
01/01/2022

AR 7

This protocol has been altered from the original NCEP Protocol by the local EMS Medical Director




Post-intubation / BIAD Management

Protocols AR 1, 2, 3, 5, and 6 should be utilized together (even if agency is not using Drug Assisted Airway Protocol) as they contain useful information for airway management.

Capnography Monitoring

- End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.
- EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.



Airway Respiratory Protocol Section



Post-intubation / BIAD Management

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Neuro**
- **Patients requiring advanced airways and ventilation commonly experience pain and anxiety.**
- **Unrelieved pain can lead to increased catecholamine release, ischemia, immunosuppression, and prolonged hospitalization.**
- **Ventilated patients cannot communicate pain / anxiety and providers are poor at recognizing pain / anxiety.**
- **Vital signs such as tachycardia and / or hypertension can provide clues to inadequate sedation, however they both are not always reliable indicators of patient's lack of adequate sedation.**
- **Pain must be addressed first, before anxiety. Opioids are typically the first line agents before benzodiazepines. Ketamine is also a reasonable first choice agent.**
- **Ventilator / Ventilation strategies will need to be tailored to individual patient presentations. Medical director can indicate different strategies above.**
- In general ventilation with BVM should cause chest rise. With mechanical ventilation a reasonable tidal volume should be about 6 mL/kg and peak pressures should be < 30 cmH2O.
- Continuous pulse oximetry and capnography should be maintained during transport for monitoring.
- Head of bed should be maintained at least 10 – 20 degrees of elevation when possible to decrease aspiration risk.
- With abrupt clinical deterioration, if mechanically ventilated, disconnect from ventilator to assess lung compliance. Search for dislodged ETT or BIAD, obstruction in tubing or airway, pneumothorax, or ETT balloon leak.
- **DOPE: Displaced tracheostomy tube / ETT, Obstructed tracheostomy tube / ETT, Pneumothorax and Equipment failure.**



Ventilator Emergencies

History


- Birth defect (tracheal atresia, tracheomalacia, craniofacial abnormalities)
- Surgical complications (damage to phrenic nerve)
- Trauma (post-traumatic brain or spinal cord injury)
- Medical condition (bronchopulmonary dysplasia, muscular dystrophy)

Signs and Symptoms

- Transport requiring maintenance of a mechanical ventilator
- Power or equipment failure at residence

Differential

- Disruption of oxygen source
- Dislodged or obstructed tracheostomy tube
- Detached or disrupted ventilator circuit
- Cardiac arrest
- Increased oxygen requirement / demand
- Ventilator failure



Airway Respiratory Protocol Section

Pearls

- **Always talk to family / caregivers as they have specific knowledge and skills.**
- **If using the patient's ventilator bring caregiver knowledgeable in ventilator operation during transport.**
- Always use patient's equipment if available and functioning properly.
- Continuous pulse oximetry and end tidal CO2 monitoring must be utilized during assessment and transport.
- Unable to correct ventilator problem: Remove patient from ventilator and manually ventilate using BVM. Take patient's ventilator to hospital even if not functioning properly.
- Typical alarms:
 - Low Pressure / Apnea: Loose or disconnected circuit, leak in circuit or around tracheostomy site.
 - Low Power: Internal battery depleted.
 - High Pressure: Plugged / obstructed airway or circuit.
- **DOPE: Displaced tracheostomy tube / ETT, Obstructed tracheostomy tube / ETT, Pneumothorax and Equipment failure.**



Tracheostomy Tube Emergencies

History




- Birth defect (tracheal atresia, tracheomalacia, craniofacial abnormalities)
- Surgical complications (accidental damage to phrenic nerve)
- Trauma (post-traumatic brain or spinal cord injury)
- Medical condition (bronchial or pulmonary dysplasia, muscular dystrophy)

Signs and Symptoms

- Nasal flaring
- Chest wall retractions (with or without abnormal breath sounds)
- Attempts to cough
- Copious secretions noted coming out of the tube
- Faint breath sounds on both sides of chest despite significant respiratory effort
- AMS
- Cyanosis

Differential

- Allergic reaction
- Asthma
- Aspiration
- Septicemia
- Foreign body
- Infection
- Congenital heart disease
- Medication or toxin
- Trauma



Airway Respiratory Protocol Section

Pearls

- Always talk to family / caregivers as they have specific knowledge and skills.
- Important to ask if patient has undergone laryngectomy. This does not allow mouth/nasal ventilation by covering stoma.
- Use patients equipment if available and functioning properly.
- Estimate suction catheter size by doubling the inner tracheostomy tube diameter and rounding down.
- Suction depth: Ask family / caregiver. No more than 3 to 6 cm typically. Instill 2 – 3 mL of NS before suctioning.
- Do not suction more than 10 seconds each attempt and pre-oxygenate before and between attempts.
- DO NOT force suction catheter. If unable to pass, then tracheostomy tube should be changed.
- Always deflate tracheal tube cuff before removal. Continual pulse oximetry and EtCO2 monitoring if available.
- **DOPE:** Displaced tracheostomy tube / ETT, Obstructed tracheostomy tube / ETT, Pneumothorax and Equipment failure.



Adult Asystole / Pulseless Electrical Activity

History

- SAMPLE
- Estimated downtime
- See Reversible Causes below
- DNR, MOST, or Living Will

Signs and Symptoms

- Pulseless
- Apneic
- No electrical activity on ECG
- No heart tones on auscultation

Differential

- See Reversible Causes below



Cardiac Arrest Protocol AC 3

Criteria for Death / No Resuscitation
Review DNR / MOST Form

YES

NO

Decomposition
Rigor mortis
Dependent lividity
Blunt force trauma
Injury incompatible with life
Extended downtime with asystole

Do not begin resuscitation

Follow
Deceased Subjects
Policy

AT ANY TIME

Return of
Spontaneous
Circulation



Go to
Post Resuscitation
Protocol AC 10

Begin Continuous CPR Compressions
Push Hard (≥ 2 inches)
Push Fast (100 - 120 / min)
Change Compressors every 2 minutes
(sooner if fatigued)
(Limit changes / pulse checks ≤ 10 seconds)

Ventilate 1 breath every 6 seconds
30:2 Compression:Ventilation if no Advanced Airway
Monitor EtCO₂
if available

AED Procedure
if available

P

Cardiac Monitor



IV or IO Access Protocol UP 6

A

Epinephrine (1:10,000) 1 mg IV / IO
Repeat every 5 minutes

Search for Reversible Causes

Blood Glucose Analysis Procedure
if applicable

P



On Scene Resuscitation / Termination of Resuscitation
Protocol(s) AC 12
as indicated

Reversible Causes

Hypovolemia
Hypoxia
Hydrogen ion (acidosis)
Hypothermia
Hypo / Hyperkalemia

Tension pneumothorax
Tamponade; cardiac
Toxins
Thrombosis; pulmonary (PE)
Thrombosis; coronary (MI)

Suspected Opioid Overdose

Administer Naloxone per
Overdose / Toxic Ingestion
Protocol TE 7



Notify Destination or
Contact Medical Control





Adult Asystole / Pulseless Electrical Activity

Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional Team Focused CPR Protocol AC 11 or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.**
- **DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT), compression to ventilation ratio is 30:2. If advanced airway in place, ventilate 10 breaths per minute with continuous, uninterrupted compressions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **Passive oxygenation optional in agencies practicing Team Focused Approach / Pit-Crew Approach.**
- **Reassess and document BIAD and / or endotracheal tube placement and EtCO₂ frequently, after every move, and at transfer of care.**
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Defibrillation:** Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
- **End Tidal CO₂ (EtCO₂)**
 - If EtCO₂ is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
 - If EtCO₂ spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **Special Considerations**
 - **Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm. Defibrillation is safe at all energy levels.
 - **Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
 - **Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol TE 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
 - **Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Transcutaneous Pacing:**
 - Pacing is NOT effective in cardiac arrest and pacing in cardiac arrest does NOT increase chance of survival
 - Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.
 - Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment options.



Bradycardia; Pulse Present

History


- Past medical history
- Medications
 - Beta-Blockers
 - Calcium channel blockers
 - Clonidine
 - Digoxin
- Pacemaker

Signs and Symptoms


- HR < 60/min with hypotension, acute altered mental status, chest pain, acute CHF, seizures, syncope, or shock secondary to bradycardia
- Chest pain
- Respiratory distress
- Hypotension or Shock
- Altered mental status
- Syncope

Differential

- Acute myocardial infarction
- Hypoxia / Hypothermia
- Pacemaker failure
- Sinus bradycardia
- Head injury (elevated ICP) or Stroke
- Spinal cord lesion
- Sick sinus syndrome
- AV blocks (1°, 2°, or 3°)
- Overdose



YES



Suspected Beta-Blocker or Calcium Channel Blocker

Follow Overdose/ Toxic Ingestion Protocol TE 7

Reversible Causes

Hypovolemia
Hypoxia
Hydrogen ion (acidosis)
Hypothermia
Hypo / Hyperkalemia

Tension pneumothorax
Tamponade; cardiac
Toxins
Thrombosis; pulmonary (PE)
Thrombosis; coronary (MI)





Bradycardia; Pulse Present

Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Identifying signs and symptoms of poor perfusion caused by bradycardia are paramount.**
- **Rhythm should be interpreted in the context of symptoms and pharmacological treatment given only when symptomatic, otherwise monitor and reassess.**
- **Consider hyperkalemia with wide complex, bizarre appearance of QRS complex, and bradycardia. Give Calcium Chloride or Gluconate in addition to Sodium Bicarbonate if hyperkalemia suspected.**
- **12-Lead ECG:**
12 Lead ECG not necessary to diagnose and treat
Obtain when patient is stable and/or following rhythm conversion.
- **Unstable condition**
Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
If at any point patient becomes unstable move to unstable arm in algorithm.
- **Hypoxemia is a common cause of bradycardia. Ensure oxygenation and support respiratory effort.**
- **Atropine:**
Atropine is considered a first line agent in symptomatic bradycardia.
Ineffective and potentially harmful in cardiac transplantation. May cause paradoxical bradycardia.
- **Symptomatic bradycardia causing shock or peri-arrest condition:**
If no IV or IO access immediately available start Transcutaneous Pacing, establish IV / IO access, and then administer atropine and/or epinephrine.
Epinephrine or Dopamine may be considered if no response to Atropine.
- **Symptomatic condition**
Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
Symptomatic bradycardia usually occurs at rates < 50 beats per minute.
Search for underlying causes such as hypoxia or impending respiratory failure.
- **Serious Signs / Symptoms:**
Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute CHF.
- **Transcutaneous Pacing Procedure (TCP)**
Indicated with unstable bradycardia unresponsive to medical therapy.
If time allows transport to specialty center because transcutaneous pacing is a temporizing measure.
Transvenous / permanent pacemaker will probably be needed.
Immediate TCP with high-degree AV block (2d or 3d degree) with no IV / IO access.
- Consider treatable causes for bradycardia (Beta Blocker OD, Calcium Channel Blocker OD, etc.)

Cardiac Arrest; Adult

AT ANY TIME

Return of Spontaneous Circulation



Go to Post Resuscitation Protocol AC 10

Criteria for Death / No Resuscitation
Review DNR / MOST Form

YES

Decomposition
Rigor mortis
Dependent lividity
Blunt force trauma
Injury incompatible with life
Extended downtime with asystole

Do not begin resuscitation

Follow Deceased Subjects Policy

NO

Begin Continuous CPR Compressions
Push Hard (≥ 2 inches)
Push Fast (100 - 120 / min)
Change Compressors every 2 minutes (sooner if fatigued)
(Limit changes / pulse checks ≤ 10 seconds)

Ventilate 1 breath every 6 seconds
30:2 Compression:Ventilation if no Advanced Airway
Monitor EtCO2 if available

AED Procedure *if available*

ALS Available

P Cardiac Monitor

NO Shockable Rhythm

AED Procedure

YES

Asystole / PEA
Protocol AC 1 *as indicated*

Airway
Protocol(s) AR 1, 2, 3

NO Shockable Rhythm

YES

VF / VT
Protocol AC 9
Tachycardia
Protocol(s) AC 6, 7 *as indicated*

Airway
Protocol(s) AR 1, 2, 3

Continue CPR 2 Minutes

Repeat and reassess

Airway
Protocol(s) AR 1, 2, 3

Arrest secondary to Opioid OD?

YES

Naloxone 0.4 – 2 mg IN / IM
Peds: 0.1 mg/kg IN

Maximum 4 mg

A

NO

Termination on Scene
Protocol AC 12 *as indicated*

 **Notify Destination or Contact Medical Control** 



Cardiac Arrest; Adult

Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assign responders to predetermined tasks. Refer to optional protocol or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.**
- **DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT) compression to ventilation ratio is 30:2. If advanced airway in place, ventilate 10 breaths per minute with continuous, uninterrupted compressions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **Passive oxygenation optional in agencies practicing Team Focused Approach / Pit-Crew Approach.**
- **Reassess and document BIAD and / or endotracheal tube placement and EtCO₂ frequently, after every move, and at transfer of care.**
- **IV / IO access and drug delivery is secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Defibrillation:**
 - Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
 - Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.
 - Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- **End Tidal CO₂ (EtCO₂)**
 - If EtCO₂ is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
 - If EtCO₂ spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **Special Considerations**
 - Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm. Defibrillation is safe at all energy levels.
 - Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
 - Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol TE 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
 - Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Transcutaneous Pacing:**
 - Pacing is NOT effective in cardiac arrest and pacing in cardiac arrest does NOT increase chance of survival
- Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.
- Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment



Chest Pain: Cardiac and STEMI

History

- Age
- Medications (Viagra / sildenafil, Levitra / vardenafil, Cialis / tadalafil)
- Past medical history (MI, Angina, Diabetes, post menopausal)
- Allergies
- Recent physical exertion
- Onset / Palliation / Provocation
- Quality (crampy, constant, sharp, dull, etc.)
- Region / Radiation / Referred
- Severity (1-10)
- Time (onset / duration / repetition)

Signs and Symptoms

- CP (pain, pressure, aching, vice-like tightness)
- Location (substernal, epigastric, arm, jaw, neck, shoulder)
- Radiation of pain
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness
- **Time of Onset**
- Women:
 - More likely to have dyspnea, N/V, weakness, back or jaw pain

Differential

- Trauma vs. Medical
- Angina vs. Myocardial infarction
- Pericarditis
- Pulmonary embolism
- Asthma / COPD
- Pneumothorax
- Aortic dissection or aneurysm
- GE reflux or Hiatal hernia
- Esophageal spasm
- Chest wall injury or pain
- Pleural pain
- Overdose: Cocaine or Methamphetamine

	12 Lead ECG Procedure
B	Aspirin 81 mg x 4 PO (chewed) Or 325 mg PO
	Nitroglycerin 0.3 / 0.4 mg Sublingual Repeat every 5 minutes x 3 <i>if prescribed to patient and (BP ≥ 100)</i>
P	Cardiac Monitor

Acute MI / STEMI
See box to right

Transport based on:
STEMI!
EMS Triage and Destination Plan
Immediate Notification of Facility
Immediate Transmission of ECG
if capable
Keep Scene Time to ≤ 15 Minutes

Acute MI / STEMI

STEMI Definition:

- ≥ 1 mm ST Segment elevation in ≥ 2 contiguous leads
- ECG software diagnoses Acute MI (symptomatic)

	IV / IO Protocol UP 6
A	Nitroglycerin 0.3 / 0.4 mg SL Repeat every 5 minutes as needed
P	
	Hypotension / Shock Protocol AM 5 <i>if indicated</i>
	CHF / Pulmonary Edema Protocol AC 5 <i>if indicated</i>

B	<i>If transporting to Non PCI Center</i> Reperfusion Checklist
P	

Notify Destination or Contact Medical Control



Chest Pain: Cardiac and STEMI

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Items in Red Text are the key performance indicators for the EMS Acute Cardiac (STEMI) Care Toolkit**
- **Nitroglycerin:**
 - Avoid Nitroglycerin in any patient who has used Viagra (sildenafil) or Levitra (vardenafil) in the past 24 hours or Cialis (tadalafil) in the past 36 hours due to potential severe hypotension.
 - Nitroglycerin may cause hypotension during any type myocardial infarction. It is NOT more likely to cause hypotension in an inferior MI and should NOT be avoided unless already hypotensive.
- **STEMI (ST-Elevation Myocardial Infarction)**
 - Positive Reperfusion Checklist should be transported to the appropriate facility based on STEMI EMS Triage and Destination Plan.
 - Consider placing 2 IV sites in the left arm: Many PCI centers use the right radial artery for intervention.
 - Consider placing defibrillator pads on patient as a precaution.
 - Consider Normal Saline or Lactated Ringers bolus of 250 – 500 mL as pre-cath hydration.
 - Scene time goal is < 15 minutes.
 - Document and time-stamp facility STEMI notification and make notification as soon as possible.
 - Document the time of the 12-Lead ECG in the PCR as a Procedure along with the interpretation (Paramedic).
- **Cardiac related symptoms in men and women:**
 - Pressure, squeezing, fullness, or pain in the chest.
 - Pain or discomfort in one or both arms, the back, neck, jaw, or stomach.
 - Shortness of breath with or without chest pain.
 - Sweating, nausea, weakness, and/or lightheadedness.
 - **Women, diabetic patients, and the elderly often experience only weakness, shortness of breath, nausea/ vomiting, and back or jaw pain.**
- If patient has taken nitroglycerin without relief, consider potency of the medication.
- Monitor for hypotension after administration of nitroglycerin and opioids.
- **EMT may administer Nitroglycerin to patients already prescribed medication. May give from EMS supply.**
- Agency medical director may require Contact of Medical Control prior to administration.



CHF / Pulmonary Edema

History

- Congestive heart failure
- Past medical history
- Medications (digoxin, Lasix, Viagra / sildenafil, Levitra / vardenafil, Cialis / tadalafil)
- Cardiac history --past myocardial infarction

Signs and Symptoms

- Respiratory distress, bilateral rales
- Apprehension, orthopnea
- Jugular vein distention
- Pink, frothy sputum
- Peripheral edema, diaphoresis
- Hypotension, shock
- Chest pain

Differential

- Myocardial infarction
- Congestive heart failure
- Asthma
- Anaphylaxis
- Aspiration
- COPD
- Pleural effusion
- Pneumonia
- Pulmonary embolus
- Pericardial tamponade
- Toxic Exposure

	Airway Protocol(s) AR 1, 2, 3 as indicated
	Chest Pain and STEMI Protocol AC 4 if indicated
B	12 Lead ECG Procedure
	Nitroglycerin 0.3 / 0.4 mg Sublingual Repeat every 5 minutes x 3 if prescribed to patient and (BP >100)
P	Cardiac Monitor
	IV / IO Procedure

Assess Symptom Severity

MILD
Normal Heart Rate
Elevated or Normal BP

MODERATE / SEVERE
Elevated Heart Rate
Elevated BP

CARDIOGENIC SHOCK
Tachycardia followed by bradycardia
Hypertension followed by hypotension

A	Nitroglycerin 0.3 / 0.4 mg SL Repeat every 5 minutes

B	Airway NIPPV Procedure
A	Nitroglycerin 0.3 / 0.4 mg SL Repeat every 5 minutes
P	

B	Remove NIPPV if in place
	Adult Hypotension / Shock Protocol AM 5 if indicated

Improving
YES
NO

Notify Destination or Contact Medical Control



CHF / Pulmonary Edema

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Items in Red Text are key performance measures used to evaluate protocol compliance and care**
- **Diuretics (furosemide) and opioids have NOT been shown to improve the outcomes of EMS patients with pulmonary edema. Even though this historically has been a mainstay of EMS treatment, it is no longer routinely recommended.**
- **Nitroglycerin:**
 - **Avoid Nitroglycerin in any patient who has used Viagra (sildenafil) or Levitra (vardenafil) in the past 24 hours or Cialis (tadalafil) in the past 36 hours due to potential severe hypotension.**
 - **Nitroglycerin may cause hypotension during any type myocardial infarction. It is NOT more likely to cause hypotension in an inferior MI and should NOT be avoided unless already hypotensive.**
- **Document the time of the 12-Lead ECG in the PCR as a Procedure along with the interpretation (Paramedic).**
- **Consider myocardial infarction in all these patients. Diabetics, geriatric and female patients often have atypical pain, or only generalized complaints.**
- **Cardiac related symptoms in men and women:**
 - **Pressure, squeezing, fullness, or pain in the chest.**
 - **Pain or discomfort in one or both arms, the back, neck, jaw, or stomach.**
 - **Shortness of breath with or without chest pain.**
 - **Sweating, nausea, weakness, and/or lightheadedness.**
 - **Women, diabetic patients, and the elderly often experience only weakness, shortness of breath, nausea/ vomiting, and back or jaw pain.**
- **If patient has taken nitroglycerin without relief, consider potency of the medication.**
- **Contraindications to opioids include severe COPD and respiratory distress. Monitor the patient closely.**
- **Monitor for hypotension after administration of nitroglycerin and opioids.**
- **Allow the patient to be in their position of comfort to maximize their breathing effort.**
- **EMT may administer Nitroglycerin to patients already prescribed medication. May give from EMS supply.**
- **Agency medical director may require Contact of Medical Control.**



Adult Tachycardia

NARROW (≤ 0.11 sec)

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Most important goal is to differentiate the type of tachycardia and if STABLE or UNSTABLE and SYMPTOMATIC.**
- **12-Lead ECG:**
 - 12 Lead ECG not necessary to diagnose and treat
 - Obtain when patient is stable and/or following rhythm conversion.
- **Unstable condition**
 - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
 - If at any point patient becomes unstable move to unstable arm in algorithm.
- Search for underlying cause of tachycardia such as fever, sepsis, dyspnea, etc.
- Typical sinus tachycardia is in the range of 100 to (200 - patient's age) beats per minute.
- **Symptomatic condition**
 - Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
 - Symptomatic tachycardia usually occurs at rates ≥ 150 beats per minute.
 - Patients symptomatic with heart rates < 150 likely have impaired cardiac function such as CHF.
- **Serious Signs / Symptoms:**
 - Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute CHF.
- **If patient has history or 12 Lead ECG reveals Wolfe Parkinson White (WPW):**
 - DO NOT administer a Calcium Channel Blocker (e.g. Diltiazem) or Beta Blockers.
 - Use caution with Adenosine and give only with defibrillator available.
- **Regular Narrow-Complex Tachycardia:**
 - Vagal maneuvers and adenosine are preferred. Vagal maneuvers may convert 19% to 54 % of SVT.
 - Using passive leg raise with Valsalva is more effective.
 - Adenosine should be pushed rapidly via proximal IV site followed by 20 mL Normal Saline rapid flush.
 - Adenosine should not be used in the post-cardiac transplant patient without **Contact of Medical Control**.
 - Agencies using both calcium channel blockers and beta blockers should choose one primarily. Giving the agents sequentially requires **Contact of Medical Control**. This may lead to profound bradycardia / hypotension.
- **Irregular Narrow-Complex Tachycardia:**
 - Rate control is more important in pre-hospital setting rather than focus on rhythm conversion.
- **Synchronized Cardioversion:**
 - Recommended to treat UNSTABLE Atrial Fibrillation, Atrial Flutter and SVT.
- Monitor for hypotension after administration of Calcium Channel Blockers or Beta Blockers.
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.



Adult Monomorphic Tachycardia

Wide Complex (≥ 0.12 sec)

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Most important goal is to differentiate the type of tachycardia and if STABLE or UNSTABLE and SYMPTOMATIC.**
- **12-Lead ECG:**
 - 12 Lead ECG not necessary to diagnose and treat
 - Obtain when patient is stable and/or following rhythm conversion.
- **Monomorphic QRS:**
 - All QRS complexes in a single lead are similar in shape.
- **Polymorphic QRS:**
 - QRS complexes in a single lead will change shape from complex to complex.
- **Rhythm should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.**
- **Unstable condition**
 - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
 - If at any point patient becomes unstable move to unstable arm in algorithm.
- **Symptomatic condition**
 - Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
 - Symptomatic tachycardia usually occurs at rates ≥ 150 beats per minute. Patients symptomatic with heart rates < 150 likely have impaired cardiac function such as CHF.
- **Serious Signs / Symptoms:**
 - Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute congestive heart failure.
- Search for underlying cause of tachycardia such as fever, sepsis, dyspnea, etc.
- Typical sinus tachycardia is in the range of 100 to (220 – patients age) beats per minute.
- If patient has history or 12 Lead ECG reveals Wolfe Parkinson White (WPW), DO NOT administer a Calcium Channel Blocker (e.g., Diltiazem) or Beta Blockers. Use caution with Adenosine and give only with defibrillator available.
- **Regular Wide-Complex Tachycardia:**
 - **Unstable condition:**
 - Immediate defibrillation if pulseless and begin CPR.
 - **Stable condition:**
 - Typically VT or SVT with aberrancy. Adenosine may be given if regular and monomorphic and if defibrillator available.
 - Verapamil contraindicated in wide-complex tachycardias.
 - Agencies using Amiodarone, Procainamide and Lidocaine need choose one agent primarily. Giving multiple anti-arrhythmics requires contact of Medical Control.
 - Atrial arrhythmias with WPW should be treated with Amiodarone or Procainamide
- **Irregular Tachycardia:**
 - Wide-complex, irregular tachycardia: Do not administer calcium channel, beta blockers, or adenosine as this may cause paradoxical increase in ventricular rate. This will usually require cardioversion. Contact Medical Control.
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.



Adult Polymorphic Tachycardia

WIDE (≥ 0.12 sec) Torsades de pointes

History

- Age
- Past medical history (MI, Angina, Diabetes, post menopausal)
- Recent physical exertion
- Palpitations, irregular heart beat
- Time (onset /duration / repetition)

Signs and Symptoms

- Chest pain, heart failure, dyspnea
- AMS
- Shock, poor perfusion, hypotension
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting, dizziness

Differential

- Cardiac arrest
- Sinus Tachycardia vs. dysrhythmia
- Fever, sepsis, infection
- Pericarditis, pulmonary embolism
- Aortic dissection or aneurysm
- Overdose

**Assess tachycardia in context of clinical condition
Identify and treat underlying cause of tachycardia**

Unstable / Serious Signs and Symptoms
HR Typically ≥ 150
Hypotension, Acute AMS, Ischemic Chest Pain,
Acute CHF, Seizures, Syncope, or Shock
secondary to tachycardia

Defibrillation Procedure	
P	

B	12 Lead ECG Procedure
P	Cardiac Monitor
	IV or IO Access Protocol UP 6

Pulse Present?

Consider consultation with medical control

Exit to
Cardiac Arrest
Protocol AC 3

P	
	Monitor and Reassess

P	
	Monitor and Reassess

**Notify Destination or
Contact Medical Control**

Polymorphic QRS:
• QRS complexes in a single lead will change shape from complex to complex.



Adult Polymorphic Tachycardia

WIDE (≥ 0.12 sec) Torsades de pointes

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
 - **Most important goal is to differentiate the type of tachycardia and if STABLE or UNSTABLE and SYMPTOMATIC.**
 - **12-Lead ECG:**
 - 12 Lead ECG not necessary to diagnose and treat
 - Obtain when patient is stable and/or following rhythm conversion.
 - **Monomorphic QRS:**
 - All QRS complexes in a single lead are similar in shape.
 - **Polymorphic QRS:**
 - QRS complexes in a single lead will change shape from complex to complex.
 - **Rhythm should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.**
 - **Unstable condition**
 - Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
 - If at any point patient becomes unstable move to unstable arm in algorithm.
 - **Symptomatic condition**
 - Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.
 - Symptomatic tachycardia usually occurs at rates ≥ 150 beats per minute. Patients symptomatic with heart rates < 150 likely have impaired cardiac function such as CHF.
 - **Serious Signs / Symptoms:**
 - Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute congestive heart failure.
 - Search for underlying cause of tachycardia such as fever, sepsis, dyspnea, etc.
 - Typical sinus tachycardia is in the range of 100 to (220 – patients age) beats per minute.
 - If patient has history or 12 Lead ECG reveals Wolfe Parkinson White (WPW), DO NOT administer a Calcium Channel Blocker (e.g., Diltiazem) or Beta Blockers. Use caution with Adenosine and give only with defibrillator available.
 - **Polymorphic / Irregular Tachycardia:**
 - This situation is usually unstable and immediate defibrillation is warranted.
 - If QT length is known, use for decision-making. Prolonged QT length defined as > 500 msec.
 - QT length < 500 msec:
 - Arrhythmia more likely related to ischemia or infarction and Magnesium not likely helpful.
 - May quickly deteriorate into Ventricular Fibrillation.
 - Even when terminated by defibrillation, may recur, so follow with medication therapy.
 - QT prolongation > 500 msec:
 - Magnesium more likely to be helpful.
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.



Ventricular Fibrillation Pulseless Ventricular Tachycardia

Cardiac Arrest Protocol AC 3

	<p>Begin Continuous CPR Compressions Push Hard (≥ 2 inches) Push Fast (100 - 120 / min) Change Compressors every 2 minutes <i>(sooner if fatigued)</i> <i>(Limit changes / pulse checks ≤ 10 seconds)</i></p> <p>Ventilate 1 breath every 6 seconds 30:2 Compression:Ventilation if no Advanced Airway Monitor EtCO2 <i>if available</i></p> <p>AED Procedure <i>if available</i></p>
P	Defibrillation Procedure
	IV / IO Access Protocol UP 6
A	<p>Epinephrine (1:10,000) 1 mg IV / IO Repeat every 5 minutes <i>If VF / VT refractory to defibrillation, delay Epinephrine administration until after 2d defibrillation</i></p>
	Search for Reversible Causes
	<p>Continue CPR Compressions Push Hard (≥ 2 inches) Push Fast (100 - 120 / min) Change Compressors every 2 minutes <i>(sooner if fatigued)</i> <i>(Limit changes / pulse checks ≤ 10 seconds)</i></p> <p><u>If Rhythm Refractory</u> Continue CPR and give Agency specific Anti-arrhythmics and Epinephrine Continue CPR up to point where you are ready to defibrillate with device charged. Repeat pattern during resuscitation.</p>
P	
	<p>Defibrillation Procedure <i>If VF / VT refractory after 3 shocks consider Dual or Double Defibrillation</i></p>

AT ANY TIME

Return of Spontaneous Circulation

Go to Post Resuscitation Protocol AC 10

- Reversible Causes**
- Hypovolemia
 - Hypoxia
 - Hydrogen ion (acidosis)
 - Hypothermia
 - Hypo / Hyperkalemia

 - Tension pneumothorax
 - Tamponade; cardiac
 - Toxins
 - Thrombosis; pulmonary (PE)
 - Thrombosis; coronary (MI)

Adult Cardiac Protocol Section


Notify Destination or Contact Medical Control



Ventricular Fibrillation Pulseless Ventricular Tachycardia

Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional Team Focused CPR Protocol AC 11 or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.**
- **DO NOT HYPERVENTILATE: If no advanced airway (BIAD, ETT) compression to ventilation ratio is 30:2. If advanced airway in place, ventilate 10 breaths per minute with continuous, uninterrupted compressions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **Passive oxygenation optional in agencies practicing Team Focused Approach / Pit-Crew Approach.**
- **Reassess and document BIAD and / or endotracheal tube placement and EtCO₂ frequently, after every move, and at transfer of care.**
- **IV / IO access and drug delivery is secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Defibrillation:**
 - Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
 - Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.
 - Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- **End Tidal CO₂ (EtCO₂)**
 - If EtCO₂ is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
 - If EtCO₂ spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **Special Considerations**
 - Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm. Defibrillation is safe at all energy levels.
 - Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
 - Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol TE 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
 - Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Magnesium Sulfate is not routinely recommended during cardiac arrest, but may help with Torsades de points, prolonged QT, low Magnesium States (malnourished / alcoholic), and suspected digitalis toxicity**
- Return of spontaneous circulation: Heart rate should be > 60 when initiating anti-arrhythmic infusions.
- Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.
- Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment options.



Transport Destination Decision
 Post-resuscitation patient is medically complex.

Consider facility capabilities:

- 24-hour cardiac catheterization laboratory
- Medical ICU service
- Cardiology service
- Neurology service
- Pulmonology service
- Targeted Temperature Management

Reversible Causes

Hypovolemia
 Hypoxia
 Hydrogen ion (acidosis)
 Hypothermia
 Hypo / Hyperkalemia

Tension pneumothorax
 Tamponade; cardiac
 Toxins
 Thrombosis; pulmonary (PE)
 Thrombosis; coronary (MI)

Arrhythmias are common and usually self limiting after ROSC

If Arrhythmia Persists follow Rhythm Appropriate Protocol



Post Resuscitation

Pearls

- **Recommended Exam: Mental Status, Neck, Skin, Lungs, Heart, Abdomen, Extremities, Neuro**
- **Continue to search for potential cause of cardiac arrest during post-resuscitation care.**
- **Hyperventilation is a significant cause of hypotension and recurrence of cardiac arrest in the post resuscitation phase and must be avoided. Titrate FiO₂ to maintain SpO₂ of 92 - 98%.**
- **Pain/sedation:**

Patients requiring advanced airways and ventilation commonly experience pain and anxiety. Unrelieved pain can lead to increased catecholamine release, ischemia, immunosuppression, and prolonged hospitalization.

Ventilated patients cannot communicate pain / anxiety and providers are poor at recognizing pain / anxiety.

Vital signs such as tachycardia and / or hypertension can provide clues to inadequate sedation, however they both are not always reliable indicators of patient's lack of adequate sedation.

Pain must be addressed first, before anxiety. Opioids are typically the first line agents before benzodiazepines. Ketamine is also a reasonable first choice agent.
- **Ventilator / Ventilation strategies:**

Tailored to individual patient presentations. Medical Control can indicate different strategies above. In general ventilation with BVM should cause chest rise. With mechanical ventilation a reasonable tidal volume should be about 6 mL/kg and peak pressures should be < 30 cmH₂O.

Continuous pulse oximetry and capnography should be maintained during transport for monitoring. Head of bed should be maintained at least 10 – 20 degrees of elevation when possible to decrease aspiration risk.
- **EtCO₂ Monitoring:**

Initial End tidal CO₂ may be elevated immediately post-resuscitation, but will usually normalize. Goal is 35 – 45 mmHg but avoid hyperventilation to achieve.
- **Titrate fluid resuscitation and vasopressor administration to maintain SBP of 90 – 100 mmHg or Mean Arterial Pressure (MAP) of 65 – 80 mmHg.**
- **STEMI (ST-Elevation Myocardial Infarction)**

Consider placing 2 IV sites in the left arm: Many PCI centers use the right radial artery for intervention. Consider placing defibrillator pads on patient as a precaution. Document and time-stamp facility STEMI notification and make notification as soon as possible. Document the time of the 12-Lead ECG in the PCR as a Procedure along with the interpretation (Paramedic).
- **Consider transport to facility capable of managing the post-arrest patient including hypothermia therapy, cardiology / cardiac catheterization, intensive care service, and neurology services.**
- **Targeted Temperature Management (optional):**

Maintain core temperature between 32 - 36°C. Infusion of cold saline is NOT recommended in the prehospital setting. No evidence suggests improved survival with prehospital cooling.
- **The condition of post-resuscitation patients fluctuates rapidly and continuously, and they require close monitoring. Appropriate post-resuscitation management may best be planned in consultation with Medical Control.**



Team Focused CPR (Optional)

Criteria for Death / No Resuscitation
Review DNR / MOST Form

YES

NO

AT ANY TIME

Return of Spontaneous Circulation



Go to Post Resuscitation Protocol AC 10

Begin Continuous CPR Compressions
Push Hard (≥ 2 inches)
Push Fast (100 - 120 / min)
Change Compressors every 2 minutes (sooner if fatigued)
(Limit changes / pulse checks ≤ 10 seconds)

Ventilate 1 breath every 6 seconds
30:2 Compression:Ventilation if no Advanced Airway
Monitor EtCO2 if available

First Arriving BLS / ALS Responder
Initiate Compressions Only CPR
Initiate Defibrillation Automated Procedure *if available*
Call for additional resources

Second Arriving BLS / ALS Responder
Assume Compressions or
Initiate Defibrillation Automated / Manual Procedure
Place BIAD
DO NOT Interrupt Compressions
Ventilate at 6 to 8 breaths per minute

Decomposition
Rigor mortis
Dependent lividity
Blunt force trauma
Injury incompatible with life
Extended downtime with asystole

Do not begin resuscitation

Follow Deceased Subjects Policy

BLS

Third Arriving Responder
BLS or ALS

ALS

Establish Team Leader (Hierarchy)
Fire Department or Squad Officer
EMT
First Arriving Responder

Rotate with Compressor
To prevent Fatigue and effect high quality compressions
Take direction from Team Leader

Fourth / Subsequent Arriving Responders
Take direction from Team Leader



Continue Cardiac Arrest Protocol AC 3

Establish Team Leader (Hierarchy)
EMS ALS Personnel
Fire Department or Squad Officer
EMT
First Arriving Responder

A

Initiate Defibrillation Automated Procedure
Establish IV / IO Protocol UP 6
Administer Appropriate Medications
Establish Airway with BIAD if not in place

P

Initiate Defibrillation Manual Procedure
Continuous Cardiac Monitoring
Establish IV / IO Protocol UP 6
Administer Appropriate Medications
Establish Airway with BIAD if not in place



Continue Cardiac Arrest Protocol AC 3

Team Leader
ALS Personnel
Responsible for patient care
Responsible for briefing / counseling family

Incident Commander
Fire Department / First Responder Officer
Team Leader until ALS arrival
Manages Scene / Bystanders
Ensures high-quality compressions
Ensures frequent compressor change
Responsible for briefing family prior to ALS arrival

Adult Cardiac Protocol Section




Team Focused CPR (Optional)

Pearls

- This protocol is optional and given only as an example. Agencies may and are encouraged to develop their own.
- Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional protocol or development of local agency protocol.
- Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated.
- **DO NOT HYPERVENTILATE:** If no advanced airway (BIAD, ETT), compression to ventilation ratio is 30:2. If advanced airway in place, ventilate 10 breaths per minute with continuous, uninterrupted compressions.
- Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.
- Passive oxygenation optional in agencies practicing Team Focused Approach / Pit-Crew Approach.
- Reassess and document BIAD and / or endotracheal tube placement and EtCO₂ frequently, after every move, and at transfer of care.
- IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.
- IV access is preferred route. Follow IV or IO Access Protocol UP 6.
- **Defibrillation:** Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
 - Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.
 - Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.
- Discussion with Medical Control can be a valuable tool in developing a differential diagnosis and identifying possible treatment options.



On Scene Resuscitation Termination of CPR (Optional)





On Scene Resuscitation / Termination of CPR

Pearls

- **General approach:**
 1. Determine if a terminal disease is involved?
 2. Is there an advanced directive such as a DNR / MOST form?
 3. Did the patient express to your historian any desires regarding resuscitation and if so what measures?
 4. Remember a living will is not a DNR.
- Obtain a history while resuscitation efforts are ongoing. Determine the most legitimate person on scene as your information source such as a spouse, child, or sibling or Durable Health Care Power of Attorney.
- Basic and Advanced Life Support may use for treatment decisions.



Mechanical Circulatory Support LVAD, RVAD, and Bi-VAD

History

- SAMPLE
- Bridge to transplant
- Destination therapy
- Estimated downtime
- LVAD, RVAD, Bi-Vad, TAH
- DNR, MOST, or Living Will
- Contact with LVAD coordinator

Signs and Symptoms


- Unconsciousness
- Pulseless
- Apneic
- Poor capillary refill / skin color
- AMS or decreased mental status
- No electrical activity on ECG
- No heart tones on auscultation

Differential

- See Reversible Causes below
- Infection/Sepsis
- Hypovolemia
- Cardiac arrest
- Hemorrhage

Contact VAD coordinator:

- As quickly as possible for troubleshooting and treatment advice, but do not delay emergency treatment
- Follow patient specific emergency plan if present



Adult Cardiac Protocol Section



Mechanical Circulatory Support LVAD, RVAD, and Bi-VAD

History


- SAMPLE
- Bridge to transplant
- Destination therapy
- Estimated downtime
- LVAD, RVAD, Bi-Vad, TAH
- DNR, MOST, or Living Will
- Contact with LVAD coordinator

Signs and Symptoms

- Unconsciousness
- Pulseless
- Apneic
- Poor capillary refill / skin color
- AMS or decreased mental status
- No electrical activity on ECG
- No heart tones on auscultation


Differential

- See Reversible Causes below
- Infection/Sepsis
- Hypovolemia
- Cardiac arrest
- Hemorrhage





Mechanical Circulatory Support LVAD, RVAD, and Bi-VAD



- Reversible Causes**
- Hypovolemia
 - Hypoxia
 - Hydrogen ion (acidosis)
 - Hypothermia
 - Hypo / Hyperkalemia
 - Tension pneumothorax
 - Tamponade; cardiac
 - Toxins
 - Thrombosis; pulmonary (PE)
 - Thrombosis; coronary (MI)
- Suspected Opioid Overdose**
- Administer Naloxone per Overdose / Toxic Ingestion Protocol TE 7



Mechanical Circulatory Support LVAD, RVAD, and Bi-VAD

Pearls

- **Recommended exam: Mental status, skin color, capillary refill, peripheral pulses, blood pressure.**
- **Assessment of blood flow and perfusion status:**
 - **Optimal BP attained by manual BP and Doppler.**
 - **Automated BP devices can measure a BP in about 50% of attempts and is not reliable to assess perfusion**
 - **A MAP of ≥ 60 mmHg is adequate for most LVAD patients.**
 - **Skin color, skin temperature, capillary refill**
- **Mechanical Circulatory Support devices:**
 - **LVAD – Left Ventricular Assist Device**
 - **RVAD – Right Ventricular Assist Device**
 - **BiVAD – Biventricular Ventricular Assist Device**
 - **TAH – Total Artificial Heart**
- **Reasons for use:**
 - **Bridge therapy – patients awaiting transplant or anticipated recovery.**
 - **Destination therapy – advanced heart failure, not candidate for transplant, and will live rest of life with device.**
- **Pump type and assessing pulses:**
 - **Pulsatile flow pumps – older units, not commonly in use now, but generate blood flow with a pulsatile flow and patient will have a palpable pulse.**
 - **Continuous flow pumps – majority of pumps now used and create blood flow in a continuous stream, no pulsatile flow, so patient will not have a palpable pulse.**
 - **Most devices are implanted inside the chest and have an internal pump, a driveline connected from the pump to the controller unit, and a power source consisting of batteries and electrical cord for receptacles.**
- **Common complications:**
 - **Disconnection of power supply, either battery disconnect, or electrical cord to receptacle disconnection.**
 - **Driveline failure or disconnection from controller unit.**
 - **Controller failure**
 - **Blood clot formation, acute stroke, and bleeding (mucosal and gastrointestinal most common sites)**
 - **Infection**
- **Abnormal heart rhythm:**
 - **Pseudo-PEA: Normal cardiac electrical activity in a patient who is alert and well perfused with no palpable pulse.**
 - **Tachyarrhythmias are usually well tolerated.**
- **End Tidal CO₂ (EtCO₂)**
 - **If EtCO₂ is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.**
 - **If EtCO₂ spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)**
- **Transcutaneous Pacing:**
 - **Pacing is NOT effective in cardiac arrest and pacing in cardiac arrest does NOT increase chance of survival**



Allergic Reaction / Anaphylaxis

History

- Onset and location
- Insect sting or bite
- Food allergy / exposure
- Medication allergy / exposure
- New clothing, soap, detergent
- Past history of reactions
- Past medical history
- Medication history

Signs and Symptoms

- Itching or hives
- Coughing / wheezing or respiratory distress
- Chest or throat constriction
- Difficulty swallowing
- Hypotension or shock
- Edema
- N/V

Differential

- Urticaria (rash only)
- Anaphylaxis (systemic effect)
- Shock (vascular effect)
- Angioedema (drug induced)
- Aspiration / Airway obstruction
- Vasovagal event
- Asthma or COPD
- CHF

Assess Symptom Severity / Suspected Exposure to Allergen

MILD
Skin Only

A	Diphenhydramine 25 - 50 mg PO
	IV Procedure <i>if indicated</i>
	Diphenhydramine 25 - 50 mg PO / IV / IM / IO

B	Monitor and Reassess Monitor for Worsening Signs and Symptoms
---	---

MODERATE
2 + Body Systems

	Epinephrine 1:1000 IM 0.3 - 0.5 mg <i>if available</i>
	Diphenhydramine 25 - 50 mg PO <i>See Pearls</i>
B	Albuterol Nebulizer 2.5mg or Levabuterol 1.25mg Repeat as needed x 3 <i>if indicated</i>
A	Epinephrine 1:1000 0.3 - 0.5 mg IM Repeat every 5 minutes if no improvement

A	Diphenhydramine 25 - 50 mg IV / IM / IO <i>if not given PO (See Pearls)</i>
	IV / IO Procedure
	Albuterol Nebulizer 2.5mg or Levabuterol 1.25mg +/- Ipratropium 0.5 mg Repeat as needed x 3 <i>if indicated</i>
P	

SEVERE
2 + Body Systems + hypotension
Or Isolated Hypotension

	Epinephrine 1:1000 IM 0.3 - 0.5 mg <i>if available</i>
	B Albuterol 2.5mg Nebulizer or Levabuterol 1.25mg Repeat as needed x 3 <i>if indicated</i>
	Airway Protocol(s) AR 1 - 4 <i>if indicated</i>
	Hypotension / Shock Protocol AM 5 <i>if indicated</i>

Notify Destination or Contact Medical Control

Adult Medical Protocol Section



Allergic Reaction / Anaphylaxis

Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Abdominal**
- **Anaphylaxis is an acute and potentially lethal multisystem allergic reaction.**
- **Epinephrine administration:**
 - **Drug of choice and the FIRST drug that should be administered in acute anaphylaxis (Moderate / Severe Symptoms.) IM Epinephrine should be administered in priority before or during attempts at IV or IO access.**
 - **Diphenhydramine and steroids have no proven utility in Moderate / Severe anaphylaxis and may be given only After Epinephrine. Diphenhydramine and steroids should NOT delay repeated Epinephrine administration.**
 - **In Moderate and Severe anaphylaxis Diphenhydramine may decrease mental status. Oral Diphenhydramine should NOT be given to a patient with decreased mental status and / or a hypotensive patient as this may cause nausea and / or vomiting.**
- **Anaphylaxis unresponsive to repeat doses of IM epinephrine may require IV epinephrine administration by IV push or epinephrine infusion. Contact Medical Control for appropriate dosing.**
- **Symptom Severity Classification:**
 - **Mild symptoms:**
 - **Flushing, hives, itching, erythema with normal blood pressure and perfusion.**
 - **Moderate symptoms:**
 - **Flushing, hives, itching, erythema plus respiratory (wheezing, dyspnea, hypoxia) or gastrointestinal symptoms (nausea, vomiting, abdominal pain) with normal blood pressure and perfusion.**
 - **Severe symptoms:**
 - **Flushing, hives, itching, erythema plus respiratory (wheezing, dyspnea, hypoxia) or gastrointestinal symptoms (nausea, vomiting, abdominal pain) with hypotension/poor perfusion or isolated hypotension.**
- **Allergic reactions may occur with only respiratory and gastrointestinal symptoms and have no rash / skin involvement.**
- **Angioedema** is seen in moderate to severe reactions and is swelling involving the face, lips or airway structures. This can also be seen in patients taking blood pressure medications like Prinivil / Zestril (lisinopril)-typically end in **-il**.
- **Hereditary Angioedema** involves swelling of the face, lips, airway structures, extremities, and may cause moderate to severe abdominal pain. Some patients are prescribed specific medications to aid in reversal of swelling. **Paramedic may assist or administer this medication per patient / package instructions.**
- **12 lead ECG and cardiac monitoring should NOT delay administration of epinephrine.**
- **EMR / EMT may administer Epinephrine IM and may administer from EMS supply.** Agency Medical Director may require contact of medical control prior to EMR / EMT administering any medication.
- **EMR / EMT may administer Epinephrine IM via AutoInjector or manual draw-up** per Agency Medical Director.
- **EMR may administer diphenhydramine by oral route only and may administer from EMS supply.** Agency Medical Director may require contact of medical control prior to EMT / EMR administering any medication.
- **EMT may administer Albuterol if patient already prescribed and may administer from EMS supply.** Agency Medical Director may require contact of medical control prior to EMT / EMR administering any medication.
- The shorter the onset from exposure to symptoms the more severe the reaction.



Diabetic; Adult

History

- Past medical history
- Medications
- Recent blood glucose check
- Last meal



Signs and Symptoms

- Altered mental status
- Combative / irritable
- Diaphoresis
- Seizures
- Abdominal pain
- Nausea / vomiting
- Weakness
- Dehydration
- Deep / rapid breathing

Differential

- Alcohol / drug use
- Toxic ingestion
- Trauma; head injury
- Seizure
- CVA
- Altered baseline mental status

	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure <i>if indicated</i>
	IV / IO Protocol UP 6
P	Cardiac Monitor
	Altered Mental Status Protocol UP 4 <i>if indicated</i>
	Hypotension / Shock Protocol AM 5 <i>if indicated</i>
	Suspected Stroke Protocol UP 14 <i>if indicated</i>
	Seizure Protocol UP 13 <i>if indicated</i>





Diabetic; Adult

Pearls

- **Recommended exam: Mental Status, Skin, Respirations and effort, Neuro.**
- **Patients with prolonged hypoglycemia may not respond to glucagon.**
- **Do not administer oral glucose to patients that are not able to swallow or protect their airway.**
- **Quality control checks should be maintained per manufacturers recommendation for all glucometers.**
- **Patient's refusing transport to medical facility after treatment of hypoglycemia:**
 - Blood sugar must be ≥ 80 , patient has ability to eat and availability of food with responders on scene.
 - Patient must have known history of diabetes and not taking any oral diabetic agents.
 - Patient returns to normal mental status and has a normal neurological exam with no new neurological deficits.
 - Must demonstrate capacity to make informed health care decisions. See Universal Patient Care Protocol UP 1.
 - Otherwise contact medical control.
- **Hypoglycemia with Oral Agents:**
 - Patient's taking oral diabetic medications should be encouraged to allow transportation to a medical facility.
 - They are at risk of recurrent hypoglycemia that can be delayed for hours and require close monitoring even after normal blood glucose is established.
 - Not all oral agents have prolonged action so Contact Medical Control for advice.
 - Patient's who meet criteria to refuse care should be instructed to contact their physician immediately and consume a meal.
- **Hypoglycemia with Insulin Agents:**
 - Many forms of insulin now exist. Longer acting insulin places the patient at risk of recurrent hypoglycemia even after a normal blood glucose is established.
 - Not all insulin have prolonged action so Contact Medical Control for advice.
 - Patient's who meet criteria to refuse care should be instructed to contact their physician immediately and consume a meal.
- **Congestive Heart Failure patients who have Blood Glucose > 250 :**
 - Limit fluid boluses unless they have signs of volume depletion, dehydration, poor perfusion, hypotension, and/or shock.
- Contact medical control for advice.



Dialysis / Renal Failure

History


- Peritoneal or Hemodialysis
- Anemia
- Catheter access noted
- Shunt access noted
- Hyperkalemia

Signs and Symptoms

- Hypotension
- Bleeding
- Fever
- Electrolyte imbalance
- Nausea and / or vomiting
- Altered Mental Status
- Seizure
- Arrhythmia

Differential

- Congestive heart failure
- Pericarditis
- Diabetic emergency
- Sepsis
- Cardiac tamponade





Dialysis / Renal Failure

Pearls

- **Recommended exam: Mental status. Neurological. Lungs. Heart.**
- **Consider transport to medical facility capable of providing Dialysis treatment.**
- **Do not take Blood Pressure or start IV in extremity which has a shunt / fistula in place.**
- **Access of shunt indicated in the dead or near-dead patient only with no IV or IO access.**
- **If hemorrhage cannot be controlled with firm, uninterrupted direct pressure, application of tourniquet with uncontrolled dialysis fistula bleeding is indicated.**
- **Hemodialysis:**
 - Process which removes waste from the blood stream and occurs about three times each week.
 - Some patients do perform hemodialysis at home.
- **Peritoneal dialysis:**
 - If patient complains of fever, abdominal pain, and / or back pain, bring the PD fluid bag, which has drained from the abdomen, to the hospital.

Complications of Dialysis Treatment:

Hypotension:

Typically responds to small fluid bolus of 250 mL Normal Saline. May result in angina, AMS, seizure or arrhythmia.

Filtration and decreased blood levels of some medications like some seizure medications:

Disequilibrium syndrome:

Shift of metabolic waste and electrolytes causing weakness, dizziness, nausea and / or vomiting and seizures.

Equipment malfunction:

Air embolism.
Bleeding.
Electrolyte imbalance.
Fever.

- **Fever:**
 - Consider sepsis in a dialysis patient with any catheter extending outside the body.
- Always consider Hyperkalemia in all dialysis or renal failure patients.
- Sodium Bicarbonate and Calcium Chloride / Gluconate should not be mixed. Ideally give in separate lines.
- Renal dialysis patients have numerous medical problems typically. Hypertension and cardiac disease are prevalent.



Hypertension

History

- Documented Hypertension
- Related diseases: Diabetes; CVA; Renal Failure; Cardiac Problems
- Medications for Hypertension
- Compliance with Hypertensive Medications
- Erectile Dysfunction medications
- Pregnancy

Signs and Symptoms

One of these

- Systolic BP 220 or greater
- Diastolic BP 120 or greater


AND at least one of these

- Headache
- Chest Pain
- Dyspnea
- Altered Mental Status
- Seizure

Differential

- Hypertensive encephalopathy
- Primary CNS Injury
Cushing's Response with Bradycardia and Hypertension
- Myocardial Infarction
- Aortic Dissection / Aneurysm
- Pre-eclampsia / Eclampsia

Hypertension is not uncommon especially in an emergency setting. Hypertension is usually transient and in response to stress and / or pain. A hypertensive emergency is based on blood pressure along with symptoms which suggest an organ is suffering damage such as MI, CVA or renal failure. This is very difficult to determine in the pre-hospital setting in most cases. Aggressive treatment of hypertension can result in harm. Most patients, even with significant elevation in blood pressure, need only supportive care. Specific complaints such as chest pain, dyspnea, pulmonary edema or altered mental status should be treated based on specific protocols and consultation with Medical Control.



Adult Medical Protocol Section

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- Elevated blood pressure is based on two to three sets of vital signs.
- Symptomatic hypertension is typically revealed through end organ dysfunction to the cardiac, CNS or renal systems.
- All symptomatic patients with hypertension should be transported with their head elevated at 30 degrees.
- Ensure appropriate size blood pressure cuff utilized for body habitus.



Hypotension / Shock

History

- Blood loss - vaginal or gastrointestinal bleeding, AAA, ectopic
- Fluid loss - vomiting, diarrhea, fever
- Infection
- Cardiac ischemia (MI, CHF)
- Medications
- Allergic reaction
- Pregnancy
- History of poor oral intake

Signs and Symptoms


- Restlessness, confusion
- Weakness, dizziness
- Weak, rapid pulse
- Pale, cool, clammy skin
- Delayed capillary refill
- Hypotension
- Coffee-ground emesis
- Tarry stools

Differential

- Ectopic pregnancy
- Dysrhythmias
- Pulmonary embolus
- Tension pneumothorax
- Medication effect / overdose
- Vasovagal
- Physiologic (pregnancy)
- Sepsis

	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure
A	IV / IO Procedure
P	Cardiac Monitor
	Airway Protocol(s) <i>if indicated</i>
	Diabetic Protocol AM 2 <i>if indicated</i>

History and Exam Suggest Type of Shock



Adult Medical Protocol Section



Hypotension / Shock

Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- Hypotension can be defined as a systolic blood pressure of less than 90. This is not always reliable and should be interpreted in context and patients typical BP if known. Shock may be present with a normal blood pressure initially.
- Shock often is present with normal vital signs and may develop insidiously. Tachycardia may be the only manifestation.
- Consider all possible causes of shock and treat per appropriate protocol.
- For non-cardiac, non-trauma hypotension, consider Dopamine when hypotension unresponsive to fluid resuscitation.
- **Hypovolemic Shock:**
Hemorrhage, trauma, GI bleeding, ruptured aortic aneurysm or pregnancy-related bleeding.
Tranexamic Acid (TXA):
Agencies utilizing TXA must have approval from your T-RAC.
- **Cardiogenic Shock:**
Heart failure: MI, Cardiomyopathy, Myocardial contusion, Ruptured ventricular / septum / valve / toxins.
- **Distributive Shock:**
Sepsis
Anaphylactic
Neurogenic: Hallmark is warm, dry, pink skin with normal capillary refill time and typically alert.
Toxins
- **Obstructive Shock:**
Pericardial tamponade. Pulmonary embolus. Tension pneumothorax.
Signs may include hypotension with distended neck veins, tachycardia, unilateral decreased breath sounds or muffled heart sounds.
- **Acute Adrenal Insufficiency or Congenital Adrenal Hyperplasia:**
Body cannot produce enough steroids (glucocorticoids / mineralocorticoids.) May have primary or secondary adrenal disease, congenital adrenal hyperplasia, or more commonly have stopped a steroid like prednisone. Injury or illness may precipitate. Usually hypotensive with nausea, vomiting, dehydration and / or abdominal pain. **If suspected Paramedic should give Methylprednisolone 125 mg IM / IV / IO or Dexamethasone 10 mg IM / IV / IO. Use steroid agent specific to your drug list. May administer prescribed steroid carried by patient IM / IV / IO. Patient may have Hydrocortisone (Cortef or Solu-Cortef). Dose: < 1y.o. give 25 mg, 1-12 y.o. give 50 mg, and > 12 y.o. give 100 mg or dose specified by patient's physician.**



Childbirth / Labor

History


- Due date
- Time contractions started / how often
- Rupture of membranes
- Time / amount of any vaginal bleeding
- Sensation of fetal activity
- Past medical and delivery history
- Medications
- Gravida / Para Status
- High Risk pregnancy

Signs and Symptoms

- Spasmodic pain
- Vaginal discharge or bleeding
- Crowning or urge to push
- Meconium

Differential

- Abnormal presentation
 - Buttock
 - Foot
 - Hand
- Prolapsed cord
- Placenta previa
- Abruptio placenta



Adult Obstetrical Protocol Section

Delivery

Go to Newly Born Protocol AO 2



Childbirth / Labor

Pearls

- **Recommended Exam (of Mother): Mental Status, Heart, Lungs, Abdomen, Neuro**
- **Record APGAR at 1 minute and 5 minutes after birth.**
- **After delivery, massaging the uterus (lower abdomen) will promote uterine contraction and help to control post-partum bleeding.**
- Document all times (delivery, contraction frequency, and length).
- **Transport or Delivery?**
Decision to transport versus remain and deliver is multifactorial and difficult. Generally it is preferable to transport. Factors that will impact decision include: number of previous deliveries; length of previous labors; frequency of contractions; urge to push; and presence of crowning.
- **Maternal positioning for labor:**
Supine with head flat or elevated per mother's choice. Maintain flexion of both knees and hips. Elevated buttocks slightly with towel. If delivery not imminent, place mother in the left, lateral recumbent position with right side up about 10 – 20°.
- **Umbilical cord clamping and cutting:**
Place first clamp about 10 cm from infant's abdomen and second clamp about 5 cm away from first clamp.
- **Multiple Births:**
Twins occur about 1/90 births. Typically manage the same as single gestation. If imminent delivery call for additional resources, if needed. Most twins deliver at about 34 weeks so lower birth weight and hypothermia are common. Twins may share a placenta so clamp and cut umbilical cord after first delivery. Notify receiving facility immediately.
- If maternal seizures occur, refer to the Obstetrical Emergencies Protocol.
- Some perineal bleeding is normal with any childbirth. Large quantities of blood or free bleeding are abnormal.



Newly Born

History


- Due date and gestational age
- Multiple gestation (twins etc.)
- Meconium / Delivery difficulties
- Congenital disease
- Medications (maternal)
- Maternal risk factors such as substance abuse or smoking

Signs and Symptoms

- Respiratory distress
- Peripheral cyanosis or mottling (normal)
- Central cyanosis (abnormal)
- Altered level of responsiveness
- Bradycardia

Differential

- Airway failure
 - Secretions
 - Respiratory drive
- Infection
- Maternal medication effect
- Hypovolemia, Hypoglycemia, Hypothermia
- Congenital heart disease





Newly Born

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Neck, Chest, Heart, Abdomen, Extremities, Neuro**
- **Document 1 and 5 minute Apgars in PCR**
- **Most newborns requiring resuscitation respond to ventilations / BVM, compressions, and/or epinephrine. If infant not responding consider hypovolemia, pneumothorax, and/or hypoglycemia (< 40 mg/dL).**
- **Term gestation, strong cry / breathing and with good muscle tone generally will need no resuscitation. Routine suctioning is no longer recommended.**
- **Most important vital signs in the newly born are respirations / respiratory effort and heart rate.**
- **Maintain warmth of infant following delivery; cap, plastic wrap, thermal mattress, radiant heat.**
- **Meconium staining:**
 - **Infant born through meconium staining who is not vigorous: Positive pressure ventilation is recommended, direct endotracheal suctioning is no longer recommended.**
- **Expected Pulse Oximetry readings immediately following birth:**

1 minute	60 – 65%
2 minutes	65 – 70%
3 minutes	70 – 75%
4 minutes	75 – 80%
5 minutes	80 – 85%
10 minutes	85 – 95%
- **Heart rate is critical during the first few moments of life and is best assessed by 3-lead ECG.**
- **Pulse oximetry should be applied to the right upper arm, wrist, or palm.**
- **CPR in infants is 120 compressions/minute with a 3:1 compression to ventilation ratio. 2-thumbs encircling chest and supporting the back is recommended. Limit interruptions of chest compressions.**
- **Maternal sedation or narcotics will sedate infant (Naloxone NO LONGER recommended-supportive care only).**
- **D10 = D50 diluted (1 ml of D50 with 4 ml of Normal Saline)**



Obstetrical Emergency

History


- Past medical history
- Hypertension meds
- Prenatal care
- Prior pregnancies / births
- Gravida / Para

Signs and Symptoms

- Vaginal bleeding
- Abdominal pain
- Seizures
- Hypertension
- Severe headache
- Visual changes
- Edema of hands and face

Differential

- Pre-eclampsia / Eclampsia
- Placenta previa
- Placenta abruptio
- Spontaneous abortion





Obstetrical Emergency

Pearls

- **Recommended Exam: Mental Status, Abdomen, Heart, Lungs, Neuro**
- **Midazolam 5 – 10 mg IM is effective in termination of seizures. Do not delay IM administration with difficult IV or IO access.**
- **Magnesium Sulfate should be administered as quickly as possible. May cause hypotension and decreased respiratory drive, but typically in doses higher than 6 g.**
- **Any pregnant patient involved in a MVC should be seen immediately by a physician for evaluation. Greater than 20 weeks generally require 4 to 6 hours of fetal monitoring. DO NOT suggest the patient needs an ultrasound.**
- **Ectopic pregnancy:**
Implantation of fertilized egg outside the uterus, commonly in or on the fallopian tube. As fetus grows, rupture may occur. Vaginal bleeding may or may not be present. Many women with ectopic pregnancy do not know they are pregnant. Usually occurs within 5 to 10 weeks of implantation. Maintain high index of suspicion with women of childbearing age experiencing abdominal pain.
- **Preeclampsia:**
Occurs in about 6% of pregnancies. Defined by hypertension and protein in the urine. RUQ pain, epigastric pain, N/V, visual disturbances, headache, and hyperreflexia are common symptoms.
In the setting of pregnancy, hypertension is defined as a BP greater than 140 systolic or greater than 90 diastolic, or a relative increase of 30 systolic and 20 diastolic from the patient's normal (pre-pregnancy) blood pressure.
Risk factors: < 20 years of age, first pregnancy, multigestational pregnancy, gestational diabetes, obesity, personal or family history of gestational hypertension.
- **Eclampsia:**
Seizures occurring in the context of preeclampsia. Remember, women may not have been diagnosed with preeclampsia.
- Maintain patient in a left lateral position, right side up 10 - 20° to minimize risk of supine hypotensive syndrome.
- Ask patient to quantify bleeding - number of pads used per hour.



Blast Injury / Incident

History

- Type of exposure (heat, gas, chemical)
- Inhalation injury
- Time of Injury
- Past medical history / Medications
- Other trauma
- Loss of Consciousness
- Tetanus/Immunization status

Signs and Symptoms

- Burns, pain, swelling
- Dizziness
- Loss of consciousness
- Hypotension/shock
- Airway compromise/distress could be indicated by hoarseness/ wheezing / Hypotension

Differential

- Superficial (1st Degree) red - painful (Don't include in TBSA)
- Partial Thickness (2nd Degree) blistering
- Full Thickness (3rd Degree) painless/charred or leathery skin
- Thermal injury
- Chemical – Electrical injury
- Radiation injury
- Blast injury

Nature of Device: Agent / Amount. Industrial Explosion. Terrorist Incident. Improvised Explosive Device.

Method of Delivery: Incendiary / Explosive

Nature of Environment: Open / Closed.

Distance from Device: Intervening protective barrier. Other environmental hazards,

Evaluate for: Blunt Trauma / Crush Injury / Compartment Syndrome / Traumatic Brain Injury / Concussion / Tympanic Membrane Rupture / Abdominal hemorrhage or Evisceration, Blast Lung Injury and Penetrating Trauma.

Scene Safety / Quantify and Triage Patients / Load and Go with Assessment / Treatment Enroute

Accidental / Intentional Explosions
(See Pearls)

	Triage Protocol UP 2 as indicated
	Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 as indicated
	Multiple Trauma Protocol TB 6 if indicated
A	IV / IO Procedure if indicated
P	Cardiac Monitor if indicated
	Thermal Burn Protocol TB 9 Chemical and Electrical Burn Protocol TB 2 if indicated
	Crush Injury Protocol TB 3 if indicated
	Radiation Incident Protocol TB 7 if indicated

Blast Lung Injury

YES

Age Appropriate
Airway Protocol(s) AR 4, 7
as indicated

NO

	Decontamination Procedure if indicated
	Pain Control Protocol UP 11 if indicated
Rapid Transport to appropriate destination using Trauma and Burn: EMS Triage and Destination Plan	

Notify Destination or Contact Medical Control

Trauma and Burn Protocol Section



Blast Injury / Incident

Pearls

• **Types of Blast Injury:**

- Primary Blast Injury: From pressure wave.
- Secondary Blast Injury: Impaled objects. Debris which becomes missiles / shrapnel.
- Tertiary Blast Injury: Patient falling or being thrown / pinned by debris.
- Most Common Cause of Death: Secondary Blast Injuries.

• **Triage of Blast Injury patients:**

- Blast Injury Patients with Burn Injuries Must be Triageed using the Thermal / Chemical / Electrical Burn Destination Guidelines for Critical / Serious / Minor Trauma and Burns
- Patients may be hard of hearing due to tympanic membrane rupture.

• **Care of Blast Injury Patients:**

- Patients may suffer multi-system injuries including blunt and penetrating trauma, shrapnel, barotrauma, burns, and toxic chemical exposure.
- Consider airway burns which should prompt early and aggressive airway management.
- Cover open chest wounds with semi-occlusive dressing.
- Use Lactated Ringers (if available) for all Critical or Serious Burns.
- Minimize IV fluids resuscitation in patients with no sign of shock or poor perfusion.

• **Blast Lung Injury:**

- Blast Lung Injury is characterized by respiratory difficulty and hypoxia. Can occur (rarely) in patients without external thoracic trauma. More likely in enclosed space or in close proximity to explosion.
- Symptoms: Dyspnea, hemoptysis, cough, chest pain, wheezing and hemodynamic instability.
- Signs: Apnea, tachypnea, hypopnea, hypoxia, cyanosis and diminished breath sounds.
- Air embolism should be considered and patient transported prone and in slight left-lateral decubitus position.
- Blast Lung Injury patients may require early intubation but positive pressure ventilation may exacerbate the injury, avoid hyperventilation.
- Air transport may worsen lung injury as well and close observation is mandated. Tension pneumothorax may occur requiring chest decompression. Be judicious with fluids as volume overload may worsen lung injury.

• **Accidental Explosions or Intentional Explosions:**

- All explosions or blasts should be considered intentional until determined otherwise.**
- Attempt to determine source of the blast to include any potential threat for aerosolization of hazardous materials.
- Evaluate scene safety to include the source of the blast that may continue to spill explosive liquids or gases.
- Consider structural collapse / Environmental hazards / Fire.
- Conditions that led to the initial explosion may be returning and lead to a second explosion.
- Greatest concern is potential threat for a secondary device.
- Patients who can, typically will attempt to move as far away from the explosive source as they safely can.
- Evaluate surroundings for suspicious items; unattended back packs or packages, or unattended vehicles.
- If patient is unconscious or there is(are) fatality(fatalities) and you are evaluating patient(s) for signs of life: Before moving note if there are wires coming from the patient(s), or it appears the patient(s) is(are) lying on a package/pack, or bulky item, do not move the patient(s), quickly back away and immediately notify a law enforcement officer.**
- If there are no indications the patient is connected to a triggering mechanism for a secondary device, expeditiously remove the patient(s) from the scene and begin transport to the hospital.
- Protect the airway and cervical spine, however, beyond the primary survey, care and a more detailed assessment should be deferred until the patient is in the ambulance.
- If there are signs the patient was carrying the source of the blast, notify law enforcement immediately and most likely, a law enforcement officer will accompany your patient to the hospital.



Chemical and Electrical Burn

History

- Type of exposure (heat, gas, chemical)
- Inhalation injury
- Time of Injury
- Past medical history / Medications
- Other trauma
- Loss of Consciousness
- Tetanus/Immunization status

Signs and Symptoms


- Burns, pain, swelling
- Dizziness
- Loss of consciousness
- Hypotension/shock
- Airway compromise/distress could be indicated by hoarseness/ wheezing / Hypotension

Differential

- Superficial (1st Degree) red - painful (Don't include in TBSA)
- Partial Thickness (2nd Degree) blistering
- Full Thickness (3rd Degree) painless/charred or leathery skin
- Thermal injury
- Chemical – Electrical injury
- Radiation injury
- Blast injury

Assure Chemical Source is NOT Hazardous to Responders.
Assure Electrical Source is NO longer in contact with patient before touching patient.

Assess Burn / Concomitant Injury Severity



	Age Appropriate Airway Protocol(s) AR 1, 2, 3, 4, 5, 6, 7 <i>if indicated</i>
	Identify Contact Points <u>Eye Involvement</u> Irrigate Involved Eye(s) with Normal Saline for 15 – 30 minutes May repeat as needed
	<u>Chemical Exposure / Burn</u> Flush Contact Area with Normal Saline for 15 minutes
	Decontamination Procedure <i>if indicated</i>
	Age Appropriate Cardiac Protocol(s) <i>if indicated</i>
	Thermal Burn Protocol TB 9

Rapid Transport to appropriate destination using
Trauma and Burn:
EMS Triage and Destination Plan

Notify Destination or Contact Medical Control

Trauma and Burn Protocol Section

Chemical and Electrical Burn

Pearls

- **Recommended Exam: Mental Status, HEENT, Neck, Heart, Lungs, Abdomen, Extremities, Back, and Neuro**
- **Green, Yellow and Red In burn severity do not apply to Triage systems.**
- **Refer to Rule of Nines: Remember the extent of the obvious external burn from an electrical source does not always reflect more extensive internal damage not seen.**
- **Chemical Burns:**
 - Refer to Decontamination Procedure.
 - Normal Saline or Sterile Water is preferred, however if not available, do not delay irrigation and use tap water. Other water sources may be used based on availability.
 - Flush the area as soon as possible with the cleanest readily available water or saline solution using copious amounts of fluids.
- **Electrical Burns:**
 - DO NOT contact patient until you are certain the source of the electrical shock is disconnected.
 - Attempt to locate contact points (generally there will be two or more.) A point where the patient contacted the source and a point(s) where the patient is grounded.
 - Sites will generally be full thickness.
 - Do not refer to as entry and exit sites or wounds.**
 - Cardiac Monitor: Anticipate ventricular or atrial irregularity including VT, VF, atrial fibrillation and / or heart blocks.
 - Attempt to identify the nature of the electrical source (AC / DC), the amount of voltage and the amperage the patient may have been exposed to during the electrical shock.



Crush Syndrome Trauma

History


- Entrapped and crushed under heavy load > 30 minutes
- Extremity / body crushed
- Building collapse, trench collapse, industrial accident, pinned under heavy equipment

Signs and Symptoms

- Hypotension
- Hypothermia
- Abnormal ECG findings
- Pain
- Anxiety

Differential

- Entrapment without crush syndrome
- Vascular injury with perfusion deficit
- Compartment syndrome
- Altered mental status



Trauma and Burn Protocol Section



Crush Syndrome Trauma

Pearls

- **Recommended exam: Mental Status, Musculoskeletal, Neuro**
- **Scene safety is of paramount importance as typical scenes pose hazards to rescuers. Call for appropriate resources.**
- **Lowest blood pressure by age: < 31 days: > 60 mmHg. 31 days to 1 year: > 70 mmHg. Greater than 1 year: 70 + 2 x age in years.**
- **Pediatric IV Fluid maintenance rate: 4 mL per first 10 kg of weight + 2 mL per second 10 kg of weight + 1 mL for every additional kg in weight.**
- **Crush syndrome typically manifests after 2 – 4 hours of crush injury, but may present in < 1 hour.**
- **Fluid resuscitation:**
 - **If access to patient and initiation of IV fluids occurs after 2 hours, give 2 liters of IV fluids in adults and 20 mL/kg of IV fluids in pediatrics and then begin > 2 hour dosing regimen.**
- **Consider all possible causes of shock and treat per appropriate protocol. Majority of decompensation in pediatrics is airway related.**
- **Decreasing heart rate and hypotension occur late in children and are signs of imminent cardiac arrest.**
- **Shock may be present with a normal blood pressure initially.**
- **Shock often is present with normal vital signs and may develop insidiously. Tachycardia may be the only manifestation.**
- **Consider all possible causes of shock and treat per appropriate protocol.**
- **Patients may become hypothermic even in warm environments.**
- **Hyperkalemia from crush syndrome can produce ECG changes described in protocol, but may also be a bizarre, wide complex rhythm. Wide complex rhythms should also be treated using the VF/Pulseless VT Protocol.**



Extremity Trauma

History


- Type of injury
- Mechanism: crush / penetrating / amputation
- Time of injury
- Open vs. closed wound / fracture
- Wound contamination
- Medical history
- Medications

Signs and Symptoms

- Pain, swelling
- Deformity
- Altered sensation / motor function
- Diminished pulse / capillary refill
- Decreased extremity temperature

Differential

- Abrasion
- Contusion
- Laceration
- Sprain
- Dislocation
- Fracture
- Amputation



Pearls

- **Recommended Exam: Mental Status, Extremity, Neuro**
- Peripheral neurovascular status is important
- In amputations, time is critical. Transport and notify medical control immediately, so that the appropriate destination can be determined.
- Hip dislocations and knee and elbow fracture / dislocations have a high incidence of vascular compromise.
- Urgently transport any injury with vascular compromise.
- Blood loss may be concealed or not apparent with extremity injuries.
- Lacerations must be evaluated for repair within 6 hours from the time of injury.
- Multiple casualty incident: Tourniquet Procedure may be considered first instead of direct pressure.



Head Trauma

History

- Time of injury
- Mechanism (blunt vs. penetrating)
- Loss of consciousness
- Bleeding
- Past medical history
- Medications
- Evidence for multi-trauma

Signs and Symptoms

- Pain, swelling, bleeding
- Altered mental status
- Unconscious
- Respiratory distress / failure
- Vomiting
- Major traumatic mechanism of injury
- Seizure

Differential


- Skull fracture
- Brain injury (Concussion, Contusion, Hemorrhage or Laceration)
- Epidural hematoma
- Subdural hematoma
- Subarachnoid hemorrhage
- Spinal injury
- Abuse

	Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 <i>if indicated</i>
	Obtain and Record GCS
	Supplemental oxygen Maintain SpO2 ≥ 90% Preferably ≥ 94%
	Prevent Oxygen desaturation events < 90%
	Blood Glucose Analysis Procedure
B	Maintain EtCO2 35 – 45 mmHg
A	IV / IO Procedure <i>if indicated</i>
P	Cardiac Monitor
	Altered Mental Status Protocol UP 4 <i>if indicated</i>
	Multiple Trauma Protocol TB 6 <i>if indicated</i>
	Age Appropriate Hypotension / Shock Protocol AM 5 / PM 3 <i>if indicated</i>
	Seizure Protocol UP 13 <i>if indicated</i>
	Spinal Motion Restriction Procedure / Protocol TB 8 <i>if indicated</i>
	Pain Control Protocol UP 11 <i>if indicated</i>
	Monitor and Reassess

**DO NOT ROUTINELY
HYPERVENTILATE**

**Evidence of
Brain Herniation:**
Unilateral or Bilateral Dilatation of
Pupils / Posturing

Hyperventilate to maintain
EtCO2 30 – 35 mmHg
See Pearls



Rapid Transport to appropriate destination
using
**Trauma and Burn:
EMS Triage and Destination Plan**



**Notify Destination or
Contact Medical Control**

Trauma and Burn Protocol Section



Head Trauma

Eye Opening Response	Verbal Response	Motor Response
4 = Spontaneous	5 = Oriented	6 = Obeys commands
3 = To verbal stimuli	4 = Confused	5 = Localizes pain
2 = To pain	3 = Inappropriate words	4 = Withdraws from pain
1 = None	2 = Incoherent	3 = Flexion to pain or decorticate
	1 = None	2 = Extension to pain or decerebrate
		1 = None

Pearls

- **Recommended Exam: Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Back, Neuro**
- **GCS is a key performance measure used in the EMS Acute Trauma Care Toolkit.**
- **A single episode of hypoxia and / or hypotension can significantly increase morbidity and mortality with head injury.**
- **Hyperventilation in head injury:**
 - Hyperventilation lowers CO₂ and causes vasoconstriction leading to increased intracranial pressure (ICP) and should not be done routinely.**
 - Use in patient with evidence of herniation (blown pupil, decorticate / decerebrate posturing, bradycardia, decreasing GCS).**
 - If hyperventilation is needed, ventilate at 14 – 18 / minute to maintain EtCO₂ between 30 - 35 mmHg.**
 - Short term option only used for severe head injury typically GCS ≤ 8 or unresponsive.**
- **Do not place in Trendelenburg position as this may increase ICP and worsen blood pressure.**
- **Poorly fitted cervical collars may also increase ICP when applied too tightly.**
- **In areas with short transport times, Drug Assisted Airway protocol is not recommended for patients who are spontaneously breathing and who have oxygen saturations of ≥ 90% with supplemental oxygen including BIAD / BVM.**
- **Hypotension:**
 - Limit IV fluids unless patient is hypotensive.
 - Increased intracranial pressure (ICP) may cause hypertension and bradycardia (Cushing's Response).
 - Usually indicates injury or shock unrelated to the head injury and should be aggressively treated.
 - Fluid resuscitation should be titrated to maintain at least a systolic BP of > 70 + 2 x the age in years.
 - Lowest blood pressure by age: < 31 days: > 60 mmHg. 31 days to 1 year: > 70 mmHg. Greater than 1 year: 70 + 2 x age in years.**
- **An important item to monitor and document is a change in the level of consciousness by serial examination.**
- **Consider Restraints if necessary for patient's and/or personnel's protection per the Restraint Procedure.**
- **Concussions:**
 - Traumatic brain injuries involving any of a number of symptoms including confusion, LOC, vomiting, or headache.
 - Any prolonged confusion or mental status abnormality which does not return to normal within 15 minutes or any documented loss of consciousness should be evaluated by a physician ASAP.
 - EMS Providers should not make return-to-play decisions when evaluating an athlete with suspected concussion. This is outside the scope of practice.**



Multiple Trauma

History

- Time and mechanism of injury
- Damage to structure or vehicle
- Location in structure or vehicle
- Others injured or dead
- Speed and details of MVC
- Restraints / protective equipment
- Past medical history
- Medications


Signs and Symptoms

- Pain, swelling
- Deformity, lesions, bleeding
- Altered mental status or unconscious
- Hypotension or shock
- Arrest

Differential (Life threatening)

- Chest: Tension pneumothorax
Flail chest
Pericardial tamponade
Open chest wound
Hemothorax
- Intra-abdominal bleeding
- Pelvis / Femur fracture
- Spine fracture / Cord injury
- Head injury (see Head Trauma)
- Extremity fracture / Dislocation
- HEENT (Airway obstruction)
- Hypothermia

	Age Appropriate Airway Protocol(s) AR 1, 2, 3, 5, 6 if indicated
	Control External Hemorrhage Consider Pelvic Binding Splint Suspected Fractures
P	Chest Decompression-Needle Procedure if indicated
	Obtain and Record GCS
A	IV / IO Procedure
P	Cardiac Monitor
	Head Injury Protocol TB 5 if indicated
	Altered Mental Status Protocol UP 4 if indicated
	Spinal Motion Restriction Procedure / Protocol TB 8 if indicated
	Pain Control Protocol UP 11 if indicated





Multiple Trauma

Pearls

- **Recommended Exam:** Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Extremities, Back, Neuro
- **Items in Red Text are key performance measures used in the EMS Acute Trauma Care Toolkit**
- **Transport Destination is chosen based on the EMS System Trauma Plan with EMS pre-arrival notification.**
- **Scene times should not be delayed for procedures. These should be performed en route when possible.**
Rapid transport of the unstable trauma patient to the appropriate facility is the goal.
- **Control external hemorrhage and prevent hypothermia by keeping patient warm.**
- **Consider Chest Decompression with signs of shock and injury to torso and evidence of tension pneumothorax.**
- **Trauma Triad of Death:**
Metabolic acidosis / Coagulopathy / Hypothermia
Appropriate resuscitation measures and keeping patient warm regardless of ambient temperature helps to mitigate metabolic acidosis, coagulopathy, and hypothermia.
- **Bag valve mask is an acceptable method of managing the airway if pulse oximetry can be maintained $\geq 90\%$**
- **Tranexamic Acid (TXA):**
Agencies utilizing TXA must have approval from your T-RAC.
- **Trauma in Pregnancy:**
Providing optimal care for the mother = optimal care for the fetus. After 20 weeks gestation (fundus at or above umbilicus) transport patient on left side with 10 – 20° of elevation.
- **Pediatric Trauma:**
Age specific blood pressure 0 – 28 days > 60 mmHg, 1 month - 1 year > 70 mmHg, 1 - 10 years > 70 + (2 x age)mmHg and 11 years and older > 90 mmHg.
- **Geriatric Trauma:**
Evaluate with a high index of suspicion.
Often occult injuries are more difficult to recognize and patients can decompensate unexpectedly with little warning.
Risk of death with trauma increases after age 55.
SBP < 110 may represent shock / poor perfusion in patients over age 65.
Low impact mechanisms, such as ground level falls might result in severe injury especially in age over 65.
- See Regional Trauma Guidelines when declaring Trauma Activation.
- Severe bleeding from an extremity not rapidly controlled with direct pressure may necessitate the application of a tourniquet.
- Maintain high-index of suspicion for domestic violence or abuse, pediatric non-accidental trauma, or geriatric abuse.



Radiation Incident

History

- Type of exposure (heat, gas, chemical)
- Inhalation injury
- Time of Injury
- Past medical history / Medications
- Other trauma
- Loss of Consciousness
- Tetanus/Immunization status


Signs and Symptoms

- Burns, pain, swelling
- Dizziness
- Loss of consciousness
- Hypotension/shock
- Airway compromise/distress could be indicated by hoarseness/ wheezing / Hypotension

Differential

- Superficial (1st Degree) red - painful (Don't include in TBSA)
- Partial Thickness (2nd Degree) blistering
- Full Thickness (3rd Degree) painless/charred or leathery skin
- Thermal injury
- Chemical – Electrical injury
- Radiation injury
- Blast injury

Scene Safety / Quantify and Triage Patients / Load and Go with Assessment / Treatment Enroute




Trauma and Burn Protocol Section

Collateral Injury: Most all injuries immediately seen will be a result of collateral injury, such as heat from the blast, trauma from concussion, treat collateral injury based on typical care for the type of injury displayed.

Quality: Determine exposure type; external irradiation, external contamination with radioactive material, internal contamination with radioactive material.

Quantify: Determine exposure (generally measured in Grays/Gy). Information may be available from those on site who have monitoring equipment, do not delay transport to acquire this information.



Time Phases of Radiation Injury
(Exposure Dose vs Clinical Outcome)

Exposure Dose (Gy)	Prodrome Severity	Manifest Illness - Symptom Severity			Prognosis
		Hematologic	Gastrointestinal	Neurologic	
0.5 to 1.0	+	+	0	0	Survival almost certain
1.0 to 2.0	+ / ++	+	0	0	Survival >90 percent
2.0 to 3.5	++	++	0	0	Probable survival
3.5 to 5.5	+++	+++	+	0	Death in 50% at 3.5 to 6 wks
5.5 to 7.5	+++	+++	++	0	Death probable in 2-3 wks
7.5 to 10	+++	+++	+++	0*	Death probable in 1-2.5 wks
10 to 20	+++	+++	+++	+++	Death certain in 5-12 days
> 20	+++	+++	+++	+++**	Death certain in 2-5 days

Abbreviations: Gy: dose in Grey;
0: no effects; +: mild; ++: moderate; +++: severe or marked

* Hypotension

** Also cardiovascular collapse, fever, shock


Modified from: Waselenko, JK, MacVittie, TJ, Blakely, WF, et al. Medical management of the acute radiation syndrome: Recommendations of the strategic national stockpile radiation working group. Ann Int Med 2004; 140:1039.

Pearls

- Dealing with a patient with a radiation exposure can be a frightening experience. Do not ignore the ABC's, a dead but decontaminated patient is not a good outcome. Refer to the Decontamination Procedure for more information.
- Normal Saline or Sterile Water is preferred, however if not available, do not delay irrigation using tap water. Other water sources may be used based on availability. Flush the area as soon as possible with the cleanest readily available water or saline solution using copious amounts of fluids.
- Three methods of exposure:**
 - External irradiation
 - External contamination
 - Internal contamination
- Two classes of radiation:**
 - Ionizing radiation (greater energy) is the most dangerous and is generally in one of three states: Alpha Particles, Beta Particles and Gamma Rays.
 - Non-ionizing (lower energy) examples include microwaves, radios, lasers and visible light.
- Radiation burns with early presentation are unlikely, it is more likely this is a combination event with either thermal or chemical burn being presented as well as a radiation exposure. Where the burn is from a radiation source, it indicates the patient has been exposed to a significant source, (> 250 rem).
- Patients experiencing radiation poisoning are not contagious. Cross contamination is only a threat with external and internal contamination.
- Typical ionizing radiation sources in the civilian setting include soil density probes used with roadway builders and medical uses such as x-ray sources as well as radiation therapy. Sources used in the production of nuclear energy and spent fuel are rarely exposure threats as is military sources used in weaponry. Nevertheless, these sources are generally highly radioactive and in the unlikely event they are the source, consequences could be significant and the patient's outcome could be grave.
- The three primary methods of protection from radiation sources:**
 - Limiting time of exposure
 - Distance from
 - Shielding from the source
- Dirty bombs ingredients generally include previously used radioactive material and combined with a conventional explosive device to spread and distribute the contaminated material.
- Refer to Decontamination Procedure / WMD / Nerve Agent Protocol for dirty contamination events.
- If there is a time lag between the time of exposure and the encounter with EMS, key clinical symptom evaluation includes: Nausea/ Vomiting, hypothermia/hyperthermia, diarrhea, neurological/cognitive deficits, headache and hypotension.
- This event may require an activation of the National Radiation Injury Treatment Network, RITN. UNC Hospitals, Wake Forest-Baptist and Duke are the NC hospitals, with burns managed at UNC and Wake Forest.



Selective Spinal Motion Restriction



Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Patients meeting all the above criteria do not require spinal motion restriction. However, patients who fail one or more criteria above require spinal motion restriction, but does NOT require use of the long spine board for immobilization.**
- **Long spine boards are NOT considered standard of care in most cases of potential spinal injury. Spinal motion restriction with cervical collar and securing patient to cot, while padding all void areas is appropriate.**
- **True spinal immobilization is not possible. Spine protection and spinal motion restriction do not equal long spine board.**
- **Spinal motion restriction is always utilized in at-risk patients. These include cervical collar, securing to stretcher, minimizing movement / transfers and maintenance of in-line spine stabilization during any necessary movement / transfers. This includes the elderly or others with body or spine habitus preventing them from lying flat.**
- **Consider spinal motion restriction in patients with arthritis, cancer, dialysis, underlying spine or bone disease.**
- Range of motion (ROM) is tested by touching chin to chest (look down), extending neck (look up), and turning head from side to side (shoulder to shoulder) without posterior cervical mid-line pain. ROM should NOT be assessed if patient has midline spinal tenderness. Patient's range of motion should not be assisted.
- **EMR may participate in spinal motion restriction per Agency Medical Director**
- **Immobilization on a long spine board is not necessary where:**
 - Penetrating trauma to the head, neck or torso with no signs / symptoms of spinal injury.
- **Concerning mechanisms that may result in spinal column injury:**
 - Fall from ≥ 3 feet and/or ≥ 5 stairs or steps
 - MVC ≥ 30 mph, rollover, and/or ejection
 - Motorcycle, bicycle, other mobile device, or pedestrian-vehicle crash
 - Diving or axial load to spine
 - Electric shock



Thermal Burn

History

- Type of exposure (heat, gas, chemical)
- Inhalation injury
- Time of Injury
- Past medical history and Medications
- Other trauma
- Loss of Consciousness
- Tetanus/Immunization status

Signs and Symptoms

- Burns, pain, swelling
- Dizziness
- Loss of consciousness
- Hypotension/shock
- Airway compromise/distress could be indicated by hoarseness/wheezing

Differential

- Superficial (1st Degree) red - painful (Don't include in TBSA)
- Partial Thickness (2nd Degree) blistering
- Full Thickness (3rd Degree) painless/charred or leathery skin
- Thermal injury
- Chemical – Electrical injury
- Radiation injury
- Blast injury

Assess Burn / Concomitant Injury Severity

< 5% TBSA 2nd/3rd Degree Burn
 No inhalation injury, Not Intubated,
 Normotensive
 GCS 14 or Greater
 Minor Burn


5-15% TBSA 2nd/3rd Degree Burn
 Suspected inhalation injury or requiring
 intubation for airway stabilization
 Hypotension or GCS 13 or Less
*(When reasonably accessible,
 transport to a Burn Center)*
 Serious Burn

>15% TBSA 2nd/3rd Degree Burn
 Burns with Multiple Trauma
 Burns with definitive airway
 compromise
*(When reasonably accessible,
 transport to a Burn Center)*
 Critical Burn

	Airway Protocol(s) AR 1, 2, 3, 5, 6 as indicated	
	Multiple Trauma Protocol TB 6 if indicated	
	Remove Rings, Bracelets / Constricting Items	
	Dry Clean Sheet or Dressings	
A	IV / IO Procedure Consider 2 IV sites if greater than 15 % TBSA	
	Normal Saline 0.25 mL / kg (x % TBSA) / hr for up to the first 8 hours. <i>(More info below)</i> Lactated Ringers if available	
	Pain Control Protocol UP 11 if indicated	
	Carbon Monoxide / Cyanide Protocol TE 2 if indicated	
	Monitor and Reassess	
	Rapid Transport to appropriate destination using Trauma and Burn: EMS Triage and Destination Plan	
	Notify Destination or Contact Medical Control	

1. Lactated Ringers preferred over Normal Saline. Use if available, if not change over once available.
2. Formula example; an 80 kg (176 lbs.) patient with 50% TBSA will need 1000 cc of fluid per hour.

Thermal Burn



Body Part	Age				
	0 yr	1 yr	5 yr	10 yr	15 yr
a = 1/2 of head	9 1/2	8 1/2	6 1/2	5 1/2	4 1/2
b = 1/2 of 1 thigh	2 3/4	3 1/4	4	4 1/4	4 1/2
c = 1/2 of 1 lower leg	2 1/2	2 1/2	2 3/4	3	3 1/4

Rule of Nines

- Seldom do you find a complete isolated body part that is injured as described in the Rule of Nines.
- More likely, it will be portions of one area, portions of another, and an approximation will be needed.
- For the purpose of determining the extent of serious injury, differentiate the area with minimal or 1st degree burn from those of partial (2nd) or full (3rd) thickness burns.
- For the purpose of determining Total Body Surface Area (TBSA) of burn, include only Partial and Full Thickness burns. Report the observation of other superficial (1st degree) burns but do not include those burns in your TBSA estimate.
- Some texts will refer to 4th 5th and 6th degree burns. There is significant debate regarding the actual value of identifying a burn injury beyond that of the superficial, partial and full thickness burn at least at the level of emergent and primary care. For our work, all are included in Full Thickness burns.
- Other burn classifications in general include:
 - 4th referring to a burn that destroys the dermis and involves muscle tissue.
 - 5th referring to a burn that destroys dermis, penetrates muscle tissue, and involves tissue around the bone.
 - 6th referring to a burn that destroys dermis, destroys muscle tissue, and penetrates or destroys bone tissue.


Estimate spotty areas of burn by using the size of the patient's palm as 1 %

Pearls

- **Recommended Exam: Mental Status, HEENT, Neck, Heart, Lungs, Abdomen, Extremities, Back, and Neuro**
- **Green, Yellow and Red In burn severity do not apply to the Start / JumpStart Triage System.**
- **Critical or Serious Burns:**
 - > 5-15% total body surface area (TBSA) 2nd or 3rd degree burns, or 3rd degree burns > 5% TBSA for any age group, or circumferential burns of extremities, or electrical or lightning injuries, or suspicion of abuse or neglect, or inhalation injury, or chemical burns, or burns of face, hands, perineum, or feet*
- Require direct transport to a Burn Center. Local facility should be utilized only if distance to Burn Center is excessive or critical interventions such as airway management are not available in the field.
- Burn patients are trauma patients, evaluate for multisystem trauma.
- Assure whatever has caused the burn is no longer contacting the injury. (Stop the burning process!)
- Early intubation is required when the patient experiences significant inhalation injuries.
- Circumferential burns to extremities are dangerous due to potential vascular compromise secondary to soft tissue swelling.
- Burn patients are prone to hypothermia - never apply ice or cool the burn, must maintain normal body temperature.
- Evaluate the possibility of geriatric abuse with burn injuries in the elderly.
- Never administer IM pain injections to a burn patient.



Traumatic Arrest (Optional)



Trauma and Burn Protocol Section



Traumatic Arrest (Optional)

Pearls

- **Recommended Exam:** Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Extremities, Back, Neuro
- **Withholding resuscitative efforts with blunt and penetrating trauma victims who meet criteria is appropriate.**
- **If transport time to Trauma Center is < 15 minutes use of ECG monitor may delay resuscitation.**
- **Rhythm determination is more helpful in rural settings or where transport to nearest facility is > 15 minutes. Omit from algorithm where appropriate.**
- **Organized rhythms for the purposes of this protocol include Ventricular Tachycardia, Ventricular Fibrillation and PEA.**
- **Wide, bizarre rhythms such as Idioventricular and severely bradycardic rhythms < 40 BPM are not organized rhythms.**
- **First arriving EMS personnel should make the assessment concerning agonal respirations, pulselessness, asystole or PEA < 40, pupillary reflexes and spontaneous body movements.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated. Consider early IO placement if available and difficult IV anticipated.**
- **DO NOT HYPERVENTILATE:** If no advanced airway (BIAD, ETT) compressions to ventilations are 30:2. If advanced airway in place ventilate 8 – 10 breaths per minute.
- **ALS procedures should optimally be performed during rapid transport.**
- **Time considerations:**
 - **From the time cardiac arrest is identified, if CPR is performed \geq 15 minutes with no ROSC consider termination of resuscitation.**
 - **From the time cardiac arrest is identified, if transport time to closest Trauma Center is > 15 minutes consider termination of resuscitation.**
- **Lightning strike, drowning or in situations causing hypothermia resuscitation should be initiated.**
- **Where multiple lightning strike victims are found used Reverse Triage: Begin CPR where apneic / pulseless**
- **Agencies utilizing Targeted Temperature Management Protocol should not cool the trauma patient, but rather make every effort to maintain warmth.**



Pediatric Asystole / PEA

History

- Events leading to arrest
- Estimated downtime
- SAMPLE
- Existence of terminal illness
- Airway obstruction
- Hypothermia
- Suspected abuse

Signs and Symptoms

- Pulseless
- Apneic
- No electrical activity on ECG
- No heart tones on auscultation

Differential

- Respiratory failure
- Foreign body
- Infection (croup, epiglottitis)
- Congenital heart disease
- See Reversible Causes below

Pediatric Pulseless Arrest Protocol

Criteria for Death / No Resuscitation Review DNR / MOST Form

YES

NO

Decomposition
Rigor mortis
Dependent lividity
Blunt force trauma
Injury incompatible with life
Extended downtime with asystole

Do not begin resuscitation

Follow Deceased Subjects Policy

AT ANY TIME

Return of Spontaneous Circulation

Go to Post Resuscitation Protocol

	<p>Begin Continuous CPR Compressions Push Hard ($\geq 1/3$ AP Diameter of Chest) (1.5 inches Infant / 2 inches in Children) Push Fast (100 - 120 / min) Change Compressors every 2 minutes (sooner if fatigued) <i>(Limit changes / pulse checks ≤ 10 seconds)</i></p> <p>Ventilation rate: 1 breath every 2 seconds when age < 1 1 breathe every 3 seconds when age ≥ 1 15:2 Compression:Ventilation if no Advanced Airway</p>
	AED Procedure <i>if available</i>
P	Cardiac Monitor
	IV or IO Access Protocol UP 6
A	<p>Epinephrine 1:10,000 0.01 mg/kg IV / IO Maximum Single Dose 1mg Or Epinephrine 1:1000 0.1 mg / kg ETT Maximum 2.5 mg Repeat every 5 minutes</p>
	Search for Reversible Causes
	Blood Glucose Analysis Procedure <i>if applicable</i>

Reversible Causes

Hypovolemia
Hypoxia
Hydrogen ion (acidosis)
Hypothermia
Hypo / Hyperkalemia

Tension pneumothorax
Tamponade; cardiac
Toxins
Thrombosis; pulmonary (PE)
Thrombosis; coronary (MI)

Notify Destination or Contact Medical Control

Pediatric Cardiac Protocol Section



Pediatric Asystole / PEA

Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks.**
- **Refer to optional protocol AC 11 or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated. Compress $\geq 1/3$ anterior-posterior diameter of chest, in infants 1.5 inches and in children 2 inches.**
- **Majority of pediatric arrests stem from a respiratory insult or hypoxic event. Compressions should be coupled with ventilations.**
- **When advanced airway not in place perform 15 compressions with 2 ventilations.**
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.**
- **DO NOT HYPERVENTILATE:**
If advanced airway in place ventilate:
 - **Age < 1 year: 1 breath every 2 seconds with continuous, uninterrupted compressions.**
 - **Age ≥ 1 year: 1 breath every 3 seconds with continuous, uninterrupted compressions.**
- **Airway is a more important intervention in pediatric arrests. This should be accomplished quickly with BVM or BIAD.**
- **Patient survival is often dependent on proper ventilation and oxygenation / airway Interventions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **High-Quality CPR:**
 - Make sure chest compressions are being delivered at 100 – 120 / min.
 - Make sure chest compressions are adequate depth for age and body habitus.
 - Make sure you allow full chest recoil with each compression to provide maximum perfusion.
 - Minimize all interruptions in chest compressions to < 10 seconds.
 - Use AED or apply ECG monitor / defibrillator as soon as available.
- **Defibrillation:** Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
- **End Tidal CO₂ (EtCO₂)**
 - If EtCO₂ is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
 - If EtCO₂ spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Special Considerations**
 - **Maternal Arrest** - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm. Defibrillation is safe at all energy levels.
 - **Renal Dialysis / Renal Failure** - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
 - **Opioid Overdose** - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol UP 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
 - **Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike** – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.**



Pediatric Bradycardia With Poor Perfusion

History


- Past medical history
- Foreign body exposure
- Respiratory distress or arrest
- Apnea
- Possible toxic or poison exposure
- Congenital disease
- Medication (maternal or infant)

Signs and Symptoms

- Decreased heart rate
- Delayed capillary refill or cyanosis
- Mottled, cool skin
- Hypotension or arrest
- Altered level of consciousness

Differential

- Respiratory failure, Foreign body, Secretions, Infection (croup, epiglottitis)
- Hypovolemia (dehydration)
- Congenital heart disease
- Trauma
- Tension pneumothorax
- Hypothermia
- Toxin or medication
- Hypoglycemia
- Acidosis



Reversible Causes

Hypovolemia
 Hypoxia
 Hydrogen ion (acidosis)
 Hypothermia
 Hypo / Hyperkalemia
 Hypoglycemia

Tension pneumothorax
 Tamponade; cardiac
 Toxins
 Thrombosis; pulmonary (PE)
 Thrombosis; coronary (MI)

Suspected Beta-Blocker or Calcium Channel Blocker

Follow Pediatric Toxicology Protocol



Pediatric Bradycardia With Poor Perfusion

Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Bradycardia is often associated with hypoxia so insure patent airway, breathing, and circulation as needed.**
- **Begin CPR immediately with persistent bradycardia and poor perfusion despite adequate oxygenation and ventilation.**
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.**
- **Rhythm should be interpreted in the context of symptoms and pharmacological treatment given only when symptomatic, otherwise monitor and reassess.**
- **Consider hyperkalemia with wide complex, bizarre appearance of QRS complex, and bradycardia.**
- **12-Lead ECG:**
 - **12 Lead ECG not necessary to diagnose and treat**
 - **Obtain when patient is stable and/or following rhythm conversion.**
- **Unstable condition**
 - **Condition which acutely impairs vital organ function and cardiac arrest may be imminent.**
 - **If at any point patient becomes unstable move to unstable arm in algorithm**
- **Epinephrine is first drug choice for persistent, symptomatic bradycardia.**
- **Atropine:**
 - **Second choice, unless there is evidence of increased vagal tone or a primary AV conduction block, then give atropine first.**
 - **Ineffective and potentially harmful in cardiac transplantation. May cause paradoxical bradycardia.**
- **Symptomatic bradycardia causing shock or peri-arrest condition:**
 - **If no IV or IO access immediately available, start Transcutaneous Pacing, establish IV / IO access, and then administer epinephrine.**
 - **Epinephrine should be administered followed Atropine if no response.**
- **Symptomatic condition**
 - **Arrhythmia is causing symptoms such as palpitations, lightheadedness, or dyspnea, but cardiac arrest is not imminent.**
 - **Symptomatic bradycardia usually occurs at rates < 50 beats per minute.**
 - **Search for underlying causes such as hypoxia or impending respiratory failure.**
- **Serious Signs / Symptoms:**
 - **Hypotension. Acutely altered mental status. Signs of shock / poor perfusion. Chest pain with evidence of ischemia (STEMI, T wave inversions or depressions.) Acute CHF.**
- **Transcutaneous Pacing Procedure (TCP)**
 - **Indicated with unstable bradycardia unresponsive to medical therapy.**
 - **If time allows transport to specialty center because transcutaneous pacing is a temporizing measure.**
 - **Transvenous / permanent pacemaker will probably be needed.**
 - **Immediate TCP with high-degree AV block (2d or 3d degree) with no IV / IO access.**
- **Most maternal medications pass through breast milk to the infant so maintain high-index of suspicion for OD-toxins.**
- **Hypoglycemia, severe dehydration and narcotic effects may produce bradycardia. Many other agents a child ingests can cause bradycardia, often is a single dose.**



Pediatric Pulmonary Edema / CHF

History


- Congenital Heart Disease
- Chronic Lung Disease
- Congestive heart failure
- Past medical history

Signs/Symptoms

- Infant: Respiratory distress, poor feeding, lethargy, weight gain, +/- cyanosis
- Child/Adolescent: Respiratory distress, bilateral rales, apprehension, orthopnea, jugular vein distention (rare), pink, frothy sputum, peripheral edema, diaphoresis, chest pain
- Hypotension, shock

Differential

- Congestive heart failure
- Asthma
- Anaphylaxis
- Aspiration
- Pleural effusion
- Pneumonia
- Pulmonary embolus
- Pericardial tamponade
- Toxic Exposure



Pearls

- **Recommended exam: Mental status, Respiratory, Cardiac, Skin, Neuro**
- **Contact Medical Control early in the care of the pediatric cardiac patient.**
- **Most children with CHF have a congenital heart defect, obtain a precise past medical history.**
- **Congenital heart disease varies by age:**
 - < 1 month: Tetralogy of Fallot, Transposition of the great arteries, Coarctation of the aorta.
 - 2 – 6 months: Ventricular septal defects (VSD), Atrioseptal defects (ASD).
 - Any age: Myocarditis, Pericarditis, SVT, heart blocks.
- **Treatment of Congestive Heart Failure / Pulmonary edema may vary depending on the underlying cause and may include the following with consultation by Medical Control:**
 - Morphine Sulfate: 0.1 mg/kg IV / IO. Max single dose 5mg/dose
 - Fentanyl: 1 mcg/kg IV / IO. Max single dose 50 mcg.
 - Nitroglycerin: Dose determined after consultation of Medical Control.
 - Lasix 1 mg/kg IV / IO.
 - Agency specific vasopressor.
- Do not assume all wheezing is pulmonary, especially in a cardiac child: avoid albuterol unless strong history of recurrent wheezing secondary to pulmonary etiology (discuss with Medical Control)



Pediatric Cardiac Arrest

(Ages 31 days to 15 years)

History


- Time of arrest
- Medical history
- Medications
- Possibility of foreign body
- Hypothermia

Signs and Symptoms

- Unresponsive
- Cardiac arrest

Differential

- Respiratory failure: Foreign body, Secretions, Infection (croup, epiglottitis)
- Hypovolemia (dehydration)
- Congenital heart disease
- Trauma
- Tension pneumothorax, cardiac tamponade, pulmonary embolism
- Hypothermia
- Toxin or medication
- Electrolyte abnormalities (Glucose, K)
- Acidosis





Pediatric Cardiac Arrest

(Ages 31 days to 15 years)

Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional protocol or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated. Compress \geq 1/3 anterior-posterior diameter of chest, in infants 1.5 inches and in children 2 inches.**
- **Majority of pediatric arrests stem from a respiratory insult or hypoxic event. Compressions should be coupled with ventilations.**
- **When advanced airway not in place perform 15 compressions with 2 ventilations.**
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.**
- **DO NOT HYPERVENTILATE:**
If advanced airway in place ventilate:
Age < 1 year: 1 breath every 2 seconds with continuous, uninterrupted compressions.
Age \geq 1 year: 1 breath every 3 seconds with continuous, uninterrupted compressions.
- **Patient survival is often dependent on proper ventilation and oxygenation / airway Interventions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **High-Quality CPR:**
Make sure chest compressions are being delivered at 100 – 120 / min.
Make sure chest compressions are adequate depth for age and body habitus.
Make sure you allow full chest recoil with each compression to provide maximum perfusion.
Minimize all interruptions in chest compressions to < 10 seconds.
Use AED or apply ECG monitor / defibrillator as soon as available.
- **Defibrillation:**
Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.
Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- **End Tidal CO₂ (EtCO₂)**
If EtCO₂ is < 10 mmHg, improve chest compressions. Goal is \geq 20 mmHg.
If EtCO₂ spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Special Considerations**
Maternal Arrest - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm.
Defibrillation is safe at all energy levels.
Renal Dialysis / Renal Failure - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
Opioid Overdose - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol UP 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.**



Pediatric Tachycardia

Narrow Complex (≤ 0.09 sec)

History


- Past medical history
- Medications or Toxic Ingestion (Aminophylline, Diet pills, Thyroid supplements, Decongestants, Digoxin)
- Drugs (nicotine, cocaine)
- Congenital Heart Disease
- Respiratory Distress
- Syncope or Near Syncope

Signs and Symptoms

- Heart Rate: Child > 180 /bpm
Infant > 220 /bpm
- Pale or Cyanosis
- Diaphoresis
- Tachypnea
- Vomiting
- Hypotension
- Altered Level of Consciousness
- Pulmonary Congestion
- Syncope

Differential

- Heart disease (Congenital)
- Hypo / Hyperthermia
- Hypovolemia or Anemia
- Electrolyte imbalance
- Anxiety / Pain / Emotional stress
- Fever / Infection / Sepsis
- Hypoxia, Hypoglycemia
- Medication / Toxin / Drugs (see HX)
- Pulmonary embolus
- Trauma, Tension Pneumothorax





Pediatric Tachycardia

Narrow Complex (≤ 0.09 sec)

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Monomorphic QRS:**
All QRS complexes in a single lead are similar in shape.
- **Polymorphic QRS:**
QRS complexes in a single lead will change from complex to complex.
- Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.
- Rhythm should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.
- **12-Lead ECG:**
12-Lead ECG not necessary to diagnose and treat.
Obtain when patient is stable and/or following rhythm conversion.
When administering adenosine, obtaining a continuous 12-Lead can be helpful to physicians.
- **Unstable condition:**
Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
If at any point patient becomes unstable move to unstable arm in algorithm
If IV or IO access is in place, may administer adenosine and repeat, prior to synchronized cardioversion.
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.
- **Serious Signs and Symptoms:**
Respiratory distress / failure.
Signs of shock / poor perfusion with or without hypotension.
AMS
Sudden collapse with rapid, weak pulse
- **Narrow Complex Tachycardia (≤ 0.09 seconds):**
Sinus tachycardia: P waves present. Variable R-R waves. Infants usually < 220 beats / minute. Children usually < 180 beats / minute.
SVT: > 90 % of children with SVT will have a narrow QRS (≤ 0.09 seconds.) P waves absent or abnormal. R-R waves not variable. Usually abrupt onset. Infants usually > 220 beats / minute. Children usually > 180 beats / minute.
Atrial Flutter / Fibrillation
- **Vagal Maneuvers:**
Breath holding. Blowing a glove into a balloon. Have child blow out "birthday candles" or through an obstructed straw. Infants: May put a bag of ice water over the upper half of the face careful not to occlude the airway.
- Separating the child from the caregiver may worsen the child's clinical condition.
- Monitor for respiratory depression and hypotension associated if Diazepam, Lorazepam, or Midazolam is used.
- Continuous pulse oximetry is required for all SVT Patients if available.



Pediatric Tachycardia

Wide Complex (> 0.09 sec)

History


- Past medical history
- Medications or Toxic Ingestion (Aminophylline, Diet pills, Thyroid supplements, Decongestants, Digoxin)
- Drugs (nicotine, cocaine)
- Congenital Heart Disease
- Respiratory Distress
- Syncope or Near Syncope

Signs and Symptoms

- Heart Rate: Child > 180/bpm
Infant > 220/bpm
- Pale or Cyanosis
- Diaphoresis
- Tachypnea
- Vomiting
- Hypotension
- Altered Level of Consciousness
- Pulmonary Congestion
- Syncope

Differential

- Heart disease (Congenital)
- Hypo / Hyperthermia
- Hypovolemia or Anemia
- Electrolyte imbalance
- Anxiety / Pain / Emotional stress
- Fever / Infection / Sepsis
- Hypoxia, Hypoglycemia
- Medication / Toxin / Drugs (see HX)
- Pulmonary embolus
- Trauma, Tension Pneumothorax





Pediatric Tachycardia

Wide Complex (> 0.09 sec)

Pearls

- **Recommended Exam: Mental Status, Skin, Neck, Lung, Heart, Abdomen, Back, Extremities, Neuro**
- **Monomorphic QRS:**
All QRS complexes in a single lead are similar in shape.
- **Polymorphic QRS:**
QRS complexes in a single lead will change from complex to complex.
- Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.
- Rhythm should be interpreted in the context of symptoms and pharmacological or electrical treatment given only when symptomatic, otherwise monitor and reassess.
- **12-Lead ECG:**
12-Lead ECG not necessary to diagnose and treat.
Obtain when patient is stable and/or following rhythm conversion.
When administering adenosine, obtaining a continuous 12-Lead can be helpful to physicians.
- **Unstable condition:**
Condition which acutely impairs vital organ function and cardiac arrest may be imminent.
If at any point patient becomes unstable move to unstable arm in algorithm
- Document all rhythm changes with monitor strips and obtain monitor strips with each therapeutic intervention.
- **Serious Signs and Symptoms:**
Respiratory distress / failure.
Signs of shock / poor perfusion with or without hypotension.
AMS
Sudden collapse with rapid, weak pulse
- **Serious Signs and Symptoms:**
Respiratory distress / failure.
Signs of shock / poor perfusion with or without hypotension.
AMS
Sudden collapse with rapid, weak pulse
- **Wide Complex Tachycardia (≥ 0.09 seconds):**
SVT with aberrancy.
VT: Uncommon in children. Rates may vary from near normal to > 200 / minute. Most children with VT have underlying heart disease / cardiac surgery / long QT syndrome / cardiomyopathy.
Amiodarone 5 mg / kg over 20 – 60 minutes or Procainamide 15 mg / kg over 30 – 60 minutes IV / IO are recommended agents. They should not be administered together. Consultation with Medical Control is advised when these agents are considered.
- **Torsades de Pointes / Polymorphic (multiple shaped) Tachycardia:**
Rate is typically 150 to 250 beats / minute.
Associated with long QT syndrome, hypomagnesaemia, hypokalemia, many cardiac drugs.
May quickly deteriorate to VT.
Separating the child from the caregiver may worsen the child's clinical condition.
- Monitor for respiratory depression and hypotension associated if Diazepam, Lorazepam, or Midazolam is used.
- Continuous pulse oximetry is required for all SVT Patients if available.



Pediatric Ventricular Fibrillation Pulseless Ventricular Tachycardia

History

- Events leading to arrest
- Estimated downtime
- Past medical history
- Medications
- Existence of terminal illness
- Airway obstruction
- Hypothermia

Signs and Symptoms

- Unresponsive
- Cardiac Arrest

Differential

- Respiratory failure / Airway obstruction
- Hyper / hypokalemia, Hypovolemia
- Hypothermia, Hypoglycemia, Acidosis
- Tension pneumothorax, Tamponade
- Toxin or medication
- Thrombosis: Coronary / Pulmonary Embolism
- Congenital heart disease

Pediatric Pulseless Arrest Protocol PC 4

	<p>Begin Continuous CPR Compressions Push Hard ($\geq 1/3$ AP Diameter of Chest) (1.5 inches Infant / 2 inches in Children) Push Fast (100 - 120 / min) Change Compressors every 2 minutes (sooner if fatigued) <i>(Limit changes / pulse checks ≤ 10 seconds)</i></p> <p>Ventilation rate: 1 breath every 2 seconds when age < 1 1 breathe every 3 seconds when age ≥ 1 15:2 Compression:Ventilation if no Advanced Airway</p>
	Automated Defibrillation Procedure
P	Defibrillation Manual Procedure <ul style="list-style-type: none"> • First shock: 2 J / Kg • Second shock: 4 J / Kg • Subsequent shocks ≥ 4 J / kg Maximum 10 J / kg or adult dose
	IV / IO Protocol UP 6
A	<p>Epinephrine 1:10,000 0.01 mg/kg IV / IO Maximum 1mg Or Epinephrine 1:1000 0.1 mg / kg ETT Maximum 2.5 mg Repeat every 3 – 5 minutes</p>
	<p>If Rhythm Refractory to defibrillation</p> <ul style="list-style-type: none"> • Continue CPR and give Agency specific Anti-arrhythmic(s) in a drug-shock-drug-shock pattern. • Continue CPR up to point where you are ready to defibrillate with device charged. <p>Repeat pattern during resuscitation.</p>
P	<p>Amiodarone 5mg/kg See Pearls</p>

<p>Reversible Causes</p> <p>Hypovolemia Hypoxia Hydrogen ion (acidosis) Hypothermia Hypo / Hyperkalemia Hypoglycemia</p> <p>Tension pneumothorax Tamponade; cardiac Toxins Thrombosis; pulmonary (PE) Thrombosis; coronary (MI)</p>

AT ANY TIME

Return of Spontaneous Circulation

Go to Post Resuscitation Protocol

Notify Destination or Contact Medical Control



Pediatric Ventricular Fibrillation Pulseless Ventricular Tachycardia

Pearls

- **Team Focused Approach / Pit-Crew Approach recommended; assigning responders to predetermined tasks. Refer to optional protocol or development of local agency protocol.**
- **Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated. Compress $\geq 1/3$ anterior-posterior diameter of chest, in infants 1.5 inches and in children 2 inches.**
- **Majority of pediatric arrests stem from a respiratory insult or hypoxic event. Compressions should be coupled with ventilations.**
- **When advanced airway not in place perform 15 compressions with 2 ventilations.**
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.**
- **DO NOT HYPERVENTILATE:**
If advanced airway in place ventilate:
Age < 1 year: 1 breath every 2 seconds with continuous, uninterrupted compressions.
Age ≥ 1 year: 1 breath every 3 seconds with continuous, uninterrupted compressions.
- **Patient survival is often dependent on proper ventilation and oxygenation / airway Interventions.**
- **Do not interrupt compressions to place endotracheal tube. Consider BIAD first to limit interruptions.**
- **High-Quality CPR:**
Make sure chest compressions are being delivered at 100 – 120 / min.
Make sure chest compressions are adequate depth for age and body habitus.
Make sure you allow full chest recoil with each compression to provide maximum perfusion.
Minimize all interruptions in chest compressions to < 10 seconds.
Use AED or apply ECG monitor / defibrillator as soon as available.
- **Defibrillation:**
Follow manufacture's recommendations concerning defibrillation / cardioversion energy when specified.
Charge defibrillator during chest compressions, near the end of 2-minute cycle, to decrease peri-shock pause.
Following defibrillation, provider should immediately restart chest compressions with no pulse check until end of next cycle.
- **End Tidal CO₂ (EtCO₂)**
If EtCO₂ is < 10 mmHg, improve chest compressions. Goal is ≥ 20 mmHg.
If EtCO₂ spikes, typically > 40 mmHg, consider Return of Spontaneous Circulation (ROSC)
- **IV / IO access and drug delivery are secondary to high-quality chest compressions and early defibrillation.**
- **IV access is preferred route. Follow IV or IO Access Protocol UP 6.**
- **Special Considerations**
Maternal Arrest - Treat mother per appropriate protocol with immediate notification to Medical Control and rapid transport preferably to obstetrical center if available and proximate. Place mother supine and perform Manual Left Uterine Displacement moving uterus to the patient's left side. IV/IO access preferably above diaphragm.
Defibrillation is safe at all energy levels.
Renal Dialysis / Renal Failure - Refer to Dialysis / Renal Failure Protocol AM 3 caveats when faced with dialysis / renal failure patient experiencing cardiac arrest.
Opioid Overdose - If suspected, administer Naloxone per Overdose / Toxic Ingestion Protocol UP 7 while ensuring airway, oxygenation, ventilations, and high-quality chest compressions.
Drowning / Suffocation / Asphyxiation / Hanging / Lightning Strike – Hypoxic associated cardiac arrest and prompt attention to airway and ventilation is priority followed by high-quality and continuous chest compressions and early defibrillation.
- **Success is based on proper planning and execution. Procedures require space and patient access. Make room to work.**



Pediatric Post Resuscitation

History

- Respiratory arrest
- Cardiac arrest

Signs/Symptoms

- Return of pulse

Differential


- Continue to address specific differentials associated with the original dysrhythmia

	Pediatric Airway Protocol(s) AR 5 - 7 as needed
	Monitor Vital Signs / Reassess
	Blood Glucose Analysis Procedure
	Optimize Ventilation and Oxygenation <ul style="list-style-type: none"> • Maintain SpO2 ≥ 92 – 98% • Advanced airway if indicated • Age Appropriate Respiratory Rate • Remove Impedance Threshold Device DO NOT HYPERVENTILATE
	ETCO2 ideally 35 – 45 mm Hg
B	12 Lead ECG Procedure
	IV or IO Protocol UP 6
P	Cardiac Monitor
	Pediatric Diabetic Protocol PM 2 if indicated
	Pediatric Hypotension / Shock Protocol PM 3 if indicated
	Pediatric Bradycardia Protocol PC 2 if indicated
	Pediatric Tachycardia Protocol PC 5, 6 as indicated

Hypotension Age Based
0 – 31 Days < 60 mmHg
1 Month to 1 Year < 70 mmHg
> than 1 Year < 70 + (2 x age) mmHg

Arrhythmias are common and usually self limiting after ROSC

If Arrhythmia Persists follow Rhythm Appropriate Protocol



P	Continue Antiarrhythmic Utilized Refer to Appropriate Pediatric Arrhythmia Protocol

Post-intubation / BIAD Management Protocol AR 8

Notify Destination or Contact Medical Control



Pediatric Post Resuscitation

Pearls

- **Recommended Exam: Mental Status, Neck, Skin, Lungs, Heart, Abdomen, Extremities, Neuro**
- **Goals of care are to preserve neurologic function, prevent secondary organ damage, treat the underlying cause of illness, and optimize prehospital care. Frequent reassessment is necessary.**
- **Hyperventilation is a significant cause of hypotension and recurrence of cardiac arrest in the post resuscitation phase and must be avoided. Titrate FiO₂ to maintain SpO₂ of 92 - 98%.**
- **Use length-based or weight-based pediatric resuscitation system for medication, equipment, cardioversion, and defibrillation guidance. Pediatric paddles should be used in children < 10 kg.**
- **Pain/sedation:**
 - Patients requiring advanced airways and ventilation commonly experience pain and anxiety. Unrelieved pain can lead to increased catecholamine release, ischemia, immunosuppression, and prolonged hospitalization.
 - Ventilated patients cannot communicate pain / anxiety and providers are poor at recognizing pain / anxiety.
 - Vital signs such as tachycardia and / or hypertension can provide clues to inadequate sedation, however they both are not always reliable indicators of patient's lack of adequate sedation.
 - Pain must be addressed first, before anxiety. Opioids are typically the first line agents before benzodiazepines. Ketamine is also a reasonable first choice agent.
- **Ventilator / Ventilation strategies:**
 - Tailored to individual patient presentations. Medical Control can indicate different strategies above.
 - In general ventilation with BVM should cause chest rise. With mechanical ventilation a reasonable tidal volume should be about 6 mL/kg and peak pressures should be < 30 cmH₂O.
 - Continuous pulse oximetry and capnography should be maintained during transport for monitoring.
 - Head of bed should be maintained at least 10 – 20 degrees of elevation when possible to decrease aspiration risk.
- **EtCO₂ Monitoring:**
 - Initial End tidal CO₂ may be elevated immediately post-resuscitation, but will usually normalize.
 - Goal is 35 – 45 mmHg but DO NOT hyperventilate to achieve.
 - EtCO₂ should be continually monitored with advanced airway in place.
- **Administer resuscitation fluids and vasopressor agents to maintain SBP at targets listed on page 1. This table represents minimal SBP targets.**
- **Targeted Temperature Management is recommended in pediatrics, but prehospital use is not associated with improved outcomes. Transport to facility capable of intensive pediatric care.**
- **Consider transport to facility capable of managing the post-arrest patient including hypothermia therapy, cardiology / cardiac catheterization, intensive care service, and neurology services.**
- **The condition of post-resuscitation patients fluctuates rapidly and continuously, and they require close monitoring. Appropriate post-resuscitation management may best be planned in consultation with Medical Control.**



Pediatric Allergic Reaction

History


- Onset and location
- Insect sting or bite
- Food allergy / exposure
- Medication allergy / exposure
- New clothing, soap, detergent
- Past medical history / reactions
- Medication history

Signs and Symptoms

- Itching or hives
- Coughing / wheezing or respiratory distress
- Chest or throat constriction
- Difficulty swallowing
- Hypotension or shock
- Edema

Differential

- Urticaria (rash only)
- Anaphylaxis (systemic effect)
- Shock (vascular effect)
- Angioedema (drug induced)
- Aspiration / Airway obstruction
- Vasovagal event
- Asthma / COPD / CHF





Pediatric Allergic Reaction

Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs**
- **Anaphylaxis is an acute and potentially lethal multisystem allergic reaction.**
- **Epinephrine administration:**
 - **Drug of choice and the FIRST drug that should be administered in acute anaphylaxis (Moderate / Severe Symptoms.) IM Epinephrine should be administered in priority before or during attempts at IV or IO access.**
 - **Diphenhydramine and steroids have no proven utility in Moderate / Severe anaphylaxis and may be given only After Epinephrine. Diphenhydramine and steroids should NOT delay repeated Epinephrine administration.**
 - **In Moderate and Severe anaphylaxis Diphenhydramine may decrease mental status. Oral Diphenhydramine should NOT be given to a patient with decreased mental status and / or a hypotensive patient as this may cause nausea and / or vomiting.**
- **Anaphylaxis unresponsive to repeat doses of IM epinephrine may require IV epinephrine administration by IV push or epinephrine infusion. Contact Medical Control for appropriate dosing.**
- **Symptom Severity Classification:**
 - **Mild symptoms:**
 - **Flushing, hives, itching, erythema with normal blood pressure and perfusion.**
 - **Moderate symptoms:**
 - **Flushing, hives, itching, erythema plus respiratory (wheezing, dyspnea, hypoxia) or gastrointestinal symptoms (nausea, vomiting, abdominal pain) with normal blood pressure and perfusion.**
 - **Severe symptoms:**
 - **Flushing, hives, itching, erythema plus respiratory (wheezing, dyspnea, hypoxia) or gastrointestinal symptoms (nausea, vomiting, abdominal pain) with hypotension and poor perfusion.**
- **Allergic reactions may occur with only respiratory and gastrointestinal symptoms and have no rash / skin involvement.**
- **Angioedema is seen in moderate to severe reactions and is swelling involving the face, lips or airway structures. This can also be seen in patients taking blood pressure medications like Prinivil / Zestril (lisinopril) -typically end in -il.**
- **Fluids and Medication titrated to maintain a SBP $>70 + (\text{age in years} \times 2)$ mmHg.**
- **EMR / EMT may administer Epinephrine IM and may administer from EMS supply.** Agency Medical Director may require contact of medical control prior to EMR / EMT administering any medication.
- **EMR may administer diphenhydramine by oral route only and may administer from EMS supply.** Agency Medical Director may require contact of medical control prior to EMT / EMR administering any medication.
- **EMT may administer Albuterol if patient already prescribed and may administer from EMS supply.** Agency Medical Director may require contact of medical control prior to EMT / EMR administering any medication.
- **Patients with moderate and severe reactions should receive a 12 lead ECG and should be continually monitored, but this should NOT delay administration of epinephrine.**
- **The shorter the onset from exposure to symptoms the more severe the reaction.**



Pediatric Diabetic

History

- Past medical history
- Medications
- Recent blood glucose check
- Last meal


Signs and Symptoms

- Altered mental status
- Combative / irritable
- Diaphoresis
- Seizures
- Abdominal pain
- Nausea / vomiting
- Weakness
- Dehydration
- Deep / rapid breathing

Differential

- Alcohol / drug use
- Toxic ingestion
- Trauma; head injury
- Seizure
- CVA
- Altered baseline mental status.

	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure <i>if indicated</i>
A	IV / IO Procedure
P	Cardiac Monitor
	Altered Mental Status Protocol UP 4 <i>if indicated</i>
	Hypotension / Shock Protocol AM 5 <i>if indicated</i>
	Seizure Protocol UP 13 <i>if indicated</i>





Pediatric Diabetic

Pearls

- **Recommended Exam: Mental Status, HEENT, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Patients with prolonged hypoglycemia may not respond to glucagon.**
- **Do not administer oral glucose to patients that are not able to swallow or protect their airway.**
- **Quality control checks should be maintained per manufacturers recommendation for all glucometers.**
- **D10 / D25 Preparation:**
 - **D10: Remove 10 mL of D50 from a D50 vial. Add 40 mL of NS with the 10 mL of D50 – total volume 50 mL.**
 - **D10: Alternative, Discard 40 mL from the D50 vial and draw up 40 mL of NS – total volume 50 mL.**
 - **D25: Remove 25 mL of D50 and draw up 25 mL of NS – total volume 50 mL.**
- **Patient's refusing transport to medical facility after treatment of hypoglycemia:**

Adult caregiver must be present with pediatric patient.

Blood sugar must be ≥ 80 , patient has ability to eat and availability of food with responders on scene.

Patient must have known history of diabetes and not taking any oral diabetic agents.

Patient returns to normal mental status and has a normal neurological exam with no new neurological deficits.

Must demonstrate capacity to make informed health care decisions. See Universal Patient Care Protocol UP-1.

Otherwise contact medical control.

- **Hypoglycemia with Oral Agents:**

Patients taking oral diabetic medications should be strongly encouraged to allow transportation to a medical facility. They are at risk of recurrent hypoglycemia that can be delayed for hours and require close monitoring even after normal blood glucose is established. Not all oral agents have prolonged action so Contact Medical Control for advice. Patients who meet criteria to refuse care should be instructed to contact their physician immediately and consume a meal.
- **Hypoglycemia with Insulin Agents:**

Many forms of insulin now exist. Longer acting insulin places the patient at risk of recurrent hypoglycemia even after a normal blood glucose is established. Not all insulins have prolonged action so Contact Medical Control for advice. Patients who meet criteria to refuse care should be instructed to contact their physician immediately and consume a meal.



Pediatric Hypotension / Shock

History

- Blood loss
- Fluid loss
- Vomiting
- Diarrhea
- Fever
- Infection

Signs and Symptoms


- Restlessness, confusion, weakness
- Dizziness
- Tachycardia
- Hypotension (Late sign)
- Pale, cool, clammy skin
- Delayed capillary refill
- Dark-tarry stools

Differential

- Shock
 - Hypovolemic
 - Cardiogenic
 - Septic
 - Neurogenic
 - Anaphylactic
- Trauma
- Infection
- Dehydration
- Congenital heart disease
- Medication or Toxin

	Blood Glucose Analysis Procedure
A	IV / IO Procedure
P	Cardiac Monitor
	Pediatric Airway Protocol(s) <i>if indicated</i>
	Diabetic Protocol PM 2 <i>if indicated</i>

History and Exam Suggest Type of Shock





Hypotension / Shock

Pearls

- **Recommended Exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro**
- **Lowest blood pressure by age: < 31 days: > 60 mmHg. 31 days to 1 year: > 70 mmHg. Greater than 1 year: 70 + 2 x age in years.**
- **Consider all possible causes of shock and treat per appropriate protocol. Majority of decompensation in pediatrics is airway related.**
- **Decreasing heart rate and hypotension occur late in children and are signs of imminent cardiac arrest.**
- **Shock may be present with a normal blood pressure initially.**
- **Shock often is present with normal vital signs and may develop insidiously. Tachycardia may be the only manifestation.**
- **Consider all possible causes of shock and treat per appropriate protocol.**
- **Hypovolemic Shock:**
Hemorrhage, trauma, GI bleeding, ruptured aortic aneurysm or pregnancy-related bleeding.
- **Cardiogenic Shock:**
Heart failure: MI, Cardiomyopathy, Myocardial contusion, Ruptured ventricular / septum / valve / toxins.
- **Distributive Shock:**
Septic
Anaphylactic
Neurogenic: Hallmark is warm, dry, pink skin with normal capillary refill time and typically alert.
Toxic
- **Obstructive Shock:**
Pericardial tamponade. Pulmonary embolus. Tension pneumothorax.
Signs may include hypotension with distended neck veins, tachycardia, unilateral decreased breath sounds or muffled heart sounds.
- **Acute Adrenal Insufficiency or Congenital Adrenal Hyperplasia:**
Body cannot produce enough steroids (glucocorticoids / mineralocorticoids.) May have primary or secondary adrenal disease, congenital adrenal hyperplasia, or more commonly have stopped a steroid like prednisone. Injury or illness may precipitate. Usually hypotensive with nausea, vomiting, dehydration and / or abdominal pain. **If suspected Paramedic should give Methylprednisolone 125 mg IM / IV / IO or Dexamethasone 10 mg IM / IV / IO. Use steroid agent specific to your drug list. May administer prescribed steroid carried by patient IM / IV / IO. Patient may have Hydrocortisone (Cortef or Solu-Cortef). Dose: < 1y.o. give 25 mg, 1-12 y.o. give 50 mg, and > 12 y.o. give 100 mg or dose specified by patient's physician.**



Bites and Envenomations

History

- Type of bite / sting
- Description / photo for identification
- Time, location, size of bite / sting
- Previous reaction to bite / sting
- Domestic vs. Wild
- Tetanus and Rabies risk
- Immunocompromised patient

Signs and Symptoms

- Rash, skin break, wound
- Pain, soft tissue swelling, redness
- Blood oozing from the bite wound
- Evidence of infection
- Shortness of breath, wheezing
- Allergic reaction, hives, itching
- Hypotension or shock


Differential

- Animal bite
- Human bite
- Snake bite (poisonous)
- Spider bite (poisonous)
- Insect sting / bite (bee, wasp, ant, tick)
- Infection risk
- Rabies risk
- Tetanus risk

Call for help / additional resources
Stage until scene safe

If Needed
Carolinas Poison Control
1-800-222-1222

	General Wound Care Procedure
	Immobilize Injury
	Remove any constricting clothing / bands / jewelry
A	IV / IO Procedure <i>if indicated</i>
	Age Appropriate Trauma Protocol(s) TB 4, 5, 6 <i>if indicated</i>
	Age Appropriate Allergic Reaction/ Anaphylaxis Protocol AM 1 / PM 1 <i>if indicated</i>
	Age Appropriate Hypotension / Shock Protocol AM 5 / PM 3 <i>if indicated</i>
	Pain Control Protocol UP 11 <i>if indicated</i>
	Extremity Trauma Protocol TB 4 <i>if indicated</i>



Toxin-Environmental Protocol Section



Bites and Envenomations

Pearls

- **Recommended Exam: Mental Status, Skin, Extremities (Location of injury), and a complete Neck, Lung, Heart, Abdomen, Back, and Neuro exam if systemic effects are noted**
- **Immunocompromised patients are at an increased risk for infection: diabetes, chemotherapy, transplant patients.**
- **Consider contacting the North Carolina Poison Control Center for guidance (1-800-222-1222).**
- **Do not put responders in danger attempting to capture and animal or insect for identification purposes.**
- **Evidence of infection: swelling, redness, drainage, fever, red streaks proximal to wound.**
- **Human bites:**
Human bites have higher infection rates than animal bites due to normal mouth bacteria.
- **Dog / Cat / Carnivore bites:**
Carnivore bites are much more likely to become infected and all have risk of Rabies exposure.
Cat bites may progress to infection rapidly due to a specific bacteria (*Pasteurella multocida*).
- **Snake bites:**
Poisonous snakes in this area are generally of the pit viper family: rattlesnake and copperhead.
Coral snake bites are rare: Very little pain but very toxic. "Red on yellow - kill a fellow, red on black - venom lack."
Amount of envenomation is variable, generally worse with larger snakes and early in spring.
- **Spider bites:**
Black Widow spider bites tend to be minimally painful, but over a few hours, muscular pain and severe abdominal pain may develop (spider is black with red hourglass on belly).
Brown Recluse spider bites are minimally painful to painless. Little reaction is noted initially but tissue necrosis at the site of the bite develops over the next few days (brown spider with fiddle shape on back).



Carbon Monoxide / Cyanide

History

- Smoke inhalation
- Ingestion of cyanide
- Eating large quantity of fruit pits
- Industrial exposure
- Trauma
- Reason: Suicide, criminal, accidental
- Past Medical History
- Time / Duration of exposure

Signs and Symptoms

- AMS
- Malaise, weakness, flu like illness
- Dyspnea
- GI Symptoms; N/V; cramping
- Dizziness
- Seizures
- Syncope
- Reddened skin
- Chest pain

Differential

- Diabetic related
- Infection
- MI
- Anaphylaxis
- Renal failure / dialysis problem
- Head injury / trauma
- Co-ingestant or exposures

	Immediately Remove from Exposure
	Appropriate Airway Protocol(s) 1 - 7 as indicated
	High Flow Oxygen
	Blood Glucose Analysis Procedure
B	12 Lead ECG Procedure
A	IV/ IO Procedure
P	Cardiac Monitor / CO Monitor
	Altered Mental Status Protocol UP 4 if indicated
	Age Appropriate Diabetic Protocol AM 2 / PM 2 if indicated
	Age Appropriate Multiple Trauma Protocol TB 6 Head Injury TB 5 if indicated
	Age Appropriate Hypotension / Shock Protocol AM 5 / PM 3 if indicated

High Suspicion of Cyanide

YES

P

NO

Continue Care
Continue High Flow Oxygen

Monitor and Reasses

Notify Destination or Contact Medical Control

Pearls

- **Recommended exam: Neuro, Skin, Heart, Lungs, Abdomen, Extremities**
- **Scene safety is priority.**
- Consider CO and Cyanide with any product of combustion
- Normal environmental CO level does not exclude CO poisoning.
- Symptoms present with lower CO levels in pregnancy, children and the elderly.
- Continue high flow oxygen regardless of pulse ox readings.



Drowning

History


- Submersion in water regardless of depth
- Possible history of trauma
Slammed into shore wave break
- Duration of submersion / immersion
- Temperature of water or possibility of hypothermia

Signs and Symptoms

- Unresponsive
- Mental status changes
- Decreased or absent vital signs
- Foaming / Vomiting
- Coughing, Wheezing, Rales, Rhonchi, Stridor
- Apnea

Differential

- Trauma
- Pre-existing medical problem
Hypoglycemia
Cardiac Dysrhythmia
- Pressure injury (SCUBA diving)
Barotrauma
Decompression sickness
- Post-immersion syndrome



Toxic-Environmental Protocol Section

Pearls

- **Recommended Exam: Respiratory, Mental status, Trauma Survey, Skin, Neuro**
- **Drowning is the process of experiencing respiratory impairment (any respiratory symptom) from submersion / immersion in a liquid.**
- **Begin with BVM ventilations, if patient does not tolerate then apply appropriate mode of supplemental oxygen.**
- **Ensure scene safety. Drowning is a leading cause of death among would-be rescuers.**
- **When feasible, only appropriately trained and certified rescuers should remove patients from areas of danger.**
- **Regardless of water temperature – resuscitate all patients with known submersion time of ≤ 25 minutes.**
- **Regardless of water temperature – If submersion time ≥ 1 hour consider moving to recovery phase instead of rescue.**
- **Foam is usually present in airway and may be copious, DO NOT waste time attempting to suction. Ventilate with BVM through foam (suction water and vomit only when present.)**
- **Cardiac arrest in drowning is caused by hypoxia, airway and ventilation are equally important to high-quality CPR.**
- **Encourage transport of all symptomatic patients (cough, foam, dyspnea, abnormal lung sounds, hypoxia) due to potential worsening over the next 6 hours.**
- Predicting prognosis in prehospital setting is difficult and does not correlate with mental status. Unless obvious death, transport.
- Hypothermia is often associated with drowning and submersion injuries even with warm ambient conditions.
- Drowning patient typically has <1 – 3 mL/kg of water in lungs (does not require suction) Primary treatment is reversal of hypoxia.
- Spinal motion restriction is usually unnecessary. When indicated it should not interrupt ventilation, oxygenation and / or CPR.



Hyperthermia

History

- Age, very young and old
- Exposure to increased temperatures and / or humidity
- Past medical history / Medications
- Time and duration of exposure
- Poor PO intake, extreme exertion
- Fatigue and / or muscle cramping

Signs and Symptoms

- Altered mental status / coma
- Hot, dry or sweaty skin
- Hypotension or shock
- Seizures
- Nausea

Differential

- Fever (Infection)
- Dehydration
- Medications
- Hyperthyroidism (Thyroid Storm)
- Delirium tremens (DT's)
- Heat cramps, exhaustion, stroke
- CNS lesions or tumors

Temperature Measurement Procedure **if available**

Temperature Measurement should NOT delay treatment of hyperthermia

Remove from heat source to cool environment
Cooling measures
Remove tight clothing
Blood Glucose Analysis Procedure
Age Appropriate Diabetic Protocol AM 2 / PM 2 as indicated

Assess Symptom Severity

HEAT CRAMPS
Normal to elevated body temperature
Warm, moist skin
Weakness, Muscle cramping

HEAT EXHAUSTION
Elevated body temperature
Cool, moist skin
Weakness, Anxious, Tachypnea

HEAT STROKE
Fever, usually > 104°F (40°C)
Hot, dry skin
Hypotension, AMS / Coma

PO Fluids as tolerated
Monitor and Reassess

Age Appropriate Airway Protocol(s) AR 1 - 7 as indicated
Altered Mental Status Protocol UP 4 as indicated
Active cooling measures Target Temp < 102.5° F (39°C)
B 12 Lead ECG Procedure
A IV / IO Procedure
P Cardiac Monitor
A
Age Appropriate Hypotension / Shock Protocol AM 5 / PM 3 as indicated
Monitor and Reassess

	Notify Destination or Contact Medical Control	
--	--	--

Toxic-Environmental Protocol Section



Hyperthermia

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lungs, Neuro**
- **Extremes of age are more prone to heat emergencies (i.e. young and old). Obtain and document patient temperature if able.**
- Predisposed by use of: tricyclic antidepressants, phenothiazines, anticholinergic medications, and alcohol.
- Cocaine, Amphetamines, and Salicylates may elevate body temperatures.
- Intense shivering may occur as patient is cooled.
- **Heat Cramps:**
Consists of benign muscle cramping secondary to dehydration and is not associated with an elevated temperature.
- **Heat Exhaustion:**
Consists of dehydration, salt depletion, dizziness, fever, mental status changes, headache, cramping, nausea and vomiting. Vital signs usually consist of tachycardia, hypotension, and an elevated temperature.
- **Heat Stroke:**
Consists of dehydration, tachycardia, hypotension, temperature $\geq 104^{\circ}\text{F}$ (40°C), and an altered mental status. Sweating generally disappears as body temperature rises above 104°F (40°C). The young and elderly are more prone to be dry with no sweating.
Exertional Heat Stroke:
In exertional heat stroke (athletes, hard labor), the patient may have sweated profusely and be wet on exam. **Rapid cooling takes precedence over transport as early cooling decreases morbidity and mortality.**
If available, immerse in an ice water bath for 5 – 10 minutes. Monitor rectal temperature and remove patient when temperature reaches 102.5°F (39°C). Your goal is to decrease rectal temperature below 104°F (40°C) with target of 102.5°F (39°C) within 30 minutes. Stirring the water aids in cooling.
Other methods include cold wet towels below and above the body or spraying cold water over body continuously.
- **Neuroleptic Malignant Syndrome (NMS):**
Neuroleptic Malignant Syndrome is a hyperthermic emergency which is not related to heat exposure. It occurs after taking neuroleptic antipsychotic medications. This is a rare but often lethal syndrome characterized by muscular rigidity, AMS, tachycardia and hyperthermia.
Drugs Associated with Neuroleptic Malignant Syndrome:
Prochlorperazine (Compazine), promethazine (Phenergan), clozapine (Clozaril), and risperidone (Risperdal) metoclopramide (Reglan), amoxapine (Ascendin), and lithium.
Management of NMS:
Supportive care with attention to hypotension and volume depletion.
Use benzodiazepines such as diazepam or midazolam for seizures and / or muscular rigidity.



Hypothermia / Frostbite

History


- Age, very young and old
- Exposure to decreased temperatures but may occur in normal temperatures
- Past medical history / Medications
- Drug use: Alcohol, barbituates
- Infections / Sepsis
- Length of exposure / Wetness / Wind chill

Signs and Symptoms

- Altered mental status / coma
- Cold, clammy
- Shivering
- Extremity pain or sensory abnormality
- Bradycardia
- Hypotension or shock

Differential

- Sepsis
- Environmental exposure
- Hypothyroidism
- Hypoglycemia
- CNS dysfunction
 - Stroke
 - Head injury
 - Spinal cord injury





Hypothermia / Frostbite

Pearls

- **Recommended Exam: Mental Status, Heart, Lungs, Abdomen, Extremities, Neuro**
- **NO PATIENT IS DEAD UNTIL WARM AND DEAD (Body temperature $\geq 93.2^\circ\text{F}$, 32°C .)**
- **Many thermometers do not register temperature below 93.2°F .**
- **Hypothermia categories:**
 - Mild $90 - 95^\circ\text{F}$ ($32 - 35^\circ\text{C}$)
 - Moderate $82 - 90^\circ\text{F}$ ($28 - 32^\circ\text{C}$)
 - Severe $< 82^\circ\text{F}$ ($< 28^\circ\text{C}$)
- **Mechanisms of hypothermia:**
 - Radiation: Heat loss to surrounding objects via infrared energy (60% of most heat loss.)
 - Convection: Direct transfer of heat to the surrounding air.
 - Conduction: Direct transfer of heat to direct contact with cooler objects (important in submersion.)
 - Evaporation: Vaporization of water from sweat or other body water losses.
- Contributing factors of hypothermia: Extremes of age, malnutrition, alcohol or other drug use.
- If the temperature is unable to be measured, treat the patient based on the suspected temperature.
- **CPR:**
 - Severe hypothermia may cause cardiac instability and rough handling of the patient theoretically can cause ventricular fibrillation. This has not been demonstrated or confirmed by current evidence. Intubation and CPR techniques should not be with-held due to this concern.
 - Intubation can cause ventricular fibrillation so it should be done gently by most experienced person.
 - Below 86°F (30°C) antiarrhythmics may not work and if given should be given at increased intervals. Contact medical control for direction. Epinephrine / Vasopressin can be administered. Below 86°F (30°C) pacing should not utilized.
 - Consider withholding CPR if patient has organized rhythm or has other signs of life. Contact Medical Control.
 - If the patient is below 86°F (30°C) then defibrillate 1 time if defibrillation is required. Deferring further attempts until more warming occurs is controversial. Contact medical control for direction.
 - Hypothermia may produce severe bradycardia so take at least 60 seconds to palpate a pulse.
- **Active Warming:**
 - Remove from cold environment and to warm environment protected from wind and wet conditions.
 - Remove wet clothing and provide warm blankets / warming blankets.
 - Hot packs can be activated and placed in the armpit and groin area if available. Care should be taken not to place the packs directly against the patient's skin.



Marine Envenomations / Injury

History

- Type of bite / sting
- Identification of organism
- Previous reaction to marine organism
- Immunocompromised
- Household pet

Signs and Symptoms


- Intense localized pain
- Increased oral secretions
- Nausea / vomiting
- Abdominal cramping
- Allergic reaction / anaphylaxis

Differential

- Jellyfish sting
- Sea Urchin sting
- Sting ray barb
- Coral sting
- Swimmers itch
- Cone Shell sting
- Fish bite
- Lion Fish sting

If Needed
Carolinas Poison Control
1-800-222-1222

	General Wound Care Procedure
A	IV / IO Procedure <i>if indicated</i>
P	Cardiac Monitor <i>if indicated</i>
	Drowning Protocol TE 3 <i>if indicated</i>
	Age Appropriate Allergy / Anaphylaxis Protocol AM 1 / PM 1 <i>if indicated</i>
	Age Appropriate Hypotension / Shock Protocol AM 5 / PM 3 <i>if indicated</i>
	Pain Control Protocol UP 11 <i>if indicated</i>



Toxic-Environmental Protocol Section



Marine Envenomations / Injury

Pearls

- **Ensure your safety: Avoid the organism or fragments of the organism as they may impart further sting / injury.**
- **Priority is removal of the patient from the water to prevent drowning.**
- **Coral:**
 - Coral is covered by various living organisms which are easily dislodged from the structure.
 - Victim may swim into coral causing small cuts and abrasions and the coral may enter to cuts causing little if any symptoms initially.
 - The next 24 – 48 hours may reveal an inflammatory reaction with swelling, redness, itching, tenderness and ulceration.
 - Treatment is flushing with large amounts of fresh water or soapy water then repeating
- **Jelly Fish / Anemone / Man-O-War:**
 - Wash the area with fresh seawater to remove tentacles and nematocysts.
 - Do not apply fresh water or ice as this will cause nematocysts firing as well.
 - Recent evidence does not demonstrate a clear choice of any solution that neutralizes nematocysts.
 - Vinegar (immersion for 30 seconds), 50:50 mixture of Baking Soda and Seawater, and even meat tenderizer may have similar effects.
 - Immersion in warm water for 20 minutes, 110 – 114°F (43 - 46°C), has recently been shown to be effective in pain control.
 - Shaving cream may be useful in removing the tentacles and nematocysts with a sharp edge (card).
 - Stimulation of the nematocysts by pressure or rubbing cause the nematocyst to fire even if detached from the jellyfish.
 - Lift away tentacles as scrapping or rubbing will cause nematocysts firing.
 - Typically symptoms are immediate stinging sensation on contact, intensity increases over 10 minutes.
 - Redness and itching usually occur.
 - Papules, vesicles and pustules may be noted and ulcers may form on the skin.
 - Increased oral secretions and gastrointestinal cramping, nausea, pain or vomiting may occur.
 - Muscle spasm, respiratory and cardiovascular collapse may follow.
- **Lionfish:**
 - In North Carolina this would typically occur in the home as they are often kept as pets in saltwater aquariums.
 - Remove any obvious protruding spines and irrigate area with copious amounts of saline.
 - The venom is heat labile so immersion in hot water, 110 – 114 degrees for 30 to 90 minutes is the treatment of choice but do not delay transport if indicated.
- **Stingrays:**
 - Typical injury is swimmer stepping on ray and muscular tail drives 1 – 4 barbs into victim.
 - Venom released when barb is broken.
 - Typical symptoms are immediate pain which increases over 1 – 2 hours. Bleeding may be profuse due to deep puncture wound.
 - Nausea, vomiting, diarrhea, muscle cramping and increased urination and salivation may occur.
 - Seizures, hypotension and respiratory or cardiovascular collapse may occur.
 - Irrigate wound with saline. Extract the spine or barb unless in the abdomen or thorax, contact medical control for advise.
 - Immersion in hot water if available for 30 to 90 minutes but do not delay transport.
- Patients can suffer cardiovascular collapse from both the venom and / or anaphylaxis even in seemingly minor envenomations.
- Sea creature stings and bites impart moderate to severe pain.
- Arrest the envenomation by inactivation of the venom as appropriate.
- Ensure good wound care, immobilization and pain control.



Overdose / Toxic Ingestion

History


- Ingestion or suspected ingestion of a potentially toxic substance
- Substance ingested, route, quantity
- Time of ingestion
- Reason (suicidal, accidental, criminal)
- Available medications in home
- Past medical history, medications

Signs and Symptoms

- Mental status changes
- Hypotension / hypertension
- Decreased respiratory rate
- Tachycardia, dysrhythmias
- Seizures
- S.L.U.D.G.E.
- D.U.M.B.B.E.L.S


Differential

- Tricyclic antidepressants (TCAs)
- Acetaminophen (Tylenol)
- Aspirin
- Depressants
- Stimulants
- Anticholinergic
- Cardiac medications
- Solvents, Alcohols, Cleaning agents
- Insecticides (organophosphates)





Overdose / Toxic Ingestion





Overdose / Toxic Ingestion

Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Extremities, Neuro**
- **Opioids and opiates may require higher doses of Naloxone to improve respiration, in certain circumstances up to 10 mg.**
- **Time of Ingestion:**
 1. Most important aspect is the **TIME OF INGESTION** and the substance and amount ingested and any co-ingestants.
 2. Every effort should be made to elicit this information before leaving the scene.
- **Charcoal Administration:**

The American Academy of Clinical Toxicology DOES NOT recommend the routine use of charcoal in poisonings.

 1. Consider Charcoal within the **FIRST HOUR** after ingestion. If a potentially life threatening substance is ingested or extended release agent(s) are involved and \geq one hour from ingestion contact medical control or Poison Center for direction.
 2. If NG is necessary to administer Charcoal then **DO NOT** administer unless known to be adsorbed, and airway secured by intubation and ingestion is less than **ONE HOUR** confirmed and potentially lethal.
 3. Charcoal in general should only be given to a patient who is alert and awake such that they can self-administer the medication.
- **Do not rely on patient history of ingestion, especially in suicide attempts. Make sure patient is still not carrying other medications or has any weapons.**
- **Pediatric:**
 - Age specific blood pressure 0 – 28 days > 60 mmHg, 1 month - 1 year > 70 mmHg, 1 - 10 years > 70 + (2 x age)mmHg and 11 years and older > 90 mmHg.
 - Maintenance IV Rate: By weight of child: First 10 kg = 4 mL, Second 10 kg = 2 mL, Additional kg = 1 mL. (Example: 36 kg child: First 10 kg = 40 mL, Second 10 kg = 20 mL, 16 kg remaining at 1 mL each. Total is 76 mL / hour)
- **Bring bottles, contents, emesis to ED.**
- **S.L.U.D.G.E: Salivation, Lacrimation, Urination, Defecation, GI distress, Emesis**
- **D.U.M.B.B.E.L.S: Diarrhea, Urination, Miosis, Bradycardia, Bronchorrhea, Emesis, Lacrimation, Salivation.**
- **Tricyclic:** 4 major areas of toxicity: seizures, dysrhythmias, hypotension, decreased mental status or coma; rapid progression from alert mental status to death.
- **Acetaminophen:** initially normal or nausea/vomiting. If not detected and treated, causes irreversible liver failure
- **Aspirin:** Early signs consist of abdominal pain and vomiting. Tachypnea and altered mental status may occur later. Renal dysfunction, liver failure, and or cerebral edema among other things can take place later.
- **Depressants:** decreased HR, decreased BP, decreased temperature, decreased respirations, non-specific pupils
- **Stimulants:** increased HR, increased BP, increased temperature, dilated pupils, seizures
- **Anticholinergic:** increased HR, increased temperature, dilated pupils, mental status changes
- **Cardiac Medications:** dysrhythmias and mental status changes
- **Solvents:** nausea, coughing, vomiting, and mental status changes
- **Insecticides:** increased or decreased HR, increased secretions, nausea, vomiting, diarrhea, pinpoint pupils
- **Nerve Agent Antidote kits** contain 2 mg of Atropine and 600 mg of pralidoxime in an autoinjector for self administration or patient care. These kits may be available as part of the domestic preparedness for Weapons of Mass Destruction.
- **EMR and EMT may administer naloxone by IN / IM route only and may administer from EMS supply. Agency medical director may require Contact of Medical Control prior to administration and may restrict locally.**
- **When appropriate contact the North Carolina Poison Control Center for guidance, reference Policy 18.**
- Consider restraints if necessary for patient's and/or personnel's protection per the Restraint Procedure.



WMD-Nerve Agent Protocol

History


- Exposure to chemical, biologic, radiologic, or nuclear hazard
- Potential exposure to unknown substance/hazard

Signs and Symptoms

- **S**alivation
- **L**acrimation
- **U**rination; increased, loss of control
- **D**efecation / Diarrhea
- **G**I Upset; Abdominal pain / cramping
- **E**mesis
- **M**uscle Twitching
- Seizure Activity
- Respiratory Arrest

Differential

- Nerve agent exposure (e.g., VX, Sarin, Soman, etc.)
- Organophosphate exposure (pesticide)
- Vesicant exposure (e.g., Mustard Gas, etc.)
- Respiratory Irritant Exposure (e.g., Hydrogen Sulfide, Ammonia, Chlorine, etc.)



Pearls

- **Recommended Exam: Mental Status, Skin, HEENT, Heart, Lungs, Gastrointestinal, Neuro**
- **Follow local HAZMAT protocols for decontamination and use of personal protective equipment.**
- **Adult / Pediatric Atropine Dosing Guides:**
 - Confirmed attack: Begin with 1 Nerve Agent Kit for patients less than 7 years of age, 2 Nerve Agent Kits from 8 to 14 years of age, and 3 Nerve Agent Kits for patients 15 years of age and over.
 - If Triage / MCI issues exhaust supply of Nerve Agent Kits, use pediatric atropines (if available).
 - Usual pediatric doses: 0.5 mg ≤ 40 pounds (18 kg), 1 mg dose if patient weighs between 40 to 90 pounds (18 to 40 kg), and 2 mg dose ≥ 90 pounds (≥ 40 kg).
- Each Nerve Agent Kit contains 600 mg of Pralidoxime (2-PAM) and 2 mg of Atropine.
- **Seizure Activity: Any benzodiazepine by any route is acceptable.**
- For patients with major symptoms, there is no limit for atropine dosing.
- Carefully evaluate patients to ensure they not from exposure to another agent (e.g., narcotics, vesicants, etc.)
- The main symptom that the atropine addresses is excessive secretions so atropine should be given until salivation improves.
- EMS personnel, public safety officers and EMR / EMT may carry, self-administer or administer to a patient atropine / pralidoxime by protocol. Agency medical director may require Contact of Medical Control prior to administration.




Suspected Viral Hemorrhagic Fever Ebola

Evolving Protocol:
Protocol subject to change at any time dependent on changing outbreak locations.

Monitor for protocol updates.

Viral Hemorrhagic Fevers:
Ebola is one of many.




YES →







NO ↓



NO →



YES ↓



**Notify Destination as soon and as discretely as possible
DO NOT ENTER facility with patient until instructed
Follow entry directions from hospital staff**



Special Circumstances Section



Suspected Viral Hemorrhagic Fever Ebola

PARTICULAR ATTENTION MUST BE PAID TO PROTECTING MUCOUS MEMBRANES OF THE EYES, NOSE, and MOUTH FROM SPLASHES OF INFECTIOUS MATERIAL OR SELF INOCULATION FROM SOILED PPE / GLOVES.

THERE SHOULD BE NO EXPOSED SKIN

DONNING PPE: **BEFORE** you enter the patient area.

Recommended PPE

PAPR: A PAPR with a full face shield, helmet, or headpiece. Any reusable helmet or headpiece must be covered with a single-use (disposable) hood that extends to the shoulders and fully covers the neck and is compatible with the selected PAPR.

N95 Respirator: Single-use (disposable) N95 respirator in combination with single-use (disposable) surgical hood extending to shoulders and single-use (disposable) full face shield. If N95 respirators are used instead of PAPRs, careful observation is required to ensure healthcare workers are not inadvertently touching their faces under the face shield during patient care.

Single-use (disposable) fluid-resistant or impermeable gown that extends to at least mid-calf or coverall without integrated hood. Coveralls with or without integrated socks are acceptable.

Single-use (disposable) nitrile examination gloves with extended cuffs. Two pairs of gloves should be worn. At a minimum, outer gloves should have extended cuffs.

Single-use (disposable), fluid-resistant or impermeable boot covers that extend to at least mid-calf or single-use (disposable) shoe covers. Boot and shoe covers should allow for ease of movement and not present a slip hazard to the worker.

Single-use (disposable) fluid-resistant or impermeable shoe covers are acceptable only if they will be used in combination with a coverall with integrated socks.

Single-use (disposable), fluid-resistant or impermeable apron that covers the torso to the level of the mid-calf should be used if Ebola patients have vomiting or diarrhea. An apron provides additional protection against exposure of the front of the body to body fluids or excrement. If a PAPR will be worn, consider selecting an apron that ties behind the neck to facilitate easier removal during the doffing procedure

DOFFING PPE: OUTSIDE OF PPE IS CONTAMINATED! DO NOT TOUCH

1) PPE must be carefully removed without contaminating one's eyes, mucous membranes, or clothing with potentially infectious materials.

Use great care while doffing your PPE so as not to contaminate yourself (e.g. Do not remove your N-95 facemask or eye protection BEFORE you remove your gown). There should be a dedicated monitor to observe donning and doffing of PPE. It is very easy for personnel to contaminate themselves when doffing. A dedicated monitor should observe doffing to insure it is done correctly. Follow CDC guidance on doffing.

2) PPE must be double bagged and placed into a regulated medical waste container and disposed of in an appropriate location.

3) Appropriate PPE must be worn while decontaminating / disinfecting EMS equipment or unit.

3) Re-useable PPE should be cleaned and disinfected according to the manufacturer's reprocessing instructions.

Hand Hygiene should be performed by washing with soap and water with hand friction for a minimum of 20 seconds.

Alcohol-based hand rubs may be used if soap and water are not available.

EVEN IF AN ALCOHOL-BASED HAND RUB IS USED, WASH HANDS WITH SOAP AND WATER AS SOON AS

FEASIBLE.

THE USE OF GLOVES IS NOT A SUBSTITUTE FOR HAND WASHING WITH SOAP & WATER

For any provider exposure or contamination contact occupational health.


If the patient is being transported via stretcher then a disposable sheet can be placed over them.

Pearls

- **Transmission to another individual is the greatest after a patient develops fever. Once there is fever, the viral load in the bodily fluids appears to be very high and thus a heightened level of PPE is required.**
- **Patient contact precautions are the most important consideration.**
- **Incubation period 2-21 days**
- **Ebola must be taken seriously; however using your training, protocols, procedures and proper Personal Protective Equipment (PPE), patients can be cared for safely.**
- When an infection does occur in humans, the virus can be spread in several ways to others. The virus is spread through direct contact (through broken skin or mucous membranes) with a sick person's blood or body fluids (urine, saliva, feces, vomit, and semen) objects (such as needles) that have been contaminated with infected body fluids.
- Limit the use of needles and other sharps as much as possible. All needles and sharps should be handled with extreme care and disposed in puncture-proof, sealed containers. Safety devices must be employed immediately after use.
- **Ebola Information: For a complete review of Ebola go to:**
<http://www.cdc.gov/vhf/ebola/index.html>
<http://www.cdc.gov/vhf/ebola/hcp/interim-guidance-emergency-medical-services-systems-911-public-safety-answering-points-management-patients-known-suspected-united-states.html>



Suspected Viral Hemorrhagic Fever Ebola



- 1) EMS personnel performing decontamination / disinfection should wear recommended PPE
When performing Decontamination EMS Personnel MUST wear appropriate PPE, which includes:
 - Gloves (Double glove)
 - Fluid resistant (impervious) Tyvek Like Full length (Coveralls)
 - Eye protection (Goggles)
 - N-95 face mask
 - Fluid resistant (impervious)-Head covers
 - Fluid resistant (impervious)-Shoe / Boot covers
- 2) Face protection (N-95 facemask with goggles) should be worn since tasks such as liquid waste disposal can generate splashes.
- 3) Patient-care surfaces (including stretchers, railings, medical equipment control panels, and adjacent flooring, walls and work surfaces) are likely to become contaminated and should be decontaminated and disinfected after transport.
- 4) A blood spill or spill of other body fluid or substance (e.g., feces or vomit) should be managed through removal of bulk spill matter, cleaning the site, and then disinfecting the site. For large spills, a chemical disinfectant with sufficient potency is needed to overcome the tendency of proteins in blood and other body substances to neutralize the disinfectant's active ingredient. An EPA-registered hospital disinfectant with label claims for viruses that share some technical similarities to Ebola (such as, norovirus, rotavirus, adenovirus, poliovirus) and instructions for cleaning and decontaminating surfaces or objects soiled with blood or body fluids should be used according to those instructions.
(Alternatively, a 1:10 dilution of household bleach (final working concentration of 500 parts per million or 0.5% hypochlorite solution) that is prepared fresh daily (i.e., within 12 hours) can be used to treat the spill before covering with absorbent material and wiping up. After the bulk waste is wiped up, the surface should be disinfected as described in the section above).
- 5) Contaminated reusable patient care equipment should be placed in biohazard bags (double-bagged) and labeled for decontamination and disinfection.
- 6) Reusable equipment should be cleaned and disinfected according to manufacturer's instructions by appropriately trained personnel wearing correct PPE.
- 7) Avoid contamination of reusable porous surfaces that cannot be made single use. Use only a mattress and pillow with plastic or other covering that fluids cannot get through.
- 8) To reduce exposure, all potentially contaminated textiles (cloth products) should be discarded. This includes non-fluid-impermeable pillows or mattresses. They should be considered regulated medical waste and placed in biohazard red bags. They must be double-bagged prior to being placed into regulated medical waste containers.

Pearls

- **Ebola Information:** For a complete review of Ebola EMS Vehicle Disinfection go to:
<http://www.cdc.gov/vhf/ebola/hcp/interim-guidance-emergency-medical-services-systems-911-public-safety-answering-points-management-patients-known-suspected-united-states.html>

Suspected Viral Hemorrhagic Fever Ebola

Decedent Known or suspected carrier of HVF / Ebola Requires Transportation

Only personnel trained in handling infected human remains, and wearing full PPE, should touch, or move any Ebola-infected remains.
Handling human remains should be kept to a minimum.

Donning / Doffing PPE

PPE should be in place BEFORE contact with the body

- 1) Prior to contact with body, postmortem care personnel must wear PPE consisting of: surgical scrub suit, surgical cap, impervious Tyvek-Coveralls, eye protection (e.g., face shield, goggles), facemask, shoe covers, and double surgical gloves.
- 2) Additional PPE (leg coverings,) might be required in certain situations (e.g., copious amounts of blood, vomit, feces, or other body fluids that can contaminate the environment).

PPE should be removed immediately after and discarded as regulated medical waste.

- 1) Use caution when removing PPE as to avoid contaminating the wearer.
- 2) Hand hygiene (washing your hands thoroughly with soap and water or an alcohol based hand rub) should be performed immediately following the removal of PPE. If hands are visibly soiled, use soap and water.

Preparation of Body Prior to Transport

- 1) At the site of death, the body should be wrapped in a plastic shroud. Wrapping of the body should be done in a way that prevents contamination of the outside of the shroud.
- 2) Change your gown or gloves if they become heavily contaminated with blood or body fluids.
- 3) Leave any intravenous lines or endotracheal tubes that may be present in place.
- 4) Avoid washing or cleaning the body.
- 5) After wrapping, the body should be immediately placed in a leak-proof plastic bag not less than 150 μ m thick and zippered closed. The bagged body should then be placed in another leak-proof plastic bag not less than 150 μ m thick and zippered closed before being transported to the morgue.

Surface Decontamination

- 1) Prior to transport to the morgue, perform surface decontamination of the corpse-containing body bags by removing visible soil on outer bag surfaces with EPA-registered disinfectants which can kill a wide range of viruses.
- 2) Follow the product's label instructions. Once the visible soil has been removed, reapply the disinfectant to the entire bag surface and allow to air dry.
- 3) Following the removal of the body, the patient room should be cleaned and disinfected.
- 4) Reusable equipment should be cleaned and disinfected according to standard procedures.

Transportation of VHV / Ebola Remains

PPE is required for individuals driving or riding in a vehicle carrying human remains. DO NOT handle the remains of a suspected / confirmed case of Ebola. The remains must be safely contained in a body bag where the outer surface of the body bag has been disinfected prior to the transport.

Pearls

- **Ebola Information:** For a complete review of Handling Remains of Ebola Infected Patients go to: <http://www.cdc.gov/vhf/ebola/hcp/guidance-safe-handling-human-remains-ebola-patients-us-hospitals-mortuaries.html>

SC 1

Any local EMS System changes to this document must follow the NC OEMS Protocol Change Policy and be approved by OEMS



High Consequence Pathogens (Respiratory Diseases, SARS, MERS-CoV, COVID-19)

EMD Dispatch Center Screening

1. All calls requiring response from EMS system:

Ask: Do you have FEVER AND/OR RESPIRATORY SYMPTOMS?
(cough, breathing difficulty, or other respiratory symptoms?)

EMD Systems:

PDS – Card 36 Pandemic Flu

APCO – COVID-19 Pandemic Vital Points Card

PowerPhone – Pandemic Influenza Card

Evolving Protocol:

Protocol subject to change at any time dependent on changing outbreak locations.

Monitor for protocol updates.

EMD Screen Positive

EMD Screen Negative

Negative
FR or EMS Screening

Exit to
Appropriate
Protocol(s)

Positive
FR or EMS Screening

EMS PPE

EMS
General Treatment
Considerations

Exit to
Appropriate Protocol(s)

PPE Supply Chain Disruptions:

- Prioritize respirators (N95 or equivalent) to aerosol-generating procedures until supply chain restored.
- Prioritize gowns to aerosol-generating procedures.
- It is reasonable for providers to wear a facemask during their duty-shift and change only when soiled or damaged. Adjust use based on supply chain.

Special Circumstances Section



High Consequence Pathogens (Respiratory Diseases, SARS, MERS-CoV, COVID-19)

Pearls

- **First Responders: Because community spread is now present, every patient contact should be considered to have potential for infection with COVID-19. Limit number of FR when caring for patients to limit exposures and PPE use.**
- **Place facemask on any patient complaining of respiratory problems with or without a fever.**
- **Dispatch Screening:**
 - If caller interrogation results in positive screen first responders are assigned based on local agency direction.
 - This screening process will result in many False Positive screens in order to be very sensitive.
- **First Responder and EMS Screening:**
 - Limit distance initially to ≥ 6 feet and conduct a quick screening using the EMD specific question. If this results in a positive screen, immediately place a facemask on the source patient and all providers don appropriate PPE and limit provider number to that which necessary for patient care.
- **Close Contact and Duration Definition:**
 - Healthcare provider exposure is defined as being within 6 feet for ≥ 15 minutes in a patient with suspected illness.
 - Unprotected (no or incorrect PPE) with direct contact with body fluids, including respiratory generated body fluids.
- **Transport:**
 - Occupants in cab of vehicle all should wear facemasks. Riders should be discouraged in order to limit PPE use.
 - Limit number of providers in vehicle required to provide patient care in order to limit exposures.
 - Ensure use of correct PPE for crew and passengers when aerosol-producing procedures utilized.
- **Recommend facemask and gloves with every patient contact. It is reasonable to wear eye protection on every patient contact.**
- **Reasonable to wear simple/surgical mask during entire duty-shift when not able to maintain social distance of ≥ 6 feet among fellow providers when not engaged in patient care.**
- **Negative Pressure in care compartment:**
 - Door or window available to separate driver's and care compartment space:**
 - Close door/window between driver's and care compartment and operate rear exhaust fan on full.
 - No door or window available to separate driver's and care compartment space:**
 - Open outside air vent in driver's compartment and set rear exhaust fan to full.
 - Set vehicle ventilation system to non-recirculating to bring in maximum outside air.
 - Use recirculating HEPA ventilation system if equipped.
- **Airborne precautions:**
 - Standard PPE with fit-tested N95 mask (or PAPR respirator) and utilization of a gown or coveralls, change of gloves after every patient contact, and strict hand washing precautions. This level is utilized with Aspergillus, SARS/MERS/COVID-19, Tuberculosis, Measles (rubeola) Chickenpox (varicella-zoster), Smallpox, Influenza, disseminated herpes zoster, or Adenovirus/Rhinovirus.
- **Contact precautions:**
 - Standard PPE with utilization of a gown or coveralls, change of gloves after every patient contact, and strict hand washing precautions.
 - This level is utilized with GI complaints, blood or body fluids, C diff, scabies, wound and skin infections, MRSA.
 - Clostridium difficile (C diff) is not inactivated by alcohol-based cleaners and washing with soap and water is indicated.
- **Droplet precautions:**
 - Standard PPE plus a standard surgical mask for providers who accompany patients in the treatment compartment and a surgical mask or NRB O2 mask for the patient.
 - This level is utilized when Influenza, Meningitis, Mumps, Streptococcal pharyngitis, Pertussis, Adenovirus, Rhinovirus, and undiagnosed rashes.
- **All-hazards precautions:**
 - Standard PPE plus airborne precautions plus contact precautions.
 - This level is utilized during the initial phases of an outbreak when the etiology of the infection is unknown or when the causative agent is found to be highly contagious (e.g. SARS, MERS-CoV, COVID-19).
- **COVID-19 (Novel Coronavirus): For most current criteria to guide evaluations of patients under investigation:**
 - <https://www.cdc.gov/coronavirus/2019-ncov/symptoms-testing/symptoms.html>

High Consequence Pathogens

(Respiratory Diseases, SARS, MERS-CoV, COVID-19)

Decontamination Recommendations

EMS Personnel Requires Decontamination

Driver:

- Should wear full PPE as described when caring for patient.
- Remove all PPE, except respiratory (N95, PAPR, or equivalent) and perform hand hygiene prior to entering cab to prevent contamination of driver's compartment. **Cab occupants only need to wear facemasks if respirator not already used.**

Wash hands:

- Thoroughly after transferring patient care and/or cleaning ambulance

Maintain records:

- All prehospital providers exposed to patient at the scene and during ambulance transport (self-monitoring for symptoms for 14 days is recommended, even if wearing appropriate PPE).
This does not mean the providers can no longer work.
- List all prehospital provider names (students, observers, supervisors, first response etc.) in the Patient Care Report.

EMS Equipment / Transport Unit Requires Decontamination

Safely clean vehicles used for transport:

- Follow standard operating procedures for the containment and disposal of regulated medical waste.
- Follow standard operating procedures for containing and reprocessing used linen.

Wear appropriate PPE when:

- Removing soiled linen from the vehicle. Avoid shaking the linen.
- Clean and disinfect the vehicle in accordance with agency standard operating procedures.
- Personnel performing the cleaning should wear a disposable gown and gloves (a respirator should not be needed) during the clean-up process; the PPE should be discarded after use.
- All surfaces that may have come in contact with the patient or materials contaminated during patient care (e.g., stretcher, rails, control panels, floors, walls, work surfaces) should be thoroughly cleaned and disinfected using an **EPA-registered disinfectant** appropriate for SARS, MERS-CoV, or coronavirus in healthcare settings in accordance with manufacturer's recommendations. **Keep doors open to patient care compartment while cleaning to allow air exchanges.**

EMS Provider Exposure Risk and Monitoring Recommendations

Close Contact				Close Contact			
Less than 6 feet for ≥ 15 minutes				Less than 6 feet for ≥ 15 minutes			
Source patient NOT WEARING A MASK				Source patient WEARING A MASK			
PPE Utilized	Exposure Risk	Monitoring	Work Restrictions	PPE Utilized	Exposure Risk	Monitoring	Work Restrictions
NONE	HIGH	Self-monitor Supervision	If symptomatic: Fever and Respiratory symptoms (cough, difficulty breathing or other respiratory symptoms) THEN Exclude from work: • At least 72 hours after fever resolution with no use of fever reducing medications. AND • At least 10 days since symptom onset.	NONE	MEDIUM	Self-monitor Supervision	If symptomatic: Fever and Respiratory symptoms (cough, difficulty breathing or other respiratory symptoms) THEN Exclude from work: • At least 72 hours after fever resolution with no use of fever reducing medications. AND • At least 10 days since symptom onset.
No facemask N95 or PAPR	HIGH			No facemask N95 or PAPR	MEDIUM		
No Eye Protection	MEDIUM			No Eye Protection	LOW		
No Gown/ Coveralls or Gloves	LOW			No Gown/ Coveralls or Gloves	LOW		
All recommended PPE Except facemask instead of N95 or PAPR	LOW			All recommended PPE Except facemask instead of N95 or PAPR	LOW		

Placing a simple/surgical mask on the patient within 15 minutes of contact decreases exposure risk.

Return to Work Practice and Work Restrictions (if excluded from work OR exposure to suspected or known COVID-19 patient):

- Prior to duty shift, measure temperature and assess for illness symptoms either by provider, infection control officer, or occupational or public health.
- Self-monitoring with oversight by agency's infection control officer, occupation or public health department per agency policy.
- Wear mask at all times and restrict care of immunocompromised patients (Cancer, Transplant, Steroid use) until all symptoms have resolved or 14 days after onset of illness, whichever is longest.
- Social distance: Employee should maintain 6 feet of separation as work duties permit in the workspace.
- Remove from work if employee becomes symptomatic.

- <https://www.cdc.gov/coronavirus/2019-ncov/hcp/guidance-for-ems.html>
- <https://www.cdc.gov/coronavirus/2019-ncov/healthcare-facilities/hcp-return-work.html>
- <https://www.epa.gov/pesticide-registration/list-n-disinfectants-use-against-sars-cov-2-covid-19>

Special Circumstances Section

High Consequence Pathogens (Respiratory Diseases, SARS, MERS-CoV, COVID-19)

First Responder Guidance

COVID-19 Declared Pandemic with both State and Federal Emergencies Declared

- Many systems are heavily dependent on First Responder agencies to supplement critical prehospital medical care services.
- Community spread is now evident both in NC and in the US.
- Every patient, regardless of medical or injury complaint, is at risk of COVID-19 and all should undergo routine screening questions.
- While EMD is a first step, all providers must screen every patient contact and don appropriate PPE based on clinical situation and COVID-19 screening.
- The citizens we serve continue to have a variety of illness and injury unrelated to COVID-19.
- Limiting PPE use:
First Responders should consider staging with all incidents and sending 1 provider (or more dependent on situation) into the scene to assess for fever and respiratory complaints.

Request staged resources as needed only to provide necessary medical care.

Where patients do not require immediate intervention, first responders may stay in contact with patient, but remain beyond 6 feet until EMS providers arrive to begin assessment and further care.

Consider calling patient on mobile phones to maintain contact and provide reassurance and explain current situation.

PPE Crisis or Alternative Strategies

N95 Respirators

- Use only for aerosol generating procedures (Nebulizer, NIPPV, Suctioning, BVM, BIAD, Intubation).
- Use facemasks in all other scenarios.
- Use respirators (N95 or equivalent) beyond the manufacturing expiration date when not soiled, ripped, torn, or otherwise damaged. Securing straps should also be in good repair and operational:
Visually inspect straps, nose bridge/foam, and mask in general.
Perform seal check: <https://www.youtube.com/watch?v=pGXiUyAoEd8>
- Models tested by CDC and are believed to function properly beyond expiration date:
3M: 1860, 1860s, 1870, 8210, 9010, 8000 Medline/Alpha Protech NON27501
Gerson 1730 Moldex: 1512, 2201
- Minimize providers caring for patient to the extent possible to conserve.
- Use Self-Contained Breathing Apparatus (SCBA) if needed.
- Re-use respiratory (N95 or equivalent) masks and place in paper bag between use. Do not touch inside of mask. Wash hands thoroughly before removing mask.
- When to discard a respirator (N95 or equivalent):
After using during an aerosol producing procedure.
Contamination with blood, body fluids or secretions, following close contact with known COVID-19 patient.


Gowns:

- Use only for aerosol generating procedures (Nebulizer, NIPPV, Suctioning, BVM, BIAD, Intubation).
- Use only for close patient contact, lifting, moving, or transferring where provider contacts patients body.
- May use removable and washable coveralls.

<https://www.cdc.gov/coronavirus/2019-ncov/hcp/respirators-strategy/index.html>



Hospice or Palliative Care Patient (Optional)



Special Circumstances Section



Hospice or Palliative Care Patient (Optional)

Acute Pain / Air Hunger:

SEVERITY	MEDICATION		
	Morphine (IV/IM/SQ)	Dilaudid (IV/IM/SQ)	Fentanyl (IV/IM/SQ)
Mild	2 mg	0.5 mg	25 mcg
Moderate	4 mg	1 mg	50 mcg
Severe	8 mg	2 mg	100 mcg
Titration	2 mg q 15 minutes IV	0.5 mg q 15 minutes IV	25 mcg q 15 minutes IV

Due to pain associated with IM injection, IM administration should only be used if alternative medications or routes of administration are not available. PICC lines may be accessed for use by EMS with sterile technique. May access port-a-cath if appropriate equipment is available and provider is trained.

If using IM or SQ injections, delay repeat dosing by 30 minutes to prevent dose stacking.

Consider using moderate / severe dose in opiate tolerant patients:

Opiate tolerant patients have typical daily dose of narcotic is equivalent to ≥ 60 mg of oral Morphine per day (60 OME (Oral Morphine Equivalents)).

Examples of opiate dosages equivalent to 60 mg of oral Morphine:

- | | |
|--------------------------------|--------------------------|
| 40 mg/day of Oxycodone | 60 mg/day Hydrocodone |
| 25 mcg/hr Fentanyl Transdermal | 15 mg/day of Methadone |
| 200 mg/day of Tapentadol | 16 mg/day of Oxymorphone |
| Suboxone | |

Consider total use of multiple types of opiates. If in doubt about the patient's level of opiate tolerance, or amount of total daily opiate use, treat with a lower initial dose of opiate.

Anxiety / Agitation:

SEVERITY	MEDICATION			
	Ativan (IV/IM/SQ)	Versed (IV/IM/SQ)	Valium (IV/IM/SQ)	Haldol (IV/IM/SQ)
Mild / Moderate	0.5 mg	1 mg	2 mg	2 mg
Severe	1 mg	2 mg	5 mg	4 mg

May repeat dose in 15 minutes for IV administration, or 30 minutes for IM or SQ injections.

Nausea / Vomiting:

Zofran IV / IM	Phenergan IV / IM	Haldol IV / IM / SQ	Ativan IV / IM / SQ
4 mg	25 mg	2 mg	0.5 mg

Pearls

- MOST form section A and DNR forms are equivalent – if valid, Do Not Resuscitate.**
- MOST form and DNR forms may be revoked by Health Care Power of Attorney or other appropriate surrogate decision-makers.**
- Palliative care is specialized care for patients with a chronic and/or terminal illness which focuses on managing symptom exacerbation and the stress of illness.
- Hospice care is specialized care (similar to palliative care) for patients within the last 6 months of life.
- Hospice patient may not have a DNR or MOST form completed and still be enrolled in Hospice care.
- Emergency Kits (eKit):**
May be given to patient by Hospice to use at home for acute symptom exacerbation. Each eKit is individualized and will be different for each patient but typically addresses pain, nausea/vomiting, anxiety, and/or secretions. (EMS is able to administer if within provider's scope of practice.)
- Interaction on-scene with Hospice personnel:**
Hospice nurses are valuable resource in helping patients/families make care/transport decisions. EMS should discuss care/transport decision with hospice nurse. After medication administration, if no transport occurs, care may be transferred to Hospice nurse.



Mass Vaccination/Immunization Medication Distribution

History

- Follow local public health department criteria for specific immunization or medication administered.
- Patient receiving medication or vaccination must be without evidence of active infection.
- AEMT and Paramedic providers may participate
- EMT may participate when DHHS/NCMB allows special provision during local or state emergency.

Situation

- Local implementation of this protocol must be done as a component of the EMS system's local public health department community immunization or medication distribution program.
- May initiate protocol when a community has limited public health department resources or when local or state health emergency is declared.

Review immunization/vaccination or medication guide provided by the local public health department:

- | | |
|----------|---|
| A | <ul style="list-style-type: none"> • Patient selection criteria per local public health department (may vary) • Vaccine/immunization or medication indications • Vaccine/immunization or medication contraindications • Vaccine/immunization or medication distribution procedure |
| B | <ul style="list-style-type: none"> • EMT may provide vaccinations when DHHS/NCMB allows special provision during local or state emergency. |

Confirm patient eligibility for the vaccination or medication including:

- Age
- Medical history
- Contraindications
- Allergies

Eligibility confirmed?

NO

Do not administer:

- Refer to local public health department providers/officials for further care and instructions.

Allergic Reaction or Complications



- Exit to age appropriate Protocol(s)
- Notify appropriate local public health department provider/official

Administer vaccination or medication:

- | | |
|----------|--|
| A | <ul style="list-style-type: none"> • Dose dependent on local public health department • Route dependent on local public health department (PO, IN, IM, IV, SQ) |
| B | <h3><u>Administer Over-the-Counter medication and/or vaccination (if applicable):</u></h3> <ul style="list-style-type: none"> • Undergo specific "just-in-time" training • Dose dependent on local public health department • Route dependent on local public health department (PO, IN, IM). SQ when specified by NCOEMS. • Complete required local public health department documentation • Provide post immunization or medication written instructions and monitoring |

Administer Over-the-Counter medication and/or vaccination (if applicable):

Pearls

- **Purpose:**
Provide protocol driven process for EMS providers to assist with public health immunization or medication distribution initiatives.
- **Documentation of the immunization or medication:**
Complete using local public health department approved record system.
Creation of an EMS patient care report is not required and is not required to submit to NCOEMS.
Must create a log of all patient contacts associated with the immunization or medication distribution program maintained by the EMS system.
If local public health department is maintaining a log of all patients, EMS may use the public health log and keep copies in the EMS system.
- **Injection site:**
Most common injection site for subcutaneous is tissue of an upper arm; follow procedure USP-4 otherwise.
Injection volume is limited to 1 - 2 mL per site unless specific guidance is given per local public health department.
Most common sites for intramuscular injections are upper arm, buttocks, and thighs, follow procedure USP-4.
Injection volume is limited to 1 mL in the upper arm, unless specific guidance is given per local public health department; follow procedure USP-4 otherwise.
Injection volume is limited to 2 mL (1 mL in pediatrics) in buttocks and thighs, unless specific guidance is given per local public health department; follow procedure USP-4 otherwise.



Scene Rehabilitation: General (Optional)

Pearls

- **This protocol is optional and given only as an example. Agencies may and are encouraged to develop their own.**
- **Rehabilitation officer has full authority in deciding when responders may return to duty and may adjust rest / rehabilitation time frames depending on existing conditions.**
- **Rehabilitation goals:**
 - Relief from climatic conditions.**
 - Rest, recovery, and hydration prior to incident, during, and following incident.**
 - Active and / or passive cooling or warming as needed for incident type and climate conditions.**
- **May be utilized with adult responders on fire, law enforcement, rescue, EMS and training scenes.**
- **Responders taking anti-histamines, blood pressure medication, diuretics or stimulants are at increased risk for cold and heat stress.**
- **General indications for rehabilitation:**
 - 20-minute rehabilitation following use of a second 30-minute SCBA, 45-minute SCBA or single 60-minute SCBA cylinder.
 - 20-minute rehabilitation following 40 minutes of intense work without SCBA.
- **General work-rest cycles:**
 - 10-minute self-rehabilitation following use of one 30-minute SCBA cylinder or performing 20 minutes of intense work without SCBA.
- **Serious signs / symptoms:**
 - Chest pain, dizziness, dyspnea, weakness, nausea, or headache.
 - Symptoms of heat stress (cramps) or cold stress.
 - Changes in gait, speech, or behavior.
 - Altered Mental Status.
 - Abnormal Vital Signs per agency SOP or Policy / Procedure.
- **Rehabilitation Section:**
 - Integral function within the Incident Management System.
 - Establish section such that it provides shelter / shade, privacy and freedom from smoke or other hazards
 - Large enough to accommodate expected number of personnel.
 - Separate area to remove PPE.
 - Accessible to EMS transport units and water supply.
 - Away from media agencies and spectators / bystanders.



Scene Rehabilitation: Responder (Optional)

Remove:

PPE
Body Armor
Chemical Suits
SCBA
Turnout Gear
Other equipment as indicated


Continue:

Heat and Cold Stress treatment techniques from General Rehab Section

Injury / Illness / Complaint should be treated using appropriate treatment protocol beyond need for oral or IV hydration.



NFA Age Predicted 85 % Maximum Heart Rate	
20 - 25	170
26 - 30	165
31 - 35	160
36 - 40	155
41 - 45	152
46 - 50	148
51 - 55	140
55 - 60	136
61 - 65	132



Special Operations Section

Pearls

- This protocol is optional and given only as an example. Agencies may and are encouraged to develop their own.
- Rehabilitation officer has full authority in deciding when responders may return to duty.
- Utilized when responder is not appropriate for General Rehabilitation Protocol.
- May be utilized with adult responders on fire, law enforcement, rescue, EMS and training scenes.
- Responders taking anti-histamines, blood pressure medication, diuretics or stimulants are at increased risk for cold and heat stress.
- Rehabilitation Section is an integral function within the Incident Management System.
- Establish section such that it provides shelter, privacy and freedom from smoke or other hazards.



Table of Contents

Policy

Disposition Policy Section

1. Criteria for Death or Withholding Resuscitation
2. Deceased Subjects
3. Discontinuation of Prehospital Resuscitation
4. Disposition (Patient Instructions)
5. DNR and MOST
6. Patient Without A Protocol
7. Physician on Scene
8. Opioid Overdose Misuse
9. Organ Procurement Agency Notification

Documentation Policy Section

1. Documentation and Data Quality
2. Documentation of Vital Signs

EMS Dispatch Policy Section

1. EMS Dispatch Center Time

Medical Policy Section

1. Ketamine Program Requirements

Pediatric Policy Section

1. Children With Special Healthcare Needs (NC Kidbase)
2. Infant Abandonment

Service Metric Policy Section

1. EMS Back In Service Time
2. EMS Wheels Rolling Turn-Out Time

Toxic Environmental Policy Section

1. Poison Control

Transport Policy Section

1. Air Transport
2. Safe Transport of Pediatric Patients
3. Transport



Criteria for Death / Withholding Resuscitation

Policy:

CPR, BLS and ALS treatment are to be withheld only if the patient is obviously dead (see procedure section) or a valid (*properly completed, signed, dated, and unexpired*) **North Carolina Do Not Resuscitate (DNR)** form and/or **Medical Orders for Scope of Treatment (MOST)** form is present (Disposition Policy 5).

EMS personnel shall also honor a valid **POLST (Physician Orders for Life Sustaining Treatment)**, **POST (Physician Orders for Scope of Treatment)**, **MOST and/or DNR** (*properly completed, signed, dated, and unexpired*) from another state or US military form. NCGS Article 23: 90-320.

Purpose:

The purpose of this policy is to:

- Honor those who have obviously expired prior to EMS arrival.
- To honor the terminal wishes of the patient
- To prevent the initiation of unwanted resuscitation

Procedure:

1. If a patient is in complete cardiopulmonary arrest (clinically dead) and meets one or more of the criteria below, CPR and ALS therapy need not be initiated:
 - Body decomposition
 - Rigor mortis
 - Dependent lividity
 - Blunt force trauma
 - Injury not compatible with life (i.e., decapitation, burned beyond recognition, massive open or penetrating trauma to the head or chest with obvious organ destruction)
 - Extended downtime (> 15 minutes) with Asystole on the ECG
 - Meets criteria established in AC 12 Termination of CPR Protocol
 - Meets criteria established in TB 10 Traumatic Arrest Protocol
2. If a bystander or first responder has initiated CPR or automated defibrillation prior to ALS personnel (Paramedic or AEMT) arrival and any of the above criteria (signs of obvious death) are present, the ALS provider may discontinue CPR / resuscitation efforts. All other EMS personnel levels must communicate with medical control prior to discontinuation of the resuscitative efforts unless specifically addressed in AC 12 Termination of CPR Protocol and/or TB 10 Traumatic Arrest Protocol.
3. If doubt exists, start resuscitation immediately. Once resuscitation is initiated, continue resuscitation efforts until either:
 - a) Resuscitation efforts meet the criteria for implementing the **Discontinuation of Prehospital Resuscitation Policy** (Disposition Policy 3).
 - b) Patient care responsibilities are transferred to the destination hospital staff.



Deceased Subjects

Policy:

EMS will handle the disposition of deceased subjects in a uniform, professional, and timely manner.

Purpose:

The purpose of this policy is to:

- Organize and provide for a timely disposition of any deceased subject
- Maintain respect for the deceased and family
- Allow EMS to return to service in a timely manner.

Procedure:

1. Do not remove lines or tubes from unsuccessful cardiac arrests/codes unless directed below.
2. Notify the law enforcement agency with jurisdiction if applicable.
3. If subject was found deceased by EMS, the scene is turned over to law enforcement.
4. If EMS has attempted to resuscitate the patient and then terminated the resuscitative efforts, EMS personnel should contact the primary care provider (medical cases) or medical examiner (traumatic cases or primary care provider unavailable) to provide information about the resuscitative efforts.

Cases that require notification of the Medical Examiner when death results from:

- | | |
|--|-----------|
| Accident | Poisoning |
| Homicide | Suicide |
| Violence | |
| Occurring in jail, prison, correctional institution, or in LEO custody | |
| Occurring under suspicious, unusual, or unnatural circumstances | |
| Sudden unexpected death when in otherwise good health | |
| No current primary care or specialty physician care | |

5. Transport arrangements should be made in concert with law enforcement and the family's wishes.
6. If the deceased subject's death is not under the jurisdiction of the medical examiner, any line(s) or tube(s) placed by EMS should be removed prior to transport.
7. Document the situation, name of primary care provider or Medical Examiner contacted, the patient care report form (PCR).
8. Physician Assistants and/or Nurse Practitioners may sign a North Carolina death certificate when specially authorized by their supervising physician.
9. Follow Disposition Policy 9 Organ Procurement Agency Notification



Discontinuation of Prehospital Resuscitation

Policy:

Unsuccessful cardiopulmonary resuscitation (CPR), basic life support (BLS), and other advanced life support (ALS) interventions may be discontinued prior to transport or arrival at the hospital when this policy is followed.

Purpose:

The purpose of this policy is to:

- Allow for discontinuation of prehospital resuscitation after the delivery of adequate and appropriate BLS and/or ALS therapy.

Procedure:

1. Discontinuation of CPR, BLS, and ALS intervention may be implemented **prior to contact with Medical Control if ALL** of the following criteria have been met:

- Patient must be ≥ 18 years of age
- High quality CPR administered
- Airway successfully managed:
 - Acceptable airway management techniques include orotracheal intubation, Blind Insertion Airway Device (BIAD) placement, or cricothyrotomy
 - EtCO₂ monitoring for airway confirmation utilized if available
- IV or IO access has been achieved
- No hypothermia (body temperature $\geq 93.2^{\circ}\text{F}$ or 32°C)
- Protocol AC 12 On Scene Resuscitation Termination of CPR or TB 10 Traumatic Arrest utilized as applicable
- All EMS BLS and ALS personnel involved in the patient's care agree that discontinuation of the resuscitation is appropriate

2. If all of the above criteria are not met and discontinuation of prehospital resuscitation is desired, **contact Medical Control**.

3. The **Deceased Subjects Policy** should be followed.

Document all patient care and interactions with the patient's family, personal physician, medical examiner, law enforcement, and medical control in the EMS patient care report (PCR).

Disposition (Patient Instructions)

Policy:

All patient encounters responded to by EMS will result in the accurate and timely completion of:

- The Patient Care Report (PCR) for all patients transported by EMS
- The Patient Disposition Form for all patients not transported by EMS

Purpose:

To provide for the documentation of:

- The evaluation and care of the patient
- The patient's refusal of the evaluation, treatment, and/or transportation
- The patient's disposition instructions
- The patient's EMS encounter to protect the local EMS system and its personnel from undue risk and liability.

Procedure:

1. All patient encounters, which result in some component of an evaluation, must have a Patient Care Report completed.
2. All patients who refuse any component of the evaluation or treatment, based on the complaint, must have a Disposition Form completed.
3. All patients who are NOT transported by EMS must have a Disposition (patient instruction) Form completed including the Patient Instruction Section.
4. A copy of the Patient Disposition Form should be maintained with the official Patient Care Report (PCR)



North Carolina Do Not Resuscitate and MOST Form

Policy:

CPR, BLS and ALS treatment are to be withheld only if the patient is obviously dead (see procedure section) or a valid (*properly completed, signed, dated, and unexpired*) **North Carolina Do Not Resuscitate (DNR)** form and/or **Medical Orders for Scope of Treatment (MOST)** form is present (Disposition Policy 5).

EMS personnel shall also honor a valid **POLST (Physician Orders for Life Sustaining Treatment)**, **POST (Physician Orders for Scope of Treatment)**, **MOST and/or DNR** (*properly completed, signed, dated, and unexpired*) from another state or US military form. NCGS Article 23: 90-320.

Purpose:

- Honor those who have obviously expired prior to EMS arrival.
- To honor the terminal wishes of the patient
- To prevent the initiation of unwanted resuscitation

Procedure:

1. When confronted with a patient or situation involving the NC DNR and/or MOST form(s), the following form content must be verified before honoring the form(s) request.
 - The form(s) must be either an original North Carolina DNR or North Carolina MOST form
 - The effective date and expiration date must be completed and current
 - The DNR and/or MOST Form must be signed by a physician, physician's assistant, or nurse practitioner.
 - Out-of-state or US military form:
 - Must be an original MOST, DNR, POLST (Physician Orders for Life Sustaining Treatment) or POST (Physician Orders for Scope of Treatment).
 - The effective date and expiration date must be completed and current
 - The DNR and/or MOST Form must be signed by a physician, physician's assistant, or nurse practitioner
2. A valid DNR or MOST form may be overridden by the request of (N.C.G.S. 90-21.13):
 - Court appointed guardian
 - Health care power of attorney
 - Spouse
 - Majority of patient's reasonably available parents and/or children who are ≥ 18 years old
 - Majority of patient's reasonably available siblings who are ≥ 18 years old
 - Patient's attending physician

*EMS personnel should contact **Medical Control** to obtain assistance and direction if clarification is necessary.*
3. A living will (other legal document) that identifies the patient's desire to withhold CPR or other medical care may be honored with the approval of **Medical Control**. Ideally, consultation with patient's family and personal physician is suggested as time allows.



Standards Policy: Disposition Policy Section

Patient Without a Protocol

Policy:

Anyone requesting EMS services will receive a professional evaluation, treatment, and transportation (if needed) in a systematic, orderly fashion regardless of the patient's problem or condition.

Purpose:

- To ensure the provision of appropriate medical care for every patient regardless of the patient's problem or condition.

Procedure:

1. Treatment and medical direction for all patient encounters, which can be triaged into an EMS patient care protocol, is to be initiated by protocol.
2. When confronted with an emergency or situation that does not fit into an existing EMS patient care protocol, the patient should be treated by the **Universal Patient Care Protocol** and a **Medical Control Physician** should be contacted for further instructions.

Disposition Policy 6

Revised
01/20/2022

Any local EMS System changes to this document must follow the NC OEMS Protocol Change Policy and be approved by OEMS



Physician on Scene

Policy:

The medical direction of prehospital care at the scene of an emergency is the responsibility of those most appropriately trained in providing such care. All care should be provided within the rules and regulations of the state of North Carolina.

Purpose:

- To identify a chain of command to allow field personnel to adequately care for the patient
- To assure the patient receives the maximum benefit from prehospital care
- To minimize the liability of the EMS system as well as the on-scene physician

Procedure:

1. When a non medical-control physician offers assistance to EMS or the patient is being attended by a physician with whom they do not have an ongoing patient relationship, EMS personnel must review the On-Scene Physician Form with the physician. All requisite documentation must be verified and the physician must be approved by on-line medical control.
2. When the patient is being attended by a physician with whom they have an ongoing patient relationship, EMS personnel may follow orders given by the physician if the orders conform to current EMS guidelines, and if the physician signs the PCR. Notify medical control at the earliest opportunity. Any deviation from local EMS protocols requires the physician to accompany the patient to the hospital.
3. EMS personnel may accept orders from the patient's physician over the phone with the approval of medical control. The paramedic should obtain the specific order and the physician's phone number for relay to medical control so that medical control can discuss any concerns with the physician directly.



Opioid Overdose/Misuse (Optional)

Policy:

Patients who have experienced an opioid overdose/misuse should be offered a variety of options to more appropriately manage their care where available in the community. All care should be provided within the rules and regulations of the state of North Carolina.

Purpose:

- To ensure patients are offered options for treatment of opioid misuse where available.
- Provide harm reduction measures related to opioid misuse.

Procedure:

1. Patients must be over 18 years of age and experienced unintentional overdose or misuse of an opioid medication(s) only. Patients must NOT have experienced cardiac arrest defined as administration of chest compressions by first responders or EMS during the incident.
2. The patient must regain a normal mental status and respiratory effort after the administration of naloxone, NOT have suicidal or homicidal ideations/intentions, and NOT ingested substance(s) for intentional self-harm.
3. Patients who have co-ingested other substances should be treated based on appropriate protocol. Consult Carolina Poison Center at 1-800-222-1222 for advice if needed.
4. Transport to an Emergency Department should be offered to all patients. For patients who decline transport to an Emergency Department, alternative destinations should be offered if available in the community. Options may include assistance with accessing inpatient treatment centers, outpatient facilities, mobile crisis solutions, addiction specialists, and/or other local treatment options.
5. In order to decline transport, the patient must meet the following criteria:
 - a) Be 18 years or older
 - b) Maintain a GCS of 15 (alert, and oriented to time, place, person, and situation)
 - c) Demonstrate decision-making capacity as outlined in Universal Protocol (UP 1) Pearls.
6. If patient declines transport to an Emergency Department, an additional dose of naloxone should be offered by EMS if patient consents to additional treatment. IN administration is preferable to limit the possibility of provider needle stick injury. If patient has no sober and responsible party to monitor them, EMS should offer IM administration of naloxone if patient consents to treatment. If available, a naloxone kit should be left with the patient, family, and/or friends on scene. EMS should provide brief education on how to properly use these kits and refer them to read all package related material and instructions provided by the manufacturer.
7. In addition to naloxone kits, the following items should be offered where possible/available:
 - a) Offer to properly dispose of any dirty needles following your agency policy
 - b) Provide clean needles/syringes where possible following your agency policy
 - c) Refer to a community peer support team if available
 - d) Provide literature outlining resources for substance misuse treatment programs in the community



Standards Policy: Disposition Policy Section

Organ Procurement Agency Notification

Policy:

When cardiopulmonary resuscitation (CPR), basic life support (BLS), and other advanced life support (ALS) interventions are withheld or discontinued on scene, EMS will report the death to the appropriate organ procurement organization servicing the county where death occurred in a timely manner. EMS will share information relevant to the donation process with the appropriate organ procurement organization.

Purpose:

To ensure an organ procurement organization is notified of deaths pronounced in the field by EMS in order to:

- Honor the decedent’s registered declaration of eye and/or tissue donation.
- Preserve family’s opportunity to support eye and/or tissue donation.
- Service the public health by facilitating eye and tissue donation.

Procedure:

EMS will notify the appropriate organ procurement organization of deaths pronounced outside of the hospital. Potential donors between ages of newborn – 100 years old will be referred.

Essential information to be provided to the organ procurement organization include:

- Caller name, title, and agency contact information
- Patient demographics
- Last seen alive date/time or time of death
- Circumstances of death (notify organ procurement agency even if medical examiner case)
- Medical interventions and medical history
- Next of kin name and contact information
- Who is taking custody of the decedent’s body (ex: funeral home, hospital, M.E.)
- EMS **SHOULD NOT** discuss eye or tissue donation with next of kin. Coordinators specializing in family support will attempt to contact appropriate family members about organ donation.
- Document all patient care and interactions with the patient’s family, personal physician, medical examiner, law enforcement, and medical control in the EMS electronic patient care report (ePCR).

Contact information for Organ Procurement Organizations:

LifeShare to the Carolinas		LifeNet Health	HonorBridge
1 (800) 932-4483		1 (800) 847-7831	1 (800) 252-2672
Anson	Jackson	Currituck	All other NC counties
Buncombe	Lincoln		
Burke	Macon		
Cabarrus	Madison		
Cherokee	Mecklenburg		
Clay	Polk		
Cleveland	Rutherford		
Gaston	Stanly		
Graham	Swain		
Haywood	Transylvania		
Henderson	Union		



Standards Policy: Documentation Policy Section

EMS Documentation and Data Quality

Policy:

The complete EMS documentation associated with service delivery and patient care shall be electronically recorded into a Patient Care Report (PCR) within 24 hours of the completion of the EMS event, with an EMS Data Score at/or below the state average.

Definition:

EMS documentation of a Patient Care Report (PCR) is based on the appropriate and complete documentation of the EMS data elements as required and defined within the North Carolina College of Emergency Physician's EMS Standards (www.NCCEP.org). Since each EMS event and/or patient scenario is unique, only the data elements relevant to that EMS event and/or patient scenario should be completed.

The EMS Data Score is calculated on each EMS PCR as it is electronically processed into the North Carolina PreHospital Medical Information System (PreMIS). Data Quality Scores are provided within PreMIS. The best possible score is a 0 (zero) and with each data quality error a point is added to the data quality score.

A complete Patient Care Report (PCR) must contain the following information (as it relates to each EMS event and/or patient):

- Service delivery and crew information regarding the EMS Agency's response
- Dispatch information regarding the dispatch complaint, and EMD card number
- Patient care provided prior to EMS arrival
- Patient assessment as required by each specific complaint based protocol
- Past medical history, medications, allergies, and DNR/MOST status
- Trauma and cardiac arrest information if relevant to the EMS event or patient
- All times related to the event
- All procedures and their associated time
- All medications administered with their associated time
- Disposition and/or transport information
- Communication with medical control
- Appropriate signatures (written and/or electronic)

Purpose:

The purpose of this policy is to:

- Promote timely and complete EMS documentation.
- Promote quality documentation that can be used to evaluate and improve EMS service delivery, personnel performance, and patient care to the county's citizens.
- Promote quality documentation that will decrease EMS legal and risk management liability.
- Provide a means for continuous evaluation to assure policy compliance.



Standards Policy: Documentation Policy Section

EMS Documentation and Data Quality

Procedure:

The following procedures shall be implemented to assure policy compliance:

1. The EMS Patient Care Report (PCR) shall be completed as soon as possible after the time of the patient encounter. **Documentation should be completed prior to leaving the destination facility unless call demand dictates otherwise, in which case documentation must be completed prior to the end of the personnel's shift.**
2. **A copy of the patient care report form SHOULD be provided to the receiving medical facility. If the final PCR is not available at the time the patient is left with the emergency department or other healthcare facility, an interim report such as the PreMIS Preliminary Report Form MUST be provided.**
3. The PCR must be completed in the PreMIS System or electronically submitted to the PreMIS System within 24 hours of the EMS event or patient encounter's completion. The EMS data quality feedback provided at the time of the electronic submission into PreMIS should be reviewed and when possible any identified errors will be corrected within each PCR. Each PCR may be electronically resubmitted to PreMIS as many times as needed.
4. The EMS Data Quality Scores for the EMS System, EMS Agency, and individual EMS personnel will be reviewed regularly within the EMS System Peer Review Committee.



Standards Policy: Documentation Policy Section

Documentation of Vital Signs

Policy:

Every patient encounter by EMS will be documented. Vital signs are a key component in the evaluation of any patient and a complete set of vital signs is to be documented for any patient who receives some assessment component.

Purpose:

To insure:

- Evaluation of every patient's volume and cardiovascular status
- Documentation of a complete set of vital signs

Procedure:

1. An **initial** complete set of vital signs includes:
 - Pulse rate
 - Systolic **AND** diastolic blood pressure
 - Respiratory rate
 - Pain / severity (when appropriate to patient complaint)
 - GCS for Injured Patients
2. When no ALS treatment is provided, palpated blood pressures are acceptable for **REPEAT** vital signs.
3. Based on patient condition and complaint, vital signs may also include:
 - Pulse Oximetry
 - Temperature
 - End Tidal CO₂
 - Breath Sounds
 - Level of Response
4. If the patient refuses this evaluation, the patient's mental status and the reason for refusal of evaluation must be documented. A patient disposition form must also be completed.
5. Document situations that preclude the evaluation of a complete set of vital signs.
6. Record the time vital signs were obtained.
7. Any abnormal vital sign should be repeated and monitored closely.

EMS Dispatch Center Time

Policy:

The EMS Dispatch Center Time will be less than 90 seconds, 90% of the time, for all events identified and classified as an emergent or hot (with lights and siren) response.

Definition:

The EMS Dispatch Center Time is defined as the time interval beginning with the time the initial 911 phone call rings at the 911 Communications Center requesting emergency medical services and ending with the dispatch time of the EMS Unit responding to the event.

Purpose:

The purpose of this policy is to:

- Provide the safest and most appropriate level of response to all EMS events within the EMS System.
- Provide a timely and reliable response for all EMS events within the EMS System.
- Provide quality EMS service and patient care to the county's citizens.
- Provide a means for continuous evaluation to assure policy compliance.

Procedure:

The following procedures shall be implemented to assure policy compliance:

1. A public calls into the 911 Communications Center requesting emergency medical assistance will never be required to speak with more than two persons before a formal EMS Unit is dispatched.
2. In EMS Dispatch Centers where Emergency Medical Dispatch (EMD) has been implemented, EMS Units will be dispatched by EMD certified personnel in accordance with the standards developed by the Medical Director and the Emergency Medical Dispatch Protocols.
3. EMS Units will be dispatched hot (with lights and sirens) or cold (no lights and sirens) by the 911 Call Center based on predetermined criteria. If First Responders are dispatched as a component of the EMS response, they should typically be dispatched hot (with lights and sirens).
4. Without question, exception, or hesitation, EMS Units will respond as dispatched (hot or cold). This includes both requests to respond on active calls and requests to "move-up" to cover areas of the System that have limited EMS resources available.
5. EMS Units may, at their discretion, request for a First Responder on Non-First Responder calls in situations where additional resources are required such as manpower, extreme response time of the EMS Unit, need for forcible entry, etc.



EMS Dispatch Center Time

6. EMS Units dispatched with a cold (no lights and sirens) response, will not upgrade to a hot (with lights and sirens) response **UNLESS**:
 - Public Safety personnel on-scene requests a hot (with lights and sirens) response.
 - Communications Center determines that the patient's condition has changed, and requests you to upgrade to a hot (with lights and sirens) response.
7. An EMS Unit may divert from a current cold (no lights and sirens) call to a higher priority hot (with lights and sirens) call **ONLY IF**:
 - The EMS Unit can get to the higher priority call before it can reach the lower priority call. Examples of High Priority Calls: Chest Pain, Respiratory Distress, CVA, etc.
 - The diverting EMS Unit must notify the EMS Dispatch Center that they are diverting to the higher priority call.
 - The diverting EMS Unit ensures that the EMS Dispatch Center dispatches an EMS Unit to their original call.
 - Once a call has been diverted, the next EMS Unit dispatched must respond to the original call. A call cannot be diverted more than one (1) time.
8. Any EMS Dispatch Center Time delays resulting in a prolonged EMS Dispatch Center Time for emergent hot (with lights and sirens) events will be documented in Patient Care Report (PCR) as an "EMS Dispatch Delay" as required and defined in the North Carolina College of Emergency Physicians (NCCEP) EMS Dataset Standards Document.
9. All EMS Dispatch Delays will be reviewed regularly within the EMS System Peer Review Committee.



Standards Policy: Medical Policy Section

Ketamine Program Requirements

Policy:

When administered outside of the AR 3 Airway Drug Assisted Intubation Protocol, an EMS System or Agency must be approved by the State Medical Director and follow the guidelines below when administering Ketamine.

Purpose:

The purpose of this policy is to:

- Ensure that Ketamine is administered in a safe and effective manner
- Facilitate use of Ketamine in appropriate patients
- Establish a reporting mechanism for state review
-

Procedure:

1. The EMS system or Agency must adopt NCCEP protocols unchanged or submit equivalent protocols for review.
2. Letters of support must be obtained from all receiving hospitals where patients will be delivered after administration. These letters must be submitted to the OEMS prior to approval.
3. All personnel must be trained prior to implementation.
4. All administrations must be reviewed through the established PI/QA Medical Oversight process to include hospital outcome feedback. Concerns identified must be reviewed by the Peer Review/QA committee.
5. There are two (2) components of the NCOEMS reporting process:
 - a. The EMS system or agency must submit to the OEMS a Ketamine Adverse Outcome Reporting Form and ePCR within 14 days for administrations that result in any of the following;
 - 1) Cardiac Arrest (pre-hospital or ED)
 - 2) Unanticipated intubation required after administration (pre-hospital or ED).

*Secure Ketamine Adverse Outcome Report link: <https://nc.readyop.com/fs/4ckl/786b>

- b. The EMS system or agency must submit a quarterly report to the OEMS indicating;
 - 1) The total number of administrations
 - 2) Summary of primary protocol utilizations
 - 3) Summary highlighting the PR/QA of cases that required a Ketamine Reporting Form.

*Secure Ketamine Quarterly Report link: <https://nc.readyop.com/fs/4ckG/1544>

****IF THE REPORTING LINKS ABOVE DO NOT DIRECT YOU TO AN ACTIVE FORM, PLEASE COPY AND PASTE THE LINK INTO YOUR WEB BROWSER MANUALLY****



Policy:

Medical technology, changes in the healthcare industry, and increased home health capabilities have created a special population of patients that interface with the EMS system. It is important for EMS to understand and provide quality care to children with special health care needs.

Purpose:

The purpose of this policy is to:

- Provide quality patient care and EMS services to children with special health care needs.
- Understand the need to communicate with the parents and caregivers regarding healthcare needs and devices that EMS may not have experience with.
- Promote, request, and use the “Kidbase” form, which catalogs the health care problems, needs, and issues of each child with a special healthcare need.

Procedure:

1. Caregivers who call 911 to report an emergency involving a child with special health care needs may report that the emergency involves a “Kidbase child” (if they are familiar with the NC Kidbase program) or may state that the situation involves a special needs child.
2. Responding EMS personnel should ask the caregiver of a special needs child for a copy of the “Kidbase Form”, which is the North Carolina terminology for the Emergency Information Form (EIF).
3. EMS personnel may choose to contact the child’s primary care physician for assistance with specific conditions or devices or for advice regarding appropriate treatment and/or transport of the child in the specific situation.
4. Transportation of the child, if necessary, will be made to the hospital appropriate for the specific condition of the child. In some cases this may involve bypassing the closest facility for a more distant yet more medically appropriate destination.



Standards Policy: Pediatric Policy Section

Infant Abandonment

Policy:

The North Carolina Infant Homicide Prevention Act provides a mechanism for unwanted infants to be taken under temporary custody by a law enforcement officer, social services worker, healthcare provider, or EMS personnel if an infant is presented by the parent within 7 days of birth. Emergency Medical Services will accept and protect infants who are presented to EMS in this manner, until custody of the child can be released to the Department of Social Services.

*“A law enforcement officer, a department of social services worker, a health care provider as defined in G.S. 90-21.11 at a hospital or local or district health department, or an **emergency medical technician** at a fire station shall, without a court order, take into temporary custody an infant under 7 days of age that is voluntarily delivered to the individual by the infant's parent who does not express an intent to return for the infant. An individual who takes an infant into temporary custody under this subsection shall perform any act necessary to protect the physical health and well-being of the infant and shall immediately notify the department of social services. Any individual who takes an infant into temporary custody under this subsection may inquire as to the parents' identities and as to any relevant medical history, but the parent is not required to provide this information.”*

Purpose:

To provide:

- Protection to infants that are placed into the custody of EMS under this law
- Protection to EMS systems and personnel when confronted with this issue

Procedure:

1. Initiate the Pediatric Assessment Procedure.
2. Initiate Newly Born Protocol as appropriate.
3. Initiate other treatment protocols as appropriate.
4. Keep infant warm.
5. Call local Department of Social Services or the county equivalent as soon as infant is stabilized.
6. Transport infant to medical facility as per local protocol.
7. Assure infant is secured in appropriate child restraint device for transport.
8. Document protocols, procedures, and agency notifications in the PCR.



EMS Back in Service Time

Policy:

All EMS Units transporting a patient to a medical facility shall transfer the care of the patient and complete all required operational tasks to be back in service for the next potential EMS event within 30 minutes of arrival to the medical facility, 90% of the time.

Definition:

The EMS Back in Service Time is defined as the time interval beginning with the time the transporting EMS Unit arrives at the medical facility destination and ending with the time the EMS Unit checks back in service and available for the next EMS event.

Purpose:

The purpose of this policy is to:

- Assure that the care of each EMS patient transported to a medical facility is transferred to the medical facility staff in a timely manner.
- Assure that the EMS unit is cleaned, disinfected, restocked, and available for the next EMS event in a timely manner.
- Assure that an interim or complete EMS patient care report (PCR) is completed and left with the receiving medical facility documenting, at a minimum, the evaluation and care provided by EMS for that patient (It is acceptable to leave the PreMIS Preliminary Report or equivalent if the final PCR cannot be completed before leaving the facility).
- Provide quality EMS service and patient care to the county's citizens.
- Provide a means for continuous evaluation to assure policy compliance.

Procedure:

The following procedures shall be implemented to assure policy compliance:

1. The EMS Unit's priority upon arrival at the medical facility will be to transfer the care of the patient to medical facility staff as soon as possible.
2. EMS personnel will provide a verbal patient report on to the receiving medical facility staff.
3. EMS personnel will provide an interim (PreMIS Preliminary Report or equivalent) or final Patient Care Report (PCR) to the receiving medical facility staff, prior to leaving the facility, that documents at a minimum the patient's evaluation and care provided by EMS prior to arrival at the medical facility. A complete PCR should be completed as soon as possible but should not cause a delay in the EMS Back in Service Time.
4. The EMS Unit will be cleaned, disinfected, and restocked (if necessary) during the EMS Back in Service Time interval.
5. Any EMS Back in Service Time delay resulting in a prolonged EMS Back in Service Time will be documented in Patient Care Report (PCR) as an "EMS Turn-Around Delay" as required and defined in the North Carolina College of Emergency Physicians (NCCEP) EMS Dataset Standards Document.
6. All EMS Turn-Around Delays will be reviewed regularly within the EMS System Peer Review Committee.



Standards Policy: Service Metric Policy Section

EMS Wheels Rolling (Turn-Out) Time

Policy:

The EMS Wheels Rolling (Turn-out) Time will be less than 90 seconds, 90% of the time, for all events identified and classified as an emergent or hot (with lights and siren) response.

Definition:

The EMS Wheels Rolling (Turn-out) Time is defined as the time interval beginning with the time the EMS Dispatch Center notifies an EMS Unit to respond to a specific EMS event and ending with the time the EMS Unit is moving en route to the scene of the event.

Purpose:

The purpose of this policy is to:

- Provide a timely and reliable response for all EMS events within the EMS System.
- Provide quality EMS service and patient care to the county's citizens.
- Provide a means for continuous evaluation to assure policy compliance.

Procedure:

The following procedures shall be implemented to assure policy compliance:

1. In EMS Dispatch Centers where Emergency Medical Dispatch (EMD) has been implemented, EMS Units will be dispatched by EMD certified personnel in accordance with the standards developed by the Medical Director and the Emergency Medical Dispatch Protocols.
2. The EMS Unit Wheels Rolling (Turn-out) time will be less than 90 seconds from time of dispatch, 90% of the time. If a unit fails to check en route within 2:59 (mm:ss), the next available EMS unit will be dispatched.
3. Without question, exception, or hesitation, EMS Units will respond as dispatched (hot or cold). This includes both requests to respond on active calls and requests to "move-up" to cover areas of the System that have limited EMS resources available.
4. An EMS Unit may divert from a current cold (no lights and sirens) call to a higher priority hot (with lights and sirens) call **ONLY IF**:
 - The EMS Unit can get to the higher priority call before it can reach the lower priority call. Examples of High Priority Calls: Chest Pain, Respiratory Distress, CVA, etc.
 - The diverting EMS Unit must notify the EMS Dispatch Center that they are diverting to the higher priority call.
 - The diverting EMS Unit ensures that the EMS Dispatch Center dispatches an EMS Unit to their original call.
 - Once a call has been diverted, the next EMS Unit dispatched must respond to the original call. A call cannot be diverted more than one (1) time.
5. Any EMS Wheels Rolling (Turn-out) Time delay resulting in a prolonged EMS Response Time for emergent hot (with lights and sirens) events will be documented in Patient Care Report (PCR) as an "EMS Response Delay" as required and defined in the North Carolina College of Emergency Physicians (NCCEP) EMS Dataset Standards Document.
6. All EMS Response Delays will be reviewed regularly within the EMS System Peer Review Committee.



State Poison Center

Policy:

The state poison center should be utilized by the 911 centers and the responding EMS services to obtain assistance with the prehospital triage and treatment of patients who have a potential or actual poisoning.

Purpose:

The purpose of this policy is to:

- Improve the care of patients with poisonings, envenomations, and environmental/biochemical terrorism exposures in the prehospital setting.
- Provide for the most timely and appropriate level of care to the patient, including the decision to transport or treat on the scene.
- Integrate the State Poison Center into the prehospital response for hazardous materials and biochemical terrorism responses

Procedure:

1. The 911 call center will identify and if EMD capable, complete key questions for the Overdose/Poisoning, Animal Bites/Attacks, or Carbon Monoxide/Inhalation/HazMat emergency medical dispatch complaints and dispatch the appropriate EMS services and/or directly contact the State Poison Center for consultation.
2. If no immediate life threat or need for transport is identified, EMS personnel may conference the patient/caller with the Poison Center Specialist at the **State Poison Center at 800-222-1222**. If possible, dispatch personnel should remain on the line during conference evaluation.
3. The Poison Center Specialist at the State Poison Center will evaluate the exposure and make recommendations regarding the need for on-site treatment and/or hospital transport in a timely manner. If dispatch personnel are not on-line, the Specialist will recontact the 911 center and communicate these recommendations.
4. If the patient is determined to need EMS transport, the poison center Specialist will contact the receiving hospital and provide information regarding the poisoning, including treatment recommendations. EMS may contact medical control for further instructions or to discuss transport options.
5. If the patient is determined not to require EMS transport, personnel will give the phone number of the patient/caller to the Poison Center Specialist. The Specialist will initiate a minimum of one follow-up call to the patient/caller to determine the status of patient.
6. Minimal information that should be obtained from the patient for the state poison center includes:
 - Name and age of patient
 - Time of exposure
 - Signs and symptoms
 - Substance(s) involved
 - Any treatment given
7. Minimal information which should be provided to the state poison center for mass poisonings, including biochemical terrorism and HazMat, includes:
 - Substance(s) involved
 - Signs and symptoms
 - Time of exposure
 - Any treatment given



Air Transport

Policy:

Air transport should be utilized whenever patient care can be improved by decreasing transport time or by giving advanced care not available from ground EMS services, but available from air medical transport services (i.e. blood).

Purpose:

The purpose of this policy is to:

- Improve patient care in the prehospital setting.
- Allow for expedient transport in serious, mass casualty settings.
- Provide life-saving treatment such as blood transfusion.
- Provide more timely access to interventional care in acute Stroke and ST-elevation myocardial infarction (STEMI) patients

Procedure:

Patient transportation via ground ambulance will not be delayed to wait for helicopter transportation.

If the patient is packaged and ready for transport and the helicopter is not on the ground, or within a reasonable distance, the transportation will be initiated by ground ambulance.

Air transport should be considered if any of the following criteria apply:

- High priority patient with > 20 minute transport time
- Entrapped patients with > 10 minute estimated extrication time
- Multiple casualty incident with red/yellow tag patients
- Multi-trauma or medical patient requiring life-saving treatment not available in prehospital environment (i.e., blood transfusion, invasive procedure, operative intervention)
- Time dependent medical conditions such as acute ST-elevation myocardial infarctions (STEMI) or acute Stroke that could benefit from the resources at a specialty center as per the EMS System's Stroke and STEMI Plans.

If a potential need for air transport is anticipated, but not yet confirmed, an air medical transport service can be placed on standby.

If the scene conditions or patient situation improves after activation of the air medical transport service and air transport is determined not to be necessary, paramedic or administrative personnel may cancel the request for air transport.

Minimal Information which should be provided to the air medical transport service include:

- Number of patients
- Age of patients
- Sex of patients
- Mechanism of injury or complaint (MVC, fall, etc)



Standards Policy: Transport Policy Section

Safe Transport of Pediatric Patients

Policy:

Without special considerations children are at risk of injury when transported by EMS. EMS must provide appropriate stabilization and protection to pediatric patients during EMS transport.

Purpose:

To provide:

- Provide a safe method of transporting pediatric patients within an ambulance.
- Protect the EMS system and personnel from potential harm and liability associated with the transportation of pediatric patients.

Procedure:

1. Drive cautiously at safe speeds observing traffic laws.
2. Tightly secure all monitoring devices and other equipment.
3. Insure that all pediatric patient less than 40 lbs are restrained with an approved child restraint device secured appropriately to the stretcher or captains chair.
3. Insure that all EMS personnel use the available restraint systems during the transport.
4. Transport adults and children who are not patients, properly restrained, in an alternate passenger vehicle, whenever possible.
5. Do not allow parents, caregivers, or other passengers to be unrestrained during transport.
6. NEVER attempt to hold or allow the parents or caregivers to hold the patient during transport.



Transport

Policy:

All individuals served by the EMS system will be evaluated, treated, and furnished transportation (if indicated) in the most timely and appropriate manner for each individual situation.

Purpose:

To provide:

- Rapid emergency EMS transport when needed.
- Appropriate medical stabilization and treatment at the scene when necessary
- Protection of patients, EMS personnel, and citizens from undue risk when possible.

Procedure:

1. All trauma patients with significant mechanism or history for multiple system trauma will be transported as soon as possible. The scene time should be 10 minutes or less.
2. All acute Stroke and acute ST-Elevation Myocardial Infarction patients will be transported as soon as possible. The scene time should be 10 minutes or less for acute Stroke patients and 15 minutes or less (with 12 Lead ECG) for STEMI patients
2. Other Medical patients will be transported in the most efficient manner possible considering the medical condition. Advanced life support therapy should be provided at the scene if it would positively impact patient care. Justification for scene times greater than 20 minutes should be documented.
3. No patients will be transported in initial response non-transport vehicles.
4. In unusual circumstances, transport in other vehicles may be appropriate when directed by EMS administration.



Table of Contents

Airway Section:

- AP-2. BIAD King
- AP-5. Cricothyrotomy Surgical
- AP-6. Intubation Oral Tracheal
- AP-8. Video Laryngoscopy Glide Scope
- AP-10. Tracheostomy Tube Change
- AP-11. Endotracheal Tube Introducer
- AP-12. Intubation Confirmation CO2 Detector
- AP-13. Foreign Body Obstruction

Assessment Section:

- ASP-1. Adult
- ASP-2. Pain
- ASP-3. Pediatric
- ASP-4. Blood Glucose Analysis
- ASP-5. Capnography
- ASP-6. Pulse Oximetry
- ASP-7. Reperfusion Checklist
- ASP-8. Stroke Screen LA Prehospital
- ASP-9. Temperature Measurement
- ASP-10. Orthostatic Blood Pressure
- ASP-11. Verbal De-escalation



Table of Contents

Cardiac Section:

- CSP-1. 12-Lead ECG
- CSP-2. Cardioversion
- CSP-3. Cardiac External Pacing
- CSP-4. Cardiopulmonary Resuscitation
- CSP-5. Defibrillation Automated
- CSP-6. Defibrillation Manual
- CSP-7. Defibrillation Dual or Double

Parenteral Access Section:

- PAS-2. Arterial Line Maintenance
- PAS-3. Venous Blood Draw
- PAS-4. Central Line Maintenance
- PAS-5. Epidural Maintenance
- PAS-6. Intraventricular Catheter
- PAS-7. Existing Catheters Maintenance
- PAS-8. External Jugular Access
- PAS-9. Venous Extremity
- PAS-11. Intraosseous
- PAS-12. Swan Ganz Maintenance

Respiratory Section:

- RSP-1. Suctioning Advanced
- RSP-2. Suctioning Basic
- RSP-3. Nebulizer Inhalation Therapy
- RSP-4. NIPPV
- RSP-5. Respirator Operation
- RSP-6. Ventilator Operation



Table of Contents

Universal Section:

- USP-1. Childbirth
- USP-2. Decontamination
- USP-3. Gastric Tube Insertion
- USP-4. Injection SQ/IM
- USP-5. Restraints Physical
- USP-6. Restraints Therapeutic Take Down
- USP-7. Urinary Catheterization

Wound Care/Trauma Section:

- WTP-1. Chest Decompression
- WTP-2. Spinal Motion Restriction
- WTP-3. Splinting
- WTP-4. General
- WTP-5. Hemostatic Agent
- WTP-7. Tourniquet



Airway: BIAD King

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications for Blind Insertion Airway Device (BIAD) Use:

- Inability to adequately ventilate a patient with a Bag Valve Mask or longer EMS transport distances require a more advanced airway.
- Appropriate intubation is impossible due to patient access or difficult airway anatomy.
- Inability to secure an endotracheal tube in a patient who does not have a gag reflex where at least one failed intubation attempt has occurred.
- Patient must be unconscious.

Procedure:

1. Preoxygenate the patient.
2. Select the appropriate tube size for the patient.
3. Lubricate the tube.
4. Grasp the patient's tongue and jaw with your gloved hand and pull forward.
5. Gently insert the tube rotated laterally 45-90 degrees so that the blue orientation line is touching the corner of the mouth. Once the tip is at the base of the tongue, rotate the tube back to midline. Insert the airway until the base of the connector is in line with the teeth and gums.
6. Inflate the pilot balloon with 45-90 ml of air depending on the size of the device used.
7. **Ventilate the patient while gently withdrawing the airway until the patient is easily ventilated.**
8. Auscultate for breath sounds and sounds over the epigastrium and look for the chest to rise and fall.
9. The large pharyngeal balloon secures the device.
10. **Confirm tube placement using end-tidal CO₂ detector.**
11. **EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene**
12. **It is strongly recommended that an Airway Evaluation Form be completed with any BIAD use.**

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Airway: Cricothyrotomy-Surgical

Clinical Indications:

P **PARAMEDIC** **P**

- Failed Airway Protocol
- Management of an airway when standard airway procedures cannot be performed or have failed in a patient \geq 12 years old.

Procedure:

1. Have suction and supplies available and ready.
2. Locate the cricothyroid membrane utilizing anatomical landmarks.
3. Prep the area with an antiseptic swab (Betadine).
4. Attach a 5-cc syringe to an 18G - 1 & 1/2-inch needle.
5. Insert the needle (with syringe attached) perpendicularly through the cricothyroid membrane with the needle directed posteriorly.
6. During needle insertion, gentle aspiration should be applied to the syringe. Rapid aspiration of air into the syringe indicates successful entry into the trachea. Do not advance the needle any further. Attach forceps and remove syringe.
7. With the needle remaining in place, make a 1-inch vertical incision through the skin and subcutaneous tissue above and below the needle using a scalpel. Using blunt dissection technique, expose the cricothyroid membrane. This is a bloody procedure. The needle should act as a guide to the cricothyroid membrane.
8. With the needle still in place, make a horizontal stabbing incision approx. 1/2 inch through the membrane on each side of the needle. Remove the needle.
9. Using (skin hook, tracheal hook, or gloved finger) to maintain surgical opening, insert the cuffed tube into the trachea. (Cric tube from the kit or a #6 endotracheal tube is usually sufficient).
10. Inflate the cuff with 5-10cc of air and ventilate the patient while manually stabilizing the tube.
11. All of the standard assessment techniques for insuring tube placement should be performed (auscultation, chest rise & fall, end-tidal CO₂ detector, etc.) Esophageal bulb devices are not accurate with this procedure.
12. Secure the tube.
13. If Available apply end tidal carbon dioxide monitor (Capnography) and record readings on scene, en route to the hospital, and at the hospital.
14. Document ETT size, time, result (success), and placement location by the centimeter marks either at the patient's teeth or lips on/with the patient care report (PCR). Document all devices used to confirm initial tube placement and after each movement of the patient.
15. Consider placing an NG or OG tube to clear stomach contents after the airway is secured.
- 16. It is strongly recommended that the airway (if equipment is available) be monitored continuously through Capnography and Pulse Oximetry.**
- 17. It is strongly recommended that an Airway Evaluation Form be completed with all intubations**

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Airway: Intubation Oral Tracheal

A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Inability to adequately ventilate a patient with a Bag Valve Mask or longer EMS transport distances require a more advanced airway.
- An unconscious patient without a gag reflex who is apneic or is demonstrating inadequate respiratory effort.
- A component of Drug Assisted Intubation

Procedure:

1. Prepare, position and oxygenate the patient with 100% Oxygen.
2. Select proper ET tube (and stylette, if used), have suction ready.
3. Using laryngoscope, visualize vocal cords. (Use Sellick maneuver/BURP to assist you).
4. Limit each intubation attempt to 30 seconds with BVM between attempts.
5. Visualize tube passing through vocal cords.
- 6. Confirm and document tube placement using an end-tidal CO₂ monitoring or esophageal bulb device.**
7. Inflate the cuff with 3-to10 cc of air; secure the tube to the patient's face.
8. Auscultate for bilaterally equal breath sounds and absence of sounds over the epigastrium. If you are unsure of placement, remove tube and ventilate patient with bag-valve mask.
9. Consider using a Blind Insertion Airway Device if intubation efforts are unsuccessful.
10. If Available apply end tidal carbon dioxide monitor (Capnography) and record readings on scene, en route to the hospital, and at the hospital.
11. Document ETT size, time, result (success), and placement location by the centimeter marks either at the patient's teeth or lips on/with the patient care report (PCR). Document all devices used to confirm initial tube placement. Also document positive or negative breath sounds before and after each movement of the patient.
12. Consider placing an NG or OG tube to clear stomach contents after the airway is secured with an ET tube.
- 13. End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.**
- 4. It is strongly recommended that an Airway Evaluation Form be completed with all intubations**

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Airway: Video Laryngoscopy Glidescope

Clinical Indications:

- Patient requires advanced airway.

A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Preoxygenate the patient and use in conjunction with procedure ASP - 6.
2. Select the appropriate ETT size and GlideRite Rigid Stylette for the patient. Ready suction.
3. Power on GlideScope and allow 30 seconds for anti-fog mechanism to warm.
4. Using GlideScope visualize the vocal cords and facilitate the intubation:

In the mouth: looking directly into the patient's mouth and with the VL blade in left hand, introduce GlideScope VL into the midline of the oral pharynx. Look into the mouth to prevent soft tissue damage.

At the screen: With GlideScope VL inserted, look to monitor to identify the epiglottis, then manipulate the scope to obtain the best glottic view.

In the mouth: Looking directly into the patient's mouth, not at screen, carefully guide the distal tip of the ETT into position near the tip of the GlideScope VL. Insert the ETT behind or adjacent to the VL blade.

At the screen: Look to the monitor to complete tracheal intubation. Gently rotate or angle the ETT to redirect as needed.

Avoid excessive lifting or pushing of the glottis with the VL blade.

Reducing the elevation applied to the VL blade may facilitate intubation.

Advance the ETT while simultaneously withdrawing the stylette with the thumb.

Withdraw the stylette approximately 5 cm (2 inches).

Do not insert the stylette into the larynx during intubation – this will prevent passing into the glottis.

Secure and verify the proper ETT placement.



5. Auscultate for breath sounds and sounds over the epigastrium and look for the chest to rise and fall.
6. Secure the ETT tube with tape or mechanical tube holder.
7. **Confirm tube placement using end-tidal CO₂ detector.**
8. **End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.**
12. **Complete the Airway Evaluation Form.**

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Airway: Tracheostomy Tube Change

A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Presence of Tracheostomy site.
- Urgent or emergent indication to change the tube, such as obstruction that will not clear with suction, dislodgement, or inability to oxygenate/ventilate the patient without other obvious explanation.

Procedure:

1. Have all airway equipment prepared for standard airway management, including equipment of orotracheal intubation and failed airway.
2. Have airway device (endotracheal tube or tracheostomy tube) of the same size as the tracheostomy tube currently in place as well as 0.5 size smaller available (e.g., if the patient has a #6.0 Shilley, then have a 6.0 and a 5.5 tube).
3. Lubricate the replacement tube(s) and check the cuff.
4. Remove the tracheostomy tube from mechanical ventilation devices and use a bag-valve apparatus to pre-oxygenate the patient as much as possible.
5. Once all equipment is in place, remove devices securing the tracheostomy tube, including sutures and/or supporting bandages.
6. If applicable, deflate the cuff on the tube. If unable to aspirate air with a syringe, cut the balloon off to allow the cuff to lose pressure.
7. Remove the tracheostomy tube.
8. Insert the replacement tube. Confirm placement via standard measures except for esophageal detection (which is ineffective for surgical airways).
9. If there is any difficulty placing the tube, re-attempt procedure with the smaller tube.
10. If difficulty is still encountered, use standard airway procedures such as oral bag-valve mask or endotracheal intubation (as per protocol). **More difficulty with tube changing can be anticipated for tracheostomy sites that are immature – i.e., less than two weeks old. Great caution should be exercised in attempts to change immature tracheostomy sites.**
11. Document procedure, confirmation, patient response, and any complications in the PCR

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment for this skill should include direct observation at least once per certification cycle.



Airway: Endotracheal Tube Introducer (Bougie)

Clinical Indications:

- Patients meet clinical indications for oral intubation
- Initial intubation attempt(s) unsuccessful
- Predicted difficult intubation

A	AEMT	A
P	PARAMEDIC	P

Contraindications:

- Three attempts at orotracheal intubation (utilize failed airway protocol)
- Age less than eight (8) or ETT size less than 6.5 mm

Procedure:

1. Prepare, position and oxygenate the patient with 100% oxygen;
2. Select proper ET tube without stylet, test cuff and prepare suction;
3. Lubricate the distal end and cuff of the endotracheal tube (ETT) and the distal 1/2 of the Endotracheal Tube Introducer (Bougie) (note: Failure to lubricate the Bougie and the ETT may result in being unable to pass the ETT);
4. Using laryngoscopic techniques, visualize the vocal cords if possible using Sellick's/BURP as needed;
5. Introduce the Bougie with curved tip anteriorly and visualize the tip passing the vocal cords or above the arytenoids if the cords cannot be visualized;
6. Once inserted, gently advance the Bougie until you meet resistance or "hold-up" (if you do not meet resistance you have a probable esophageal intubation and insertion should be re-attempted or the failed airway protocol implemented as indicated);
7. Withdraw the Bougie ONLY to a depth sufficient to allow loading of the ETT while maintaining proximal control of the Bougie;
8. Gently advance the Bougie and loaded ET tube until you have hold-up again, thereby assuring tracheal placement and minimizing the risk of accidental displacement of the Bougie;
9. While maintaining a firm grasp on the proximal Bougie, introduce the ET tube over the Bougie passing the tube to its appropriate depth;
10. If you are unable to advance the ETT into the trachea and the Bougie and ETT are adequately lubricated, withdraw the ETT slightly and rotate the ETT 90 degrees COUNTER clockwise to turn the bevel of the ETT posteriorly. If this technique fails to facilitate passing of the ETT you may attempt direct laryngoscopy while advancing the ETT (this will require an assistant to maintain the position of the Bougie and, if so desired, advance the ETT);
11. Once the ETT is correctly placed, hold the ET tube securely and remove the Bougie;
12. Confirm tracheal placement according to the intubation protocol, inflate the cuff with 3 to 10 cc of air, auscultate for equal breath sounds and reposition accordingly;
13. When final position is determined secure the ET tube, reassess breath sounds, apply end tidal CO₂ monitor, and record and monitor readings to assure continued tracheal intubation.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Standards Procedure (Skill) Airway Section Airway Intubation Confirmation – End-Tidal CO₂ Detector

Clinical Indications:

- The End-Tidal CO₂ detector shall be used with any Endotracheal Tube or Blind Insertion Airway Device use.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

End-tidal (EtCO₂) monitoring is mandatory following placement of an advanced airway.

Procedure:

- Attach End-Tidal CO₂ detector to the Blind Insertion Airway Device or the Endotracheal Tube.
- Note color change. A color change or CO₂ detection will be documented on each respiratory failure or cardiac arrest patient.
- The CO₂ detector shall remain in place with the airway and monitored throughout the prehospital care and transport unless continuous Capnography is used. Any loss of CO₂ detection or color change is to be documented and monitored as procedures are done to verify or correct the airway problem.
- Tube placement should be verified frequently and always with each patient move or loss of color change in the End-Tidal CO₂ detector.
- Document the procedure and the results on/with the Patient Care Report (PCR) as well as on the Airway Evaluation Form.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Airway: Foreign Body Obstruction

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Sudden onset of respiratory distress often with coughing, wheezing, gagging, or stridor due to a foreign-body obstruction of the upper airway.

Procedure:

1. Assess the degree of foreign body obstruction
 - Do not interfere with a mild obstruction allowing the patient to clear their airway by coughing.
 - In severe foreign-body obstructions, the patient may not be able to make a sound. The victim may clutch his/her neck in the universal choking sign.
2. **For an infant**, deliver 5 back blows (slaps) followed by 5 chest thrusts repeatedly until the object is expelled or the victim becomes unresponsive.
3. **For a child**, perform a subdiaphragmatic abdominal thrust (Heimlich Maneuver) until the object is expelled or the victim becomes unresponsive.
4. **For adults**, a combination of maneuvers may be required.
 - First, subdiaphragmatic abdominal thrusts (Heimlich Maneuver) should be used in rapid sequence until the obstruction is relieved.
 - If abdominal thrusts are ineffective, chest thrusts should be used. Chest thrusts should be used primarily in morbidly obese patients and in the patients who are in the late stages of pregnancy
5. If the victim becomes unresponsive, begin CPR immediately but look in the mouth before administering any ventilations. If a foreign-body is visible, remove it.
6. **Do not perform blind finger sweeps in the mouth and posterior pharynx. This may push the object farther into the airway.**
7. In unresponsive patients, AEMT and Paramedic level professionals should visualize the posterior pharynx with a laryngoscope to potentially identify and remove the foreign-body using Magill forceps.
8. Document the methods used and result of these procedures in the patient care report (PCR).

Certification Requirements:

Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Assessment: Adult

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Any patient requesting a medical evaluation that is too large to be measured with a Length-based **2017** Resuscitation Tape.

Procedure:

- Scene size-up, including universal precautions, scene safety, environmental hazards assessment, need for additional resources, by-stander safety, and patient/caregiver interaction
- Assess need for additional resources.
- Initial assessment includes a general impression as well as the status of a patient's airway, breathing, and circulation.
- Assess mental status (e.g., AVPU) and disability (e.g., GCS).
- Control major hemorrhage and assess overall priority of patient.
- Perform a focused history and physical based on patient's chief complaint.
- Assess need for critical interventions.
- Complete critical interventions and perform a complete secondary exam to include a baseline set of vital signs as directed by protocol.
- Maintain an on-going assessment throughout transport; to include patient response/possible complications of interventions, need for additional interventions, and assessment of evolving patient complaints/conditions.
- Document all findings and information associated with the assessment, performed procedures, and any administration of medications on the PCR.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Assessment / Screening Section Pain Assessment and Documentation

Clinical Indications:

- Any patient with pain.


Definitions:

- Pain is an unpleasant sensory and emotional experience associated with actual or potential tissue damage.
- Pain is subjective (whatever the patient says it is).

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

- Initial and ongoing assessment of pain intensity and character is accomplished through the patient's self report.
- Pain should be assessed and documented in the PCR during initial assessment, before starting pain control treatment, and with each set of vitals.
- Pain should be assessed using the appropriate approved scale.
- Three pain scales are available: the 0 – 10, the Wong - Baker "faces", and the FLACC.
 - 0 – 10 Scale: the most familiar scale used by EMS for rating pain with patients. It is primarily for adults and is based on the patient being able to express their perception of the pain as related to numbers. Avoid coaching the patient; simply ask them to rate their pain on a scale from 0 to 10, where 0 is no pain at all and 10 is the worst pain ever.
 - Wong – Baker "FACES" scale: this scale is primarily for use with pediatrics but may also be used with geriatrics or any patient with a language barrier. The faces correspond to numeric values from 0-10. This scale can be documented with the numeric value.



From Hockenberry MJ, Wilson D, Winkelstein ML: Wong's Essentials of Pediatric Nursing, ed. 7, St. Louis, 2005, p. 1259. Used with permission. Copyright, Mosby.

- FLACC scale: this scale has been validated for measuring pain in children with mild to severe cognitive impairment and in pre-verbal children (including infants).

CATEGORIES	SCORING		
	0	1	2
FACE	No particular expression or smile	Occasional grimace or frown, withdrawn, disinterested.	Frequent to constant quivering chin, clenched jaw.
LEGS	Normal position or relaxed.	Uneasy, restless, tense.	Kicking, or legs drawn up.
ACTIVITY	Lying quietly, normal position moves easily.	Squirming, shifting back and forth, tense.	Arched, rigid or jerking.
CRY	No cry, (awake or asleep)	Moans or whimpers; occasional complaint	Crying steadily, screams or sobs, frequent complaints.
CONSOLABILITY	Content, relaxed.	Reassured by occasional touching hugging or being talked to, distractable.	Difficulty to console or comfort

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Assessment: Pediatric

Clinical Indications:

- Any child that can be measured with a Length-based Resuscitation Tape.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

- Scene size-up, including universal precautions, scene safety, environmental hazards assessment, need for additional resources, by-stander safety, and patient/caregiver interaction
- Assess patient using the pediatric triangle of ABCs:
 - Airway and appearance: speech/cry, muscle tone, inter-activeness, look/gaze, movement of extremities
 - Work of breathing: absent or abnormal airway sounds, use of accessory muscles, nasal flaring, body positioning
 - Circulation to skin: pallor, mottling, cyanosis
- Establish spinal immobilization if suspicion of spinal injury
- Establish responsiveness appropriate for age (AVPU, GCS, etc.)
- Color code using Broselow-Luten tape
- Assess disability (pulse, motor function, sensory function, papillary reaction)
- Perform a focused history and physical exam. Recall that pediatric patients easily experience hypothermia and thus should not be left uncovered any longer than necessary to perform an exam.
- Record vital signs (BP > 3 years of age, cap refill < 3 years of age)
- Include Immunizations, Allergies, Medications, Past Medical History, last meal, and events leading up to injury or illness where appropriate.
- Treat chief complaint as per protocol

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Blood Glucose Analysis

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Patients with suspected hypoglycemia (diabetic emergencies, change in mental status, bizarre behavior, etc.)

Procedure:

1. Gather and prepare equipment.
2. Blood samples for performing glucose analysis can be obtained through a finger-stick or when possible simultaneously with intravenous access.
3. Place correct amount of blood on reagent strip or site on glucometer per the manufacturer's instructions.
4. Time the analysis as instructed by the manufacturer.
5. Document the glucometer reading and treat the patient as indicated by the analysis and protocol.
6. Repeat glucose analysis as indicated for reassessment after treatment and as per protocol.
7. Perform Quality Assurance per manufacture recommendation.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Capnography

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Capnography shall be used when available with the use of all invasive airway procedures including endotracheal, nasotracheal, cricothyrotomy, or Blind Insertion Airway Devices (BIAD).
- Capnography should also be used when possible with CPAP.

Procedure:

1. Attach capnography sensor to the BIAD, endotracheal tube, or oxygen delivery device.
2. Note CO₂ level and waveform changes. These will be documented on each respiratory failure, cardiac arrest, or respiratory distress patient.
3. The capnometer shall remain in place with the airway and be monitored throughout the prehospital care and transport.
4. Any loss of CO₂ detection or waveform indicates an airway problem and should be documented.
5. The capnogram should be monitored as procedures are performed to verify or correct the airway problem.
6. Document the procedure and results on/with the Patient Care Report (PCR) and the Airway Evaluation Form.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Pulse Oximetry

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Patients with suspected hypoxemia.

Procedure:

1. Apply probe to patient's finger or any other digit as recommended by the device manufacturer.
2. Allow machine to register saturation level.
3. Record time and initial saturation percent on room air if possible on/with the patient care report (PCR).
4. Verify pulse rate on machine with actual pulse of the patient.
5. Monitor critical patients continuously until arrival at the hospital. If recording a one-time reading, monitor patients for a few minutes as oxygen saturation can vary.
6. Document percent of oxygen saturation every time vital signs are recorded and in response to therapy to correct hypoxemia.
7. In general, normal saturation is 97-99%. Below 94%, suspect a respiratory compromise.
8. Use the pulse oximetry as an added tool for patient evaluation. Treat the patient, not the data provided by the device.
9. The pulse oximeter reading should never be used to withhold oxygen from a patient in respiratory distress or when it is the standard of care to apply oxygen despite good pulse oximetry readings, such as chest pain. Supplemental oxygen is not required if the oxyhemoglobin saturation is $\geq 94\%$, unless there are obvious signs of heart failure, dyspneic, or hypoxic to maintain to 94%.
10. Factors which may reduce the reliability of the pulse oximetry reading include but are not limited to:
 - Poor peripheral circulation (blood volume, hypotension, hypothermia)
 - Excessive pulse oximeter sensor motion
 - Fingernail polish (may be removed with acetone pad)
 - Carbon monoxide bound to hemoglobin
 - Irregular heart rhythms (atrial fibrillation, SVT, etc.)
 - Jaundice
 - Placement of BP cuff on same extremity as pulse ox probe.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Reperfusion Checklist

Clinical Indications:

Rapid evaluation of a patient with suspected acute stroke and/or acute myocardial infarction (STEMI) to:

- Determine eligibility and potential benefit from fibrinolysis..
- Rapid identification of patients who are not eligible for fibrinolysis and will require interventional therapy.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Follow the appropriate protocol for the patient’s complaint to assess and identify an acute condition which could potentially benefit from fibrinolysis. If a positive finding is noted on one of the following assessments, proceed to step 2.
 - Perform a 12-lead ECG to identify an acute ST elevation myocardial infarction (STEMI).
 - Perform the Los Angeles Pre-hospital Stroke Screen to identify an acute stroke
2. Complete the Reperfusion Check Sheet to identify any potential contraindications to fibrinolysis. (See Appendix)
 - Systolic Blood Pressure greater than 180 mm Hg
 - Diastolic Blood Pressure greater than 110 mm Hg
 - Right vs. Left Arm Systolic Blood Pressure difference of greater than 15 mm Hg
 - History of structural Central Nervous System disease (age >= 18, history of aneurysm or AV-malformation, tumors, masses, hemorrhage, etc.)
 - Significant closed head or facial trauma within the previous 3 months
 - Recent (within 6 weeks) major trauma, surgery (including laser eye surgery), gastrointestinal bleeding, or severe genital-urinary bleeding
 - Bleeding or clotting problem or on blood thinners
 - CPR performed greater than 10 minutes
 - Currently Pregnant
 - Serious Systemic Disease such as advanced/terminal cancer or severe liver or kidney failure.
3. Identify if the patient is currently in heart failure or cardiogenic shock. For these patients, a percutaneous coronary intervention is more effective.
 - Presence of pulmonary edema (rales greater than halfway up lung fields)
 - Systemic hypoperfusion (cool and clammy)
4. If any contraindication is noted using the check list and an acute Stroke is suspected by exam or a STEMI is confirmed by ECG, activate the EMS Stroke Plan or EMS STEMI Plan for fibrinolytic ineligible patients. This may require the EMS Agency, an Air Medical Service, or a Specialty Care Transport Service to transport directly to an specialty center capable of interventional care within the therapeutic window of time.
5. Record all findings in the Patient Care Report (PCR).

Certification Requirements:

Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Stroke Screen: LA Prehospital

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Suspected Stroke Patient

Procedure:

1. Assess and treat suspected stroke patients as per protocol.
2. The Los Angeles Prehospital Stroke Screen (LAPSS) form should be completed for all suspected stroke patients (see appendix). There are six screening criteria items on the LAPSS form.
3. Screen the patient for the following criteria:
 - Age over 45 years
 - No history of a seizure disorder
 - New onset of symptoms in last 24 hours
 - Patient ambulatory prior to event
 - Blood glucose between 60-400
4. The final criterion consists of performing a patient exam looking for facial droop, unilateral grip weakness/absence, or unilateral arm weakness. One of these exam components must be positive to answer “yes” on the screening form.
5. **If all of the LAPSS screening criteria are met (“yes” to all criteria OR if unknown), follow the EMS System Stroke Plan and alert the receiving hospital of a possible stroke patient as early as possible.**
6. All sections of the LAPSS form must be completed.
7. The completed LAPSS form should be attached or documented in the PCR.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Temperature Measurement

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Monitoring body temperature in a patient with suspected infection, hypothermia, hyperthermia, or to assist in evaluating resuscitation efforts.

Procedure:

- For adult patients that are conscious, cooperative, and in no respiratory distress, an oral temperature is preferred (steps 2 to 4 below). For infants or adults that do not meet the criteria above, a rectal temperature is preferred (steps 6 to 8 below).
- To obtain an oral temperature, ensure the patient has no significant oral trauma and place the thermometer under the patient’s tongue with appropriate sterile covering.
- Have the patient seal their mouth closed around thermometer.
- If using an electric thermometer, leave the device in place until there is indication an accurate temperature has been recorded (per the “beep” or other indicator specific to the device). If using a traditional thermometer, leave it in place until there is no change in the reading for at least 30 seconds (usually 2 to 3 minutes). Proceed to step 8.
- Prior to obtaining a rectal temperature, assess whether the patient has suffered any rectal trauma by history and/or brief examination as appropriate for patient’s complaint.
- To obtain a rectal temperature, cover the thermometer with an appropriate sterile cover, apply lubricant, and insert into rectum no more than 1 to 2 cm beyond the external anal sphincter.
- Follow guidelines in step 5 above to obtain temperature.
- Record time, temperature, method (oral, rectal), and scale (C° or F°) in Patient Care Report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Assessment / Screening Section

Orthostatic Blood Pressure Measurement

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Patient situations with suspected blood, fluid loss, or dehydration with no indication for spinal immobilization. Orthostatic vital signs are not routinely recommended.
- Patients ≥ 8 years of age, or patients larger than the Broselow-Luten tape
- Orthostatic Vital Signs are not sensitive nor specific for volume loss / dehydration and may induce syncope in some cases. Assessment of orthostatic vital signs are not routinely recommended. Local Medical Director should indicate and educate on situations where they may be helpful.

Procedure:

1. Gather and prepare standard sphygmomanometer and stethoscope.
2. With the patient supine, obtain pulse and blood pressure.
3. Have the patient sit upright.
4. After 30 seconds, obtain blood pressure and pulse.
5. If the systolic blood pressure falls more than 30 mmHg or the pulse rises more than 20 bpm, the patient is considered to be orthostatic.
6. If a patient experiences dizziness upon sitting or is obviously dehydrated based on history or physical exam, formal orthostatic examination should be omitted and fluid resuscitation initiated.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Verbal De-escalation

Agency Name:		SATISFACTORY <input type="checkbox"/>
Provider Name:	Paramedic	
Instructor Name:	EMT AEMT Paramedic Physician	UNSATISFACTORY <input type="checkbox"/>

Instructor:		EMR	
1. Evaluate providers skill performance using the check off list below.			
2. Circle performance indicator:		B	B
YES = Provider completed skill with no assistance from instructor.		A	A
NO = Provider unable to complete skill satisfactorily following instructor intervention.			
IL = Provider able to complete skill satisfactorily following Instructor Led (teaching) intervention.		P	P
Satisfactory performance indicated with ≥ 8 YES / IL completions. (Combination of both YES and IL)			

YES	NO	IL	Verbalizes indications for Verbal de-escalation techniques: 1. Behavioral Health Crisis 2. Behavior Activity Rating Score ≥ 5
-----	----	----	--

YES	NO	IL	Verbalizes contraindications: None
-----	----	----	------------------------------------

YES	NO	IL	Demonstrates respect of patient's personal space <ul style="list-style-type: none"> • Maintain about 6 feet of distance and do not position yourself between the patient and only exit • Both you and patient should be able to exit the room without feeling blocked-in
-----	----	----	---

YES	NO	IL	Does not provoke patient during interaction <ul style="list-style-type: none"> • <u>Your body language must convey that you want to listen and that you do not want to inflict harm</u> Your hands should be visible and open Do not face the patient head-on. Always stand at an angle Avoid prolonged staring or direct eye contact • Make sure others are not provoking the patient (providers, family members, bystanders, providers, police officers)
-----	----	----	---

YES	NO	IL	Establishes rapport, initiates and maintains verbal contact <ul style="list-style-type: none"> • One person should make and maintain verbal contact, introduce yourself and explain your role Multiple providers talking to the patient will create confusion and may escalate patient's behavior • Emphasize you are there to keep the patient safe • Ask the patient their name and how they want to be addressed
-----	----	----	---

YES	NO	IL	Use concise statements when talking <ul style="list-style-type: none"> • Agitation creates problems in a patient's ability to process information • Keep your conversation simple and short in nature allowing time for patient to process information • <u>Repeat your statements, requests, or commands to ensure understanding</u> This is called a loop, you may need to repeat 2 – 12 times before patient understands
-----	----	----	---

YES	NO	IL	Identify wants, feelings, and stress causing the crisis <ul style="list-style-type: none"> • "When you called 911, how did you think we could help you?" • "I would like to know what caused you to become upset today so we can help you" • Identifying a need can help to quickly de-escalate the situation
-----	----	----	---

YES	NO	IL	Listen closely to patient <ul style="list-style-type: none"> • You should be able to repeat back what is said by the patient "Tell me if I have all this right" "Let me make sure I understand what you said"
-----	----	----	---

YES	NO	IL	Agree or agree to disagree <ul style="list-style-type: none"> • If statements are truthful, then agree with the truth • Agree in principle, maybe patient's statement is not true, but you can agree, that in general, the idea is true • Agree with the odds, anyone may be upset by the same circumstances • Do not agree with delusions, at that point you can agree to disagree
-----	----	----	--

YES	NO	IL	Set clear limits on acceptable behavior <ul style="list-style-type: none"> • Set limits in a positive, matter-of-fact manner, and not in a threatening manner • Inform the patient that harm to self or other providers will not be tolerated • If the patient's behavior is frightening to providers, tell the patient so • Remind the patient you are there to help, keep them safe, but the providers cannot be abused in the process
-----	----	----	---

YES	NO	IL	Offer choices to patient (if available) and remain positive in your interactions <ul style="list-style-type: none"> • Offer choices that are realistic and that may provide comfort such as drinks, food, blankets, etc. • If medication is needed, offer choice between PO and IM/IV, offer medication early in encounter
-----	----	----	---

YES	NO	IL	Debrief provider team following the incident (if restraints necessary, debrief patient as well) What went well? What could have gone better? What did we learn? Who needs to know?
-----	----	----	--

Instructor notes:



Verbal De-escalation

Clinical Information for Verbal De-escalation

Objective of Procedure:

- Verbal engagement with patient and establishing collaborative relationship with patient
- Preventing violent behavior
- Avoiding use of restraints
- Reducing patient anger and frustration
- Maintaining staff and patient safety
- Enabling patients to manage their emotions and regain personal control

Scope of Practice: EMR EMT AEMT Paramedic

Indications:

1. Behavioral Health Crisis
2. Behavior Activity Rating Score ≥ 5

Contraindications:

None

Clinical Presentation:

Patient experiencing a behavioral crisis defined as:

- Significantly deviates from society's expectations and commonly held normal behavior
 - Behavior that is unusual for patient's baseline
 - Bizarre
 - Threatening
 - Dangerous to self and/or others
 - Alarming to patient, family, or bystanders
 - Interferes with the patients ability to perform basic life functions and activities of daily living
- Behavior Activity Rating Score ≥ 5

Potential Complications:

- Injury to patient, provider, or bystander
- Need to move to restraint procedure
- Exacerbation of agitated condition

Procedure references:

1. Palmer J. (2019). Joint Commission Issues De-escalation Guidebook for Healthcare Facilities and Workers. Patient Safety and Quality Healthcare (PSQH). <https://www.psqh.com/analysis/joint-commission-issues-de-escalation-guidebook-for-healthcare-facilities-and-workers/>
2. Richmond JS, Berlin JS, Fishkind AV, et al. (2012). Verbal De-escalation of the Agitated Patient: Consensus Statement of the American Association for Emergency Psychiatry Project BETA De-escalation Workgroup. West J Emerg Med 13(1):17-25. doi: 10.5811/westjem.2011.9.6864



Cardiac: 12 Lead ECG


Clinical Indications:

- Suspected cardiac patient
- Suspected tricyclic overdose
- Electrical injuries
- Syncope

B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Assess patient and monitor cardiac status.
2. Administer oxygen as patient condition warrants.
3. If patient is unstable, definitive treatment is the priority. If patient is stable or stabilized after treatment, perform a 12 Lead ECG.
4. Prepare ECG monitor and connect patient cable with electrodes.
5. Enter the required patient information (patient name, etc.) into the 12 lead ECG device.
6. Expose chest and prep as necessary. Modesty of the patient should be respected.
7. Apply chest leads and extremity leads using the following landmarks:
 - RA -Right arm
 - LA -Left arm
 - RL -Right leg
 - LL -Left leg
 - V1 -4th intercostal space at right sternal border
 - V2 -4th intercostal space at left sternal border
 - V3 -Directly between V2 and V4
 - V4 -5th intercostal space at midclavicular line
 - V5 -Level with V4 at left anterior axillary line
 - V6 -Level with V5 at left midaxillary line
8. Instruct patient to remain still.
9. Press the appropriate button to acquire the 12 Lead ECG.
10. If the monitor detects signal noise (such as patient motion or a disconnected electrode), the 12 Lead acquisition will be interrupted until the noise is removed.
11. Once acquired, transmit the ECG data by fax to the appropriate hospital.
12. Contact the receiving hospital to notify them that a 12 Lead ECG has been sent.
13. Monitor the patient while continuing with the treatment protocol.
14. Download data as per guidelines and attach a copy of the 12 lead to the PCR.
15. Document the procedure, time, and results on/with the patient care report (PCR)



Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Cardiac: Cardioversion

P PARAMEDIC P

Clinical Indications:

- Unstable patient with a tachydysrhythmia (rapid atrial fibrillation, supraventricular tachycardia, ventricular tachycardia)
- Patient is not pulseless (the pulseless patient requires unsynchronized cardioversion, i.e., defibrillation)

Procedure:

1. Ensure the patient is attached properly to a monitor/defibrillator capable of synchronized cardioversion.
2. Have all equipment prepared for unsynchronized cardioversion/defibrillation if the patient fails synchronized cardioversion and the condition worsens.
3. Consider the use of pain or sedating medications.
4. Set energy selection to the appropriate setting.
5. Set monitor/defibrillator to synchronized cardioversion mode.
6. Make certain all personnel are clear of patient.
7. Press and hold the shock button to cardiovert. Stay clear of the patient until you are certain the energy has been delivered. NOTE: It may take the monitor/defibrillator several cardiac cycles to “synchronize”, so there may a delay between activating the cardioversion and the actual delivery of energy.
8. Note patient response and perform immediate unsynchronized cardioversion/defibrillation if the patient’s rhythm has deteriorated into pulseless ventricular tachycardia/ventricular fibrillation, following the procedure for Defibrillation-Manual.
9. If the patient’s condition is unchanged, repeat steps 2 to 8 above, using escalating energy settings.
10. Repeat until maximum setting or until efforts succeed. Consider discussion with medical control if cardioversion is unsuccessful after 2 attempts.
11. Note procedure, response, and time in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle., or other mechanisms as deemed appropriate by the local EMS System.



Cardiac: External Pacing

P **PARAMEDIC** **P**

Clinical Indications:

- Patients with symptomatic bradycardia (less than 60 per minute) with signs and symptoms of inadequate cerebral or cardiac perfusion such as:
 - Chest Pain
 - Hypotension
 - Pulmonary Edema
 - Altered Mental Status, Confusion, etc.
 - Ventricular Ectopy
- Asystole, pacing must be done early to be effective.
- PEA, where the underlying rhythm is bradycardic and reversible causes have been treated.

Procedure:

1. Attach standard four-lead monitor.
2. Apply defibrillation/pacing pads to chest and back:
 - One pad to left mid chest next to sternum
 - One pad to mid left posterior chest next to spine.
3. Rotate selector switch to pacing option.
4. Adjust heart rate to 70 BPM for an adult and 100 BPM for a child.
5. Note pacer spikes on EKG screen.
6. Slowly increase output until capture of electrical rhythm on the monitor.
7. If unable to capture while at maximum current output, stop pacing immediately.
8. If capture observed on monitor, check for corresponding pulse and assess vital signs.
9. Consider the use of sedation or analgesia if patient is uncomfortable.
10. Document the dysrhythmia and the response to external pacing with ECG strips in the PCR.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Standards Procedure (Skill) Cardiac Section

Cardiac: Cardiopulmonary Resuscitation (CPR)

Clinical Indications:

- Basic life support for the patient in cardiac arrest

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Assess the patient's level of responsiveness.
2. If no response, open the patient's airway with the head-tilt, chin-lift and look, listen, and feel for respiratory effort. If the patient may have sustained C-spine trauma, use the modified jaw thrust while maintaining immobilization of the C-spine. For infants, positioning the head in the sniffing position is the most effective method for opening the airway.
3. Check for pulse (carotid for adults and older children, brachial for infants) for at least 10 seconds. If no pulse, begin chest compressions based on chart below:

Age	Location	Depth	Rate
Infant	Over sternum, between nipples (inter-mammary line), 2-3 fingers	At least 1/3 AP diameter of chest About 1.5 inches 4 cm	Continuous compressions at least 100 – 120/minute
Child	Over sternum, just cephalad from xyphoid process, heel of one hand	At least 1/3 AP diameter of chest About 2 inches 5 cm	Continuous compressions at least 100 – 120/minute
Adult	Over sternum, just cephalad from xyphoid process, hands with interlocked fingers	At least 2 inches 5 cm	Continuous compressions at least 100 – 120/minute

4. If patient is an adult, go to step 5. If no respiratory effort in a pediatric patient, give two ventilations. If air moves successfully, go to step 5. If air movement fails, proceed to the Airway Obstruction Procedure.
5. Go to Cardiac Arrest Procedure. Begin ventilations in the adult as directed in the Cardiac Arrest Procedure
6. Provide 1 breath every 6 seconds with the BVM or BIAD. Use EtCO₂ to guide your ventilations as directed in the Cardiac Arrest Protocol.
7. Chest compressions should be provided in an uninterrupted manner. Only brief interruptions (< 5 seconds with a maximum of 10 seconds) are allowed for rhythm analysis, defibrillation, and performance of procedures
8. Document the time and procedure in the Patient Care Report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Cardiac: Defibrillation-Automated

Clinical Indications:

- Patients in cardiac arrest (pulseless, non-breathing).
- Age < 8 years, use Pediatric Pads if available.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Contraindication:

- Pediatric patients who are so small that the pads cannot be placed without touching one another.

Procedure:

1. **If multiple rescuers available, one rescuer should provide uninterrupted chest compressions while the AED is being prepared for use.**
2. Apply defibrillator pads per manufacturer recommendations. Based on 2010 guidelines, place pads preferably in AP or AL position when implanted devices (pacemakers, AICDs) occupy preferred pad positions and attempt to avoid placing directly over device.
3. Remove any medication patches on the chest and wipe off any residue.
4. If necessary, connect defibrillator leads: white to the anterior chest pad and the red to the posterior pad.
5. Activate AED for analysis of rhythm.
6. **Stop CPR and clear the patient** for rhythm analysis. Keep interruption in CPR as brief as possible.
7. Defibrillate if appropriate by depressing the “shock” button. **Assertively state “CLEAR” and visualize that no one, including yourself, is in contact with the patient prior to defibrillation.** The sequence of defibrillation charges is preprogrammed for monophasic defibrillators. Biphasic defibrillators will determine the correct joules accordingly.
8. Begin CPR (chest compressions and ventilations) immediately after the delivery of the defibrillation.
9. After 2 minutes of CPR, analyze rhythm and defibrillate if indicated. Repeat this step every 2 minutes.
10. If “no shock advised” appears, perform CPR for two minutes and then reanalyze.
11. Transport and continue treatment as indicated.
12. **Keep interruption of CPR compressions as brief as possible. Adequate CPR is a key to successful resuscitation.**
13. **If pulse returns please use the Post Resuscitation Protocol**

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Standards Procedure (Skill) Cardiac Section

Cardiac: Defibrillation-Manual

P PARAMEDIC P

Clinical Indications:

- Cardiac arrest with ventricular fibrillation or pulseless ventricular tachycardia

Procedure:

1. **Ensure that Chest Compressions are adequate and interrupted only when absolutely necessary.**
2. Clinically confirm the diagnosis of cardiac arrest and identify the need for defibrillation.
3. After application of an appropriate conductive agent if needed, apply defibrillation hands free pads (recommended to allow more continuous CPR) or paddles to the patient's chest in the proper position
 - Paddles: right of sternum at 2nd ICS and anterior axillary line at 5th ICS
 - Pads: anterior-posterior positionFor patients with implanted pacers/defibrillators, paddles or pads can be in AP or AL positions. The presence of implanted pacers/defibrillators should not delay defibrillation. Attempt to avoid placing paddles or pads directly above device.

4. Set the appropriate energy level
5. Charge the defibrillator to the selected energy level. **Continue chest compressions while the defibrillator is charging.**
6. If using paddles, assure proper contact by applying 25 pounds of pressure on each paddle.
7. **Hold Compressions, assertively state, "CLEAR" and visualize that no one, including yourself, is in contact with the patient.**
8. Deliver the countershock by depressing the discharge button(s) when using paddles, or depress the **shock button** for hands free operation.
9. Immediately resume chest compressions and ventilations for 2 minutes. After 2 minutes of CPR, analyze rhythm and check for pulse only if appropriate for rhythm.
10. Repeat the procedure every two minutes as indicated by patient response and ECG rhythm.
11. Keep interruption of CPR compressions as brief as possible. Adequate CPR is a key to successful resuscitation.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Cardiac: Defibrillation-Dual or Double

Clinical Indications:

P PARAMEDIC **P**


- Cardiac arrest with persistent ventricular fibrillation or pulseless ventricular tachycardia.
- Refractory ventricular fibrillation or pulseless ventricular tachycardia where ≥ 3 shocks delivered.

Procedure:

1. Ensure that Chest Compressions are adequate and interrupted only when absolutely necessary.

2. Clinically confirm the diagnosis of cardiac arrest and identify the need for defibrillation.
3. Prepare sites for second pad set attachment and apply defibrillation hands free pads:

- Pads: First defibrillator pads in anterior-posterior position
- Pads: Second defibrillator pads in anterior-lateral position:
- Ensure pads are not in contact with one another.



For patients with implanted pacers/defibrillators:
Avoid placing paddles or pads directly above device.

4. Set the appropriate energy level and assure controls for both defibrillator / monitors are accessible to provider performing defibrillation.
5. At next pulse / rhythm check, if refractory or persistent VF/VT continues:
Charge the defibrillator to the selected energy level.
Continue chest compressions while the defibrillator is charging.
6. Optional: Agencies may provide a single shock at this point with the second defibrillator / monitor to provide a change in energy vector delivered to the heart then move to step 7 if VF / VT persists.
- 7. When both monitor / defibrillators have reached selected energy setting:**
Hold Compressions, assertively state, "CLEAR" and visualize that no one, including yourself, is in contact with the patient.
2 options at this point:
Option 1 (double simultaneous): Provider depresses both defibrillator shock buttons simultaneously.
Option 2 (dual sequential): Provider depresses monitor 1 shock button and then immediately following, depresses monitor 2 shock button.
8. Immediately resume chest compressions and ventilations for 2 minutes. After 2 minutes of CPR, analyze rhythm and check for pulse only if appropriate for rhythm.
9. Repeat the procedure every two minutes as indicated by patient response and ECG rhythm.
10. Keep interruption of CPR compressions as brief as possible. Adequate CPR is a key to successful resuscitation.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Arterial Line Maintenance

Clinical Indications:

P **PARAMEDIC** **P**

- Transport of a patient with an existing arterial line.

Procedure:

1. Make certain arterial line is secured prior to transport, including intersection of arterial catheter and IV/Monitoring lines.
2. Use available equipment for monitoring of arterial pressures via arterial line.
3. Do not use the arterial line for administration of any fluids or medications.
4. If there is any question regarding dislodgement of the arterial line and bleeding results, remove the line and apply direct pressure over the site for at least five minutes before checking to ensure hemostasis.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Parenteral Access: Venous Blood Draw

Clinical Indications:

- Collection of a patient’s blood for laboratory analysis

A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Utilize universal precautions as per OSHA.
2. Select vein and prep as usual.
3. Select appropriate blood-drawing devices.
4. Draw appropriate tubes of blood for lab testing.
5. Assure that the blood samples are labeled with the correct information (a minimum of the patients name, along with the date and time the sample was collected).
6. Deliver the blood tubes to the appropriate individual at the hospital.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Central Line Maintenance

Clinical Indications:

P **PARAMEDIC** **P**

- Transport of a patient with a central venous pressure line already in place

Procedure:

1. Prior to transportation, ensure the line is secure.
2. Medications and IV fluids may be administered through a central venous pressure line. Such infusions must be held while the central venous pressure is transduced to obtain a central venous pressure, but may be restarted afterwards.
3. Do not manipulate the central venous catheter.
4. If the central venous catheter becomes dysfunctional, does not allow drug administration, or becomes dislodged, contact medical control.
5. Document the time of any pressure measurements, the pressure obtained, and any medication administration in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Epidural Catheter Maintenance

Clinical Indications:

P **PARAMEDIC** **P**

- Presence of an epidural catheter in a patient requiring transport

Procedure:

1. Prior to transport, ensure catheter is secure and that transport personnel are familiar with medication(s) being delivered and devices used to control medication administration.
2. No adjustments in catheter position are to be attempted.
3. No adjustments in medication dosage or administration are to be attempted without direct approval from on-line medical control.
4. Report any complications immediately to on-line medical control.
5. Document the time and dose of any medication administration or rate adjustment in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Ventricular Catheter Maintenance

P PARAMEDIC P

Clinical Indications:

- Transport of a patient with an intra-ventricular catheter in place

Procedure:

1. Prior to transport, ensure the catheter is secure.
2. Prior to transport, determine from the referring hospital/physician the desired patient position (e.g., supine, head of bed elevated 30 degrees, etc.).
3. Prior to transport, determine the height at which the drain is to be maintained, given the patient position desired from #2 above (if applicable).
4. Do not manipulate or move the drain.
5. If the patient or height of the drain is altered, immediately correct based on the pre-determined configuration in step 2 and 3 above.
6. Report any problems immediately to on-line medical control.
7. Document the time and any adjustments or problems in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Existing Catheters

Clinical Indications:

P **PARAMEDIC** **P**

- Inability to obtain adequate peripheral access.
- Access of an existing venous catheter for medication or fluid administration.
- Central venous access in a patient in cardiac arrest.

Procedure:

1. Clean the port of the catheter with alcohol wipe.
2. Using sterile technique, withdraw 5-10 ml of blood and discard syringe in sharps container.
3. Using 5cc of normal saline, access the port with sterile technique and gently attempt to flush the saline.
4. If there is no resistance, no evidence of infiltration (e.g., no subcutaneous collection of fluid), and no pain experienced by the patient, then proceed to step 5. If there is resistance, evidence of infiltration, pain experienced by the patient, or any concern that the catheter may be clotted or dislodged, do not use the catheter.
5. Begin administration of medications or IV fluids slowly and observe for any signs of infiltration. If difficulties are encountered, stop the infusion and reassess.
6. Record procedure, any complications, and fluids/medications administered in the Patient Care Report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: External Jugular Access

A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- External jugular vein cannulation is indicated in a critically ill patient ≥ 8 years of age who requires intravenous access for fluid or medication administration and in whom an extremity vein is not obtainable.
- External jugular cannulation can be attempted initially in life threatening events where no obvious peripheral site is noted.

Procedure:

1. Place the patient in a supine head down position. This helps distend the vein and prevents air embolism.
2. Turn the patient's head toward the opposite side if no risk of cervical injury exists.
3. Prep the site as per peripheral IV site.
4. Align the catheter with the vein and aim toward the same side shoulder.
5. "Tourniqueting" the vein lightly with one finger above the clavicle, puncture the vein midway between the angle of the jaw and the clavicle and cannulate the vein in the usual method.
6. Attach the IV and secure the catheter avoiding circumferential dressing or taping.
7. Document the procedure, time, and result (success) on/with the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Parenteral Access: Venous-Extremity

Clinical Indications:

- Any patient where intravenous access is indicated (significant trauma, emergent or potentially emergent medical condition).

A	AEMT	A
P	PARAMEDIC	P

Procedure:

- Saline locks may be used as an alternative to an IV tubing and IV fluid in every protocol at the discretion of the ALS professional.
- Paramedic/AEMT can use intraosseous access where threat to life exists as provided for in the Venous Access-Intraosseous procedure.
- Use the largest catheter bore necessary based upon the patient's condition and size of veins.
- Fluid and setup choice is preferably:
 - Lactated Ringers with a macro drip (10 gtt/cc) for burns
 - Normal Saline with a macro drip (10 gtt/cc) for medical conditions, trauma or hypotension
 - Normal Saline with a micro drip (60 gtt/cc) for medication infusions
- Inspect the IV solution for expiration date, cloudiness, discoloration, leaks, or the presence of particles.
- Connect IV tubing to the solution in a sterile manner. Fill the drip chamber half full and then flush the tubing bleeding all air bubbles from the line.
- Place a tourniquet around the patient's extremity to restrict venous flow only.
- Select a vein and an appropriate gauge catheter for the vein and the patient's condition.
- Prep the skin with an antiseptic solution.
- Insert the needle with the bevel up into the skin in a steady, deliberate motion until the bloody flashback is visualized in the catheter.
- Advance the catheter into the vein. **Never** reinsert the needle through the catheter. Dispose of the needle into the proper container without recapping.
- Draw blood samples when appropriate.
- Remove the tourniquet and connect the IV tubing or saline lock.
- Open the IV to assure free flow of the fluid and then adjust the flow rate as per protocol or as clinically indicated.

Rates are preferably:

- Adult: KVO: 60 cc/hr (1 gtt/ 6 sec for a macro drip set)
- Pediatric: KVO: 30 cc/hr (1 gtt/ 12 sec for a macro drip set)

If shock is present:

- Adult: 500 cc fluid boluses repeated as long as lungs are dry and BP < 90. Consider a second IV line.
- Pediatric: 20 cc/kg boluses repeated PRN for poor perfusion.

- Cover the site with a sterile dressing and secure the IV and tubing.
- Label the IV with date and time, catheter gauge, and name/ID of the person starting the IV.
- Document the procedure, time and result (success) on/with the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Intraosseous

A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Rapid, regular IV access is unavailable with any of the following:
- Cardiac arrest.
- Multisystem trauma with severe hypovolemia.
- Severe dehydration with vascular collapse and/or loss of consciousness.
- Respiratory failure / Respiratory arrest.
- Burns.

Contraindications:

- Fracture proximal to proposed intraosseous site.
- History of Osteogenesis Imperfecta
- Current or prior infection at proposed intraosseous site.
- Previous intraosseous insertion or joint replacement at the selected site.

Procedure:

1. Don personal protective equipment (gloves, eye protection, etc.).
2. **Proximal tibia:** Identify anterior-medial aspect of the proximal tibia (bony prominence below the knee cap). The insertion location will be 1-2 cm (2 finger widths) below this.
Distal tibia: If this site is not suitable, and patient is an adult, identify the anterior-medial aspect of the distal tibia (2 cm proximal to the medial malleolus).
Distal femur: If this site is not suitable, and patient is a pediatric, identify the patella with the leg outstretched to prevent bending of the knee. The insertion site is approximately 1 cm above the patella and approximately 1 – 2 cm medially.
Proximal humerus: Acceptable insertion site for adult patients. Locate the insertion site 1 – 2 cm above the surgical neck on the most prominent aspect of the greater tubercle. This is located on the lateral aspect of the ball of the humerus. Direct the needle at a 45 degree angle or toward the opposite hip.
3. Prep the site recommended by the device manufacturer with providone-iodine ointment or solution.
4. For manual pediatric devices, hold the intraosseous needle at a 60 to 90 degree angle, aimed away from the nearby joint and epiphyseal plate, twist the needle handle with a rotating grinding motion applying controlled downward force until a “pop” or “give” is felt indicating loss of resistance. Do not advance the needle any further.
5. For the EZ-IO intraosseous device, hold the intraosseous needle at a 60 to 90 degree angle, aimed away from the nearby joint and epiphyseal plate, power the driver until a “pop” or “give” is felt indicating loss of resistance. Do not advance the needle any further. Utilize the yellow needle for the proximal humerus. The pink needle is only intended for use in neonatal patients.
6. For the Bone Injection Gun (BIG), find and mark the manufacturers recommended site. Position the device and pull out the safety latch. Trigger the BIG at 90° to the surface and remove the injection device.
7. Remove the stylette and place in an approved sharps container.
8. Attach a syringe filled with at least 5 cc NS; aspirate bone marrow for manual devices only, to verify placement; then inject at least 5 cc of NS to clear the lumen of the needle.
9. Attach the IV line and adjust flow rate. A pressure bag may assist with achieving desired flows.
10. Stabilize and secure the needle with dressings and tape.
11. Paramedic may administer 10 to 20 mg (1 to 2 cc) of 2% Lidocaine in adult patients who experience infusion-related pain. This may be repeated prn to a maximum of 60 mg (6 cc).
12. Following the administration of any IO medications, flush the IO line with 10 cc of IV fluid.
13. Document the procedure, time, and result (success) on/with the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation at least once per certification cycle.



Standards Procedure (Skill) Parenteral Access Section

Parenteral Access: Swan-Ganz Catheter Maintenance

P PARAMEDIC P

Clinical Indications:

- Transport of a patient with a Swan-Ganz catheter that is in place prior to transport.

Procedure:

1. Make certain catheter is secure prior to transport.
2. Under the supervision of the nurse or physician caring for the patient, make certain the transport personnel are aware of the depth at which the catheter is secured.
3. **UNDER NO CIRCUMSTANCES SHOULD TRANSPORT PERSONNEL ADVANCE THE SWAN-GANZ CATHETER.**
4. The sterile plastic sheath that surrounds the catheter should not be manipulated.
5. The ports of the catheter may be used to continue administration of medications or IV fluids that were initiated prior to transport. These should be used as any other IV port with attention to sterile technique.
6. If applicable, measurements from the catheter may be obtained during transport and used to guide care as per local protocols and medical control orders.
7. If at anytime during the transport difficulties with the function of the Swan-Ganz catheter is noted, contact medical control.
8. Document the time and any adjustments or problems associated with the catheter in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Respiratory Section

Airway: Suctioning-Advanced

A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Obstruction of the airway (secondary to secretions, blood, or any other substance) in a patient currently being assisted by an airway adjunct such as a naso-tracheal tube, endotracheal tube, Combitube, tracheostomy tube, or a cricothyrotomy tube.

Procedure:

1. Ensure suction device is in proper working order.
2. Preoxygenate the patient as is possible.
3. Attach suction catheter to suction device, keeping sterile plastic covering over catheter.
4. Using the suprasternal notch and the end of the airway into the catheter will be placed as guides, measure the depth desired for the catheter (judgment must be used regarding the depth of suctioning with cricothyrotomy and tracheostomy tubes).
5. If applicable, remove ventilation devices from the airway.
6. With the thumb port of the catheter uncovered, insert the catheter through the airway device.
7. Once the desired depth (measured in #4 above) has been reached, occlude the thumb port and remove the suction catheter slowly.
8. A small amount of Normal Saline (10 ml) may be used if needed to loosen secretions for suctioning.
9. Reattach ventilation device (e.g., bag-valve mask) and ventilate the patient
10. Document time and result in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Respiratory Section

Respiratory: Suctioning-Basic

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Obstruction of the airway (secondary to secretions, blood, or any other substance) in a patient who cannot maintain or keep the airway clear.

Procedure:

1. Ensure suction device is in proper working order with suction tip in place.
2. Preoxygenate the patient as is possible.
3. Explain the procedure to the patient if they are coherent.
4. Examine the oropharynx and remove any potential foreign bodies or material which may occlude the airway if dislodged by the suction device.
5. If applicable, remove ventilation devices from the airway.
6. Use the suction device to remove any secretions, blood, or other substance.
7. The alert patient may assist with this procedure.
8. Reattach ventilation device (e.g., bag-valve mask) and ventilate or assist the patient
9. Record the time and result of the suctioning in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Respiratory Section

Respiratory: Nebulizer Inhalation Therapy

Clinical Indications:

- Patients experiencing bronchospasm.

B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Gather the necessary equipment.
2. Assemble the nebulizer kit.
3. Instill the premixed drug (such as Albuterol or other approved drug) into the reservoir well of the nebulizer.
4. Connect the nebulizer device to oxygen at 4 - 6 liters per minute or adequate flow to produce a steady, visible mist.
5. Instruct the patient to inhale normally through the mouthpiece of the nebulizer. The patient needs to have a good lip seal around the mouthpiece.
6. The treatment should last until the solution is depleted. Tapping the reservoir well near the end of the treatment will assist in utilizing all of the solution.
7. Monitor the patient for medication effects. This should include the patient's assessment of his/her response to the treatment and reassessment of vital signs, ECG, and breath sounds.
8. Assess and document peak flows before and after nebulizer treatments.
9. Document the treatment, dose, and route on/with the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Respiratory Section

Respiratory: NIPPV

(Non-Invasive Positive Pressure)

Clinical Indications:

- Non-Invasive Positive Airway Pressure (NIPPV) is indicated in all patients whom inadequate ventilation is suspected.

This could be as a result of Pulmonary Edema, CHF, COPD, Pneumonia, or Asthma.

- Agencies may utilize Continuous and/or Bi-Level Positive Airway Pressure Devices

B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Contraindications:

- Decreased Mental Status.
- Facial features or deformities that prevent an adequate mask seal.
- Excessive respiratory secretions.

Procedure:

1. Ensure adequate oxygen supply to ventilation device.
2. Explain the procedure to the patient.
3. Consider placement of a nasopharyngeal airway.
4. Place the delivery mask over the mouth and nose. Oxygen should be flowing through the device at this point.
5. Secure the mask with provided straps starting with the lower straps until minimal air leak occurs.
6. If the Positive Pressure is adjustable on the NIPPV device adjust and slowly titrate to achieve a positive pressure as follows:

Continuous pressure device:

5 – 25 cmH₂O for Pulmonary Edema, CHF, COPD, Asthma, Drowning, possible aspiration, or pneumonia.

Bi-Level pressure device:

IPAP 10 – 15 over EPAP 5 – 7 cmH₂O for Pulmonary Edema, CHF, COPD, Asthma, Drowning, possible aspiration, or pneumonia.

During titration keep IPAP – EPAP at least a difference of 5 cmH₂O

25 cmH₂O is maximum pressure that should be utilized with NIPPV.

Increasing positive pressure can cause hypotension.

Use caution or remove and re-evaluate with Systolic Blood Pressures consistently < 100 mmHg.

7. Evaluate the response of the patient assessing breath sounds, oxygen saturation, and general appearance.
8. Titrate oxygen levels to the patient's response. Many patients respond to low FIO₂ (30-50%).
9. Encourage the patient to allow forced ventilation to occur. Observe closely for signs of complications. The patient must be breathing for use of the NIPPV device.
10. Document time and response on patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Respiratory: Respirator Operation

Clinical Indications:

- Transport of an intubated patient

A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Confirm the placement of tube as per airway protocol.
2. Ensure adequate oxygen delivery to the respirator device.
3. Preoxygenate the patient as much as possible with bag-valve mask.
4. Remove BVM and attach tube to respiration device.
5. Per instructions of device, set initial respiration values. For example, set an inspiratory:expiratory ratio of 1:4 (for every 1 second of inspiration, allow 4 seconds and expiration) with a rate of 12 to 20.
6. Assess breath sounds. Allow for adequate expiratory time. Adjust respirator setting as clinically indicated.
7. **It is required that patients on a transport ventilator should be monitored continuously through Capnography and Pulse Oximetry. The ventilatory rate should adjusted to maintain a pulse oximetry of >90 (preferably $\geq 94\%$) while maintaining a pCO₂ of 30-35.**
8. If any worsening of patient condition, decrease in oxygen saturation, or any question regarding the function of the respirator, remove the respirator and resume bag-valve mask ventilations.
9. Document time, complications, and patient response on the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Respiratory: Ventilator Operation

P PARAMEDIC P

Clinical Indications:

- Management of the ventilation of a patient during a prolonged or interfacility transport of an intubated patient.

Procedure:

1. Transporting personnel should review the operation of the ventilator with the treating personnel (physician, nurse, or respiratory therapy) in the referring facility prior to transport if possible.
2. All ventilator settings, including respiratory rate, FiO₂, mode of ventilation, and tidal volumes should be recorded prior to initiating transport. Additionally, the recent trends in oxygen saturation experienced by the patient should be noted.
3. Prior to transport, specific orders regarding any anticipated changes to ventilator settings as well as causes for significant alarm should be reviewed with the referring medical personnel as well as medical control.
4. Once in the transporting unit, confirm adequate oxygen delivery to the ventilator.
5. Frequently assess breath sounds to assess for possible tube dislodgment during transfer.
6. Frequently assess the patient's respiratory status, noting any decreases in oxygen saturation or changes in tidal volumes, peak pressures, etc.
7. Note any changes in ventilator settings or patient condition in the PCR.
8. Consider placing an NG or OG tube to clear stomach contents.
- 9. End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube.**
10. If any significant change in patient condition, including vital signs or oxygen saturation or there is a concern regarding ventilator performance/alerts, remove the ventilator from the endotracheal tube and use a bag-valve mask with 100% O₂. Contact medical control immediately.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Childbirth

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Imminent delivery with crowning

Procedure:

1. Delivery should be controlled so as to allow a slow controlled delivery of the infant. This will prevent injury to the mother and infant.
2. Support the infant's head as needed.
3. Check the umbilical cord surrounding the neck. If it is present, slip it over the head. If unable to free the cord from the neck, double clamp the cord and cut between the clamps.
4. Suction the airway with a bulb syringe.
5. Grasping the head with hands over the ears, gently pull down to allow delivery of the anterior shoulder.
6. Gently pull up on the head to allow delivery of the posterior shoulder.
7. Slowly deliver the remainder of the infant.
8. Clamp the cord 2 inches from the abdomen with 2 clamps and cut the cord between the clamps.
9. Record APGAR scores at 1 and 5 minutes.
10. Follow the **Newly Born Protocol** for further treatment.
11. The placenta will deliver spontaneously, usually within 5 minutes of the infant. Do not force the placenta to deliver.
12. Massaging the uterus may facilitate delivery of the placenta and decrease bleeding by facilitating uterine contractions.
13. Continue transport to the hospital.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Universal Section

Decontamination

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Any patient who may have been exposed to significant hazardous materials, including chemical, biological, or radiological weapons.

Procedure:

- In coordination with HazMAT and other Emergency Management personnel, establish hot, warm and cold zones of operation.
- Ensure that personnel assigned to operate within each zone have proper personal protective equipment.
- In coordination with other public safety personnel, assure each patient from the hot zone undergoes appropriate initial decontamination. This is specific to each incident; such decontamination may include:
 - Removal of patients from Hot Zone
 - Simple removal of clothing
 - Irrigation of eyes
 - Passage through high-volume water bath (e.g., between two fire apparatus) for patients contaminated with liquids or certain solids. Patients exposed to gases, vapors, and powders often will not require this step as it may unnecessarily delay treatment and/or increase dermal absorption of the agent(s).
- Initial triage of patients should occur after step #3. Immediate life threats should be addressed prior to technical decontamination.
- Assist patients with technical decontamination (unless contraindicated based on #3 above). This may include removal of all clothing and gentle cleansing with soap and water. All body areas should be thoroughly cleansed, although overly harsh scrubbing which could break the skin should be avoided.
- Place triage identification on each patient. Match triage information with each patient's personal belongings which were removed during technical decontamination. Preserve these personnel effects for law enforcement.
- Monitor all patients for environmental illness.
- Transport patients per local protocol.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Gastric Tube Insertion

Clinical Indications:

P **PARAMEDIC** **P**

- Gastric decompression in intubated patients or for administration of activated charcoal in patients with altered mental status.

Procedure:

1. Estimate insertion length by superimposing the tube over the body from the nose to the stomach.
2. Flex the neck **if not contraindicated** to facilitate esophageal passage.
3. Liberally lubricate the distal end of the tube and pass through the patient's nostril along the floor of the nasal passage. Do not orient the tip upward into the turbinates. This increases the difficulty of the insertion and may cause bleeding.
4. In the setting of an intubated patient or a patient with facial trauma, oral insertion of the tube may be considered or preferred after securing airway.
5. Continue to advance the tube gently until the appropriate distance is reached.
6. Confirm placement by injecting 20cc of air and auscultate for the swish or bubbling of the air over the stomach. Additionally, aspirate gastric contents to confirm proper placement.
7. Secure the tube.
8. Decompress the stomach of air and food either by connecting the tube to suction or manually aspirating with the large catheter tip syringe.
9. Document the procedure, time, and result (success) on/with the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Universal Section

Injections: Subcutaneous and Intramuscular

B	<i>EMT*</i>	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- When medication administration is necessary and the medication must be given via the SQ or IM route (not auto-injector), or as an alternative route in selected medications.

Procedure:

- Receive and confirm medication order or perform according to standing orders.
- Prepare equipment and medication expelling air from the syringe.
- Explain the procedure to the patient and reconfirm patient allergies.
- The most common site for subcutaneous injection is the arm.
 - Injection volume should not exceed 1 cc.
- The possible injection sites for intramuscular injections include the arm, buttock and thigh.
 - Injection volume should not exceed 1 cc for the arm
 - Injection volume should not exceed 2 cc in the thigh or buttock.
- The thigh should be used for injections in pediatric patients and injection volume should not exceed 1 cc.
- Expose the selected area and cleanse the injection site with alcohol.
- Insert the needle into the skin with a smooth, steady motion
 - SQ: 45-degree angle**
skin pinched
 - IM: 90-degree angle**
skin flattened
- Aspirate for blood
- Inject the medication.
- Withdraw the needle quickly and dispose of properly without recapping.
- Apply pressure to the site.
- Monitor the patient for the desired therapeutic effects as well as any possible side effects.
- Document the medication, dose, route, and time on/with the patient care report (PCR).

** EMT may administer Epinephrine for anaphylaxis, by IM route, if approved by the system medical director.*

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Standards Procedure (Skill) Universal Section

Restraints: Physical

Agency Name: _____

Provider Name: _____ Paramedic

Instructor Name: _____ EMT AEMT Paramedic Physician

SATISFACTORY

UNSATISFACTORY

Instructor:
 1. Evaluate providers skill performance using the check off list below.
 2. Circle performance indicator.
 YES = Provider completed skill with no assistance from instructor.
 NO = Provider unable to complete skill satisfactorily following instructor intervention.
 IL = Provider able to complete skill satisfactorily following Instructor Led (teaching) intervention.
Satisfactory performance indicated with ≥ 12 YES / IL completions. (Combination of both YES and IL)

	EMR		
B	EMT	B	
A	AEMT	A	
P	Paramedic	P	

YES NO IL	<u>Verbalizes indications for physical restraints:</u> 1. Used to ensure the physical safety of the patient, provider, or others 2. Clear and immediate danger to the patient (self), provider, or others 3. When less restrictive alternatives are unsuccessful (e.g.. verbal de-escalation) 4. Delay in restraint will subject patient (self) , providers, or others to risk of serious harm	
YES NO IL	<u>Verbalizes contraindications for physical restraints:</u> 1. Patient has medical decision-making capacity and refuses care 2. Patient is not a danger to self, provider, or others 3. Less restrictive alternatives have not been considered or used	
YES NO IL	<u>Verbalizes assessment of resource needs:</u> Request Law Enforcement if indicated Contact Medical Control if indicated Call for additional providers if indicated <u>Withdraw from scene if unsafe</u>	
YES NO IL	<u>Assemble appropriate equipment and personnel:</u> 1. 3 – 6 providers preferably 2. Don appropriate PPE 3. Soft nylon or leather restraints specifically manufactured for use as restraints	
YES NO IL	<u>Remove potential items from all providers that can be used as weapons:</u> 1. Stethoscope, shears or scissors, hemostats, writing pens, badges, pins 2. Window punch, pocket knives, communication devices	
YES NO IL	<u>Team leader assign roles to providers and discusses plans and strategies:</u> Team leader explains procedure to patient: If patient standing and will not follow directions use Procedure USP – 6. <u>If patient already on cot or flat surface:</u> <ul style="list-style-type: none"> 1 Provider to control the head and airway 1 Provider for each extremity <u>Team leader attempts verbal instructions to move patient to cot if possible:</u> <ul style="list-style-type: none"> 2 Providers take control of both wrists and elbows 2 Providers take control of both ankles and knees 1 Provider controls head/airway and 1 Provider is available for medications May place in lateral decubitus position – DO NOT place prone: 	1
YES NO IL	Soft nylon or leather manufacture restraints are applied to wrist and ankles Secure restraints to cot with quick-release tie Examine patient for potential injuries following restrain application	
YES NO IL	Both lower extremities restrained extended, cross restraints beneath lower extremities One upper extremity restrained extended by patient's side One upper extremity restrained flexed over patient's head Do not tie restraint to cot undercarriage	2
YES NO IL	Assess pulse, motor, and sensory immediately following application Perform pulse, motor, and sensory assessments every 15 minutes afterwards	
YES NO IL	Patient must remain under constant observation by EMS at all times Appropriate monitoring equipment required based on clinical circumstances	
YES NO IL	<u>Patient care report documentation requirements (restraint checklist recommended):</u> Indication for restraint use Type of restrain applied and time of application Pulse, motor, and sensory exams and time of exam	



Instructor notes:



Restraints: Physical

Clinical Information for physical restraints

Objective of Procedure:

To protect a patient from self-harm and/or protection of providers or others on scene
Used when less restrictive alternatives have failed
Used as last resort

Scope of Practice: EMR, EMT, AEMT, and Paramedic

Indications:

Physically combative patient not responding to less restrictive means of de-escalation
Immediate danger of self-harm or harm to providers, or others on scene

Contraindications:

Less restrictive techniques have not been used or considered prior to physical restraint
Intact medical decision-making capacity refusing treatment and not a danger to self or others

Clinical Presentation:

Behavioral health crisis
Altered Mental Status with combativeness
Agitation and violence

Potential Complications:

Positional asphyxiation
Injury to patient, providers, or others
Increased mental stress to patient
Injury following escape from restraints
Bodily fluid exposure

Positioning Considerations:

Do not place patient in a supine position or place objects on top of patient
One arm should be restrained above the head
Both legs should be restrained fully extended
May place in a lateral decubitus position, supine is preferred
Head of bed should be elevated to about 30°

Procedure references:

1. Kowalski JM. (2019). Physical and Chemical Restraint. Roberts and Hedges' Clinical Procedures in Emergency Medicine and Acute Care. 7th ed.(pp 1481 - 1498). Philadelphia, PA. Elsevier.
2. Heiner JD, Moore GP. (2018). The combative and difficult. Rosen's Emergency Medicine: Concepts and Clinical Practice. 9th ed. (pp 2375 - 2386). Philadelphia, PA. Elsevier.
3. Booth JS. (2018, Dec 19). Four-Point Restraint. Retrieved from <https://emedicine.medscape.com/article/1941454-overview>.
4. Bradley S. (2017). Psychiatric Emergencies. AAOS Emergency Care and Transportation of the Sick and Injured. 11th ed. (pp.802 – 827). Burlington, MA. Jones and Bartlett Learning.



Standards Procedure (Skill) Universal Section

Restraints: Therapeutic Take Down



Clinical Information for physical restraints

Objective of Procedure:

To protect a patient from self-harm and/or protection of providers or others on scene
Used when less restrictive alternatives have failed
Used as last resort

Scope of Practice: EMR, EMT, AEMT, and Paramedic

Indications:

Physically combative patient not responding to less restrictive means of de-escalation
Immediate danger of self-harm or harm to providers, or others on scene

Contraindications:

Less restrictive techniques have not been used or considered prior to physical restraint
Intact medical decision-making capacity refusing treatment and not a danger to self or others

Clinical Presentation:

Behavioral health crisis
Altered Mental Status with combativeness
Agitation and violence

Potential Complications:

Positional asphyxiation
Injury to patient, providers, or others
Increased mental stress to patient
Injury following escape from restraints
Bodily fluid exposure

Positioning Considerations:

Do not place patient in a supine position or place objects on top of patient
One arm should be restrained above the head
May place in a lateral decubitus position, supine is preferred
Head of bed should be elevated to about 30°

Procedure references:

1. Kowalski JM. (2019). Physical and Chemical Restraint. Roberts and Hedges' Clinical Procedures in Emergency Medicine and Acute Care. 7th ed.(pp 1481 - 1498). Philadelphia, PA. Elsevier.
2. Heiner JD, Moore GP. (2018). The combative and difficult. Rosen's Emergency Medicine: Concepts and Clinical Practice. 9th ed. (pp 2375 - 2386). Philadelphia, PA. Elsevier.
3. Booth JS. (2018, Dec 19). Four-Point Restraint. Retrieved from <https://emedicine.medscape.com/article/1941454-overview>.
4. Bradley S. (2017). Psychiatric Emergencies. AAOS Emergency Care and Transportation of the Sick and Injured. 11th ed. (pp.802 – 827). Burlington, MA. Jones and Bartlett Learning.



Standards Procedure (Skill) Universal Section

Urinary Catheterization

Agency Name:		SATISFACTORY <input type="checkbox"/>
Provider Name:	Paramedic	
Instructor Name:	EMT AEMT Paramedic Physician	UNSATISFACTORY <input type="checkbox"/>

Instructor:

1. Evaluate providers skill performance using the check off list below.



2. Circle performance indicator.

P
Paramedic
P

YES = Provider completed skill with no assistance from instructor.
 NO = Provider unable to complete skill satisfactorily following instructor intervention.
 IL = Provider able to complete skill satisfactorily following Instructor Led (teaching) intervention.

Satisfactory performance indicated with \geq 12 YES / IL completions. (Combination of both YES and IL)

YES	NO	IL	<p>Verbalizes indications for urinary catheterization:</p> <ol style="list-style-type: none"> 1. Monitoring patient's fluid state and/or response to therapy during transport 2. Collection of urine sample for laboratory analysis 3. Maintaining and monitoring chronic indwelling urinary catheter
YES	NO	IL	<p>Verbalizes contraindications for urinary catheterization:</p> <ol style="list-style-type: none"> 1. Suspected or known trauma to the abdomen, pelvis, or genitalia 2. Grossly bloody urine 3. Pediatric patient
YES	NO	IL	<p>Prepare patient and explain the procedure:</p> <ol style="list-style-type: none"> 1. Explain procedure to patient and the need for sterility 2. Ensure privacy and good lighting 3. Use a chaperone for assistance and observation regardless of age or gender
YES	NO	IL	<p>Assemble appropriate equipment:</p> <ol style="list-style-type: none"> 1. Use or pre-packaged commercial kit is recommended containing: Sterile gloves, sterile lubricant, sterile water Forceps, cotton balls, and cleansing pads Urinary catheter and collection bag Male catheter size: 14 – 16 Fr. Female catheter size: 12 – 14 Fr.
YES	NO	IL	<p>Insertion of urinary catheter:</p> <ol style="list-style-type: none"> 1. Wash hands before and after procedure with soap/water or alcohol-based product 2. Open urinary catheter kit maintaining sterility of contents 3. Don sterile gloves and arrange kit contents for efficient use 4. Attach sterile water syringe to catheter balloon, inflate with 10 mL, and deflate while checking for leaks 5. Attach urinary catheter to collection bag and clamp port 6. Apply drapes to patient in sterile fashion 7. Using your non-dominant hand expose the urinary meatus, hand remains in place until catheter is placed with urine flow and is now unsterile Female: spread the labia apart Male: Retract foreskin exposing entire glans and hold penis straight and perpendicular to the trunk 8. Using your dominant hand, clean the meatus with Betadine, circling out from the male meatus and top to bottom in female 10. Lubricate the catheter tip with sterile lubricant 11. Gently place the catheter tip into the urinary meatus and advance slowly until you have flow of urine, then advance an additional 2 inches Stop immediately if you feel resistance and cannot advance the catheter 12. Inflate the catheter balloon with 10 mL of sterile water and place gently traction on the catheter by withdrawing until you feel resistance. 13. Place the collection bag in a dependent condition relative to the genitalia and secure the catheter to the abdomen or thigh with device supplied
YES	NO	IL	<p>Patient care report documentation requirements:</p> <ol style="list-style-type: none"> 1. Time of insertion, volume of urine obtained on placement, and total volume at end of call 2. Any complications noted



Instructor notes:



Urinary Catheterization

Clinical Information for urinary catheterization

Objective of Procedure:

To introduce or maintain a urinary catheter when urine function has been compromised by illness or surgery.
To monitor fluid output.

Scope of Practice: Paramedic

Indications:

Monitoring patient's fluid state and/or response to therapy during transport
Collection of urine sample for laboratory analysis
Maintaining and monitoring chronic indwelling urinary catheter

Contraindications:

Suspected or known trauma to the abdomen, pelvis, or genitalia
Grossly bloody urine
Pediatric patient

Clinical Presentation:

Medical illness requiring assessment of urine output due to ongoing therapy
Chronic indwelling catheter in urethral, condom, or suprapubic location

Potential Complications:

Trauma to urethra and/or bladder
Urinary infection and/or sepsis
Bodily fluid exposure

Positioning Considerations:

Prone to head of bed elevated to about 30°
Male: Lower extremities can be extended
Female: Lower extremities with knees bent and elevated and abducted

Procedure references:

1. Ortega R, Ng L, Sekhar Pavan, et al. (2008, April 3). Female Urethral Catheterization. Retrieved from <https://www.nejm.org/doi/full/10.1056/nejmvcm0706671>
2. Thomsen TW, Setnik GS. (2006, May 25). Male Urethral Catheterization. Retrieved from <https://www.nejm.org/doi/full/10.1056/nejmvcm054648>.



Chest Decompression

P **PARAMEDIC** **P****Clinical Indications:**

- Patients with hypotension (SBP <90), clinical signs of shock, and at least one of the following signs:
 - Jugular vein distention.
 - Tracheal deviation away from the side of the injury (often a late sign).
 - Absent or decreased breath sounds on the affected side.
 - Hyper-resonance to percussion on the affected side.
 - Increased resistance when ventilating a patient.
- Patients in traumatic arrest with chest or abdominal trauma for whom resuscitation is indicated. These patients may require bilateral chest decompression even in the absence of the signs above.

Procedure:

1. Don personal protective equipment (gloves, eye protection, etc.).
2. Administer high flow oxygen.
3. Identify and prep the site:
 - Locate the second intercostals space in the mid-clavicular line on the same side as the pneumothorax.
 - If unable to place anteriorly, lateral placement may be used at the fourth ICS mid-axillary line.
 - Prepare the site with providone-iodine ointment or solution.
4. Insert the catheter (14 gauge for adults) into the skin over the third rib and direct it just over the top of the rib (superior border) into the interspace.
5. Advance the catheter through the parietal pleura until a “pop” is felt and air or blood exits under pressure through the catheter, then advance catheter only to chest wall.
6. Remove the needle, leaving the plastic catheter in place.
7. Secure the catheter hub to the chest wall with dressings and tape.
8. Consider placing a finger cut from an exam glove over the catheter hub. Cut a small hole in the end of the finger to make a flutter valve. Secure the glove finger with tape or a rubber band. (Note – don’t waste much time preparing the flutter valve; if necessary control the air flow through the catheter hub with your gloved thumb.)

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System. Assessment should include direct observation once per certification cycle.



Spinal Motion Restriction

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Need for Spinal Motion Restriction as determined by protocol.
- **Guidelines for appropriate use of long spine board (LSB) OR any equivalent device below:**

1. **Spine boards or similar rigid devices, should NOT be used during transport or during inter-facility transfers.** They should be utilized for extrication and / or patient transfers, as well as support for chest compressions. They DO NOT improve outcomes and can induce pain, agitation / anxiety, respiratory compromise, and decreased tissue perfusion at pressure points.
2. Devices such as the long or short spine board, scoop stretcher, soft-body splints, etc., should be considered extrication devices rather than transport-devices. Instead, use of Spinal Motion Restriction which includes a rigid cervical collar, manual in-line spine stabilization, maintaining spinal alignment with movement and transfers, and securing to the ambulance stretcher.
3. Penetrating trauma to head, torso, or back with no evidence of spinal injury does not require Spinal Motion Restriction.

Procedure:

1. Gather LSB, scoop, ambulance cot, or other Spinal Motion Restriction device, securing devices, and appropriate C-collar.
2. Explain the procedure to the patient and assess / record neurological exam and pulse status.
3. Place the patient in an appropriately sized C-collar while maintaining in-line stabilization of the C-spine by second provider. In-line stabilization should not involve traction / tension, but rather maintain the head in a neutral, midline position while the first rescuer applies the collar.
4. Once the collar is secure, the second rescuer should still maintain their position to ensure stabilization (the collar is helpful but will not do the job by itself.)
5. If indicated, place patient on a Spinal Motion Restriction device with log-roll or similar technique dependent on circumstances, if patient is supine or prone. During extrication or where otherwise unable to be placed prone or supine, place on Spinal Motion Restriction device by the safest method available that allows maintenance of in-line spinal stability.
6. Stabilize the patient with straps / head rolls / tape / other devices as needed. Once the head is secured to the Spinal Motion Restriction device / stretcher, the second rescuer may release manual in-line stabilization. **Once the patient arrives at the stretcher, REMOVE the rigid Spinal Motion Restriction device while maintaining spinal alignment using log-roll or multi-rescuer lift techniques and transfer and secure to the stretcher for transport.**
7. NOTE: Spinal precautions may be achieved by many methods. Never force a patient into a certain position to immobilize them. Such situations may require a second rescuer to maintain manual stabilization throughout the transport to the hospital. Special equipment such as football players in full pads and helmet may remain immobilized with helmet and pads in place.
8. Document the time of the procedure in the patient care report (PCR).

Certification Requirements:

Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Splinting

Clinical Indications:

- Immobilization of an extremity for transport, either due to suspected fracture, sprain, or injury.
- Immobilization of an extremity for transport to secure medically necessary devices such as intravenous catheters

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Assess and document pulses, sensation, and motor function prior to placement of the splint. If no pulses are present and a fracture is suspected, consider reduction of the fracture prior to placement of the splint.
2. Remove all clothing from the extremity.
3. Select a site to secure the splint both proximal and distal to the area of suspected injury, or the area where the medical device will be placed.
4. Do not secure the splint directly over the injury or device.
5. Place the splint and secure with Velcro, straps, or bandage material (e.g., kling, kerlex, cloth bandage, etc.) depending on the splint manufacturer and design.
6. Document pulses, sensation, and motor function after placement of the splint. If there has been a deterioration in any of these 3 parameters, remove the splint and reassess
7. If a femur fracture is suspected and there is no evidence of pelvic fracture or instability, the following procedure may be followed for placement of a femoral traction splint:
 - Assess neurovascular function as in #1 above.
 - Place the ankle device over the ankle.
 - Place the proximal end of the traction splint on the posterior side of the affected extremity, being careful to avoid placing too much pressure on genitalia or open wounds. Make certain the splint extends proximal to the suspected fracture. If the splint will not extend in such a manner, reassess possible involvement of the pelvis
 - Extend the distal end of the splint at least 6 inches beyond the foot.
 - Attach the ankle device to the traction crank.
 - Twist until moderate resistance is met.
 - Reassess alignment, pulses, sensation, and motor function. If there has been deterioration in any of these 3 parameters, release traction and reassess.
8. Document the time, type of splint, and the pre and post assessment of pulse, sensation, and motor function in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Wound Care-General

Clinical Indications:

- Protection and care for open wounds prior to and during transport.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Procedure:

1. Use personal protective equipment, including gloves, gown, and mask as indicated.
2. If active bleeding, elevate the affected area if possible and hold direct pressure. Do not rely on “compression” bandage to control bleeding. Direct pressure is much more effective.
3. Once bleeding is controlled, irrigate contaminated wounds with saline as appropriate (this may have to be avoided if bleeding was difficult to control). Consider analgesia per protocol prior to irrigation.
4. Cover wounds with sterile gauze/dressings. Check distal pulses, sensation, and motor function to ensure the bandage is not too tight.
5. Monitor wounds and/or dressings throughout transport for bleeding.
6. Document the wound and assessment and care in the patient care report (PCR).

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.



Wound Care-Hemostatic Agent

Clinical Indications:

- Serious hemorrhage that can not be controlled by other means.

Contraindications:

- Wounds involving open thoracic or abdominal cavities.

Procedure:

1. Apply approved non-heat-generating hemostatic agent per manufacturer’s instructions.
2. Supplement with direct pressure and standard hemorrhage control techniques.
3. Apply dressing.

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P



Wound Care-Tourniquet

	EMR	
B	EMT	B
A	AEMT	A
P	PARAMEDIC	P

Clinical Indications:

- Life threatening extremity hemorrhage that can not be controlled by other means.
- Serious or life threatening extremity hemorrhage and tactical considerations prevent the use of standard hemorrhage control techniques.

Contraindications:

- Non-extremity hemorrhage
- Proximal extremity location where tourniquet application is not practical

Procedure:

1. Place tourniquet proximal to wound
2. Tighten per manufacturer instructions until hemorrhage stops and/or distal pulses in affected extremity disappear.
3. Secure tourniquet per manufacturer instructions
4. Note time of tourniquet application and communicate this to receiving care providers
5. Dress wounds per standard wound care protocol
6. If delayed or prolonged transport and tourniquet application time > 45 minutes: consider reattempting standard hemorrhage control techniques and removing tourniquet

Certification Requirements:

- Maintain knowledge of the indications, contraindications, technique, and possible complications of the procedure. Assessment of this knowledge may be accomplished via quality assurance mechanisms, classroom demonstrations, skills stations, or other mechanisms as deemed appropriate by the local EMS System.

North Carolina Medical Board Approved Medications for Credentialed EMS Personnel

EMS personnel at any level who administer medications must do so with medical oversight. Personnel must complete appropriate medical education. All EMS System and SCTP protocols, policies and procedures must be reviewed and approved by the Medical Director of the Office of EMS

All items highlighted in “red” are required by NCCEP in all systems with EMS personnel credentialed at the specified level. Specialty Care (SCTP) required items are not listed here, as they can be found on the Specialized Ambulance Protocol Summary (SAPS) form.

Medications	EMR	EMT	AEMT	MEDIC
ACE inhibitors				X
Acetaminophen	X	X	X ¹⁵	X
Adenosine				X
Aminophylline				X
Amiodarone				X
Anti-arrhythmic				X ¹²
Antibiotics				X
Anti-emetic preparations				X
Antivirals				X
Aspirin	X	X	X	X
Atropine	X ⁴	X ⁴	X ⁴	X
Barbiturates				X
Benzodiazepine preparations				X ¹⁴
Beta agonist preparations		X ²	X	X
Beta blockers				X ¹³
Bretylum				X
CI Esterase-Inhibitors				X
Calcium channel blockers				X ¹³
Calcium chloride/gluconate				X
Charcoal		X	X	X
Clonidine				X
Clopidogrel				X
CroFab (Crotalidae Polyvalent Immune Fab)				X ⁸
Crystalloid solutions			X	X
Cyanide poisoning antidote kit				X
Digoxin				X
Diphenhydramine	X ³	X ³	X	X
Diuretics				X
Dobutamine				X
Dopamine				X
Droperidol				X
Epinephrine	X ¹	X ¹	X	X
Etomidate				X
Flumazenil				X
Glucagon			X	X
Glucose, oral	X	X	X	X
Glucose solutions			X	X
Haloperidol				X

Medications	EMR	EMT	AEMT	MEDIC
Heparin (unfractionated and low molecular weight)				X
Histamine 2 blockers			X	X
Hydroxocobalamin				X
Immunizations			X ⁶	X ⁶
Insulin				X
Ipratropium			X	X
Isoproterenol				X
Ketamine				X ⁷
Levetiracetam				X
Lidocaine				X
Magnesium sulfate				X
Mannitol				X
Methylene blue				X
Milrinone				X
N-acetylcysteine				X
Narcotic analgesics				X
Narcotic antagonists	X ^{9,10}	X ^{9,10}	X	X
Nasal spray decongestant		X	X	X
Nesiritide				X
Nitroglycerin		X ²	X	X
Nitroprusside sodium				X
Nitrous oxide				X
Non-prescription medications		X	X	X
Non-steroidal anti-inflammatory		X	X ¹⁵	X
Norepinephrine				X
Octreotide				X
Oxygen	X ⁵	X ⁵	X ⁵	X ⁵
Oxytocin				X
Paralytic agents				X ¹⁷
Phenothiazine preparations				X
Phenylephrine				X
Phenytoin preparations				X
Plasma protein fraction				X
Platelet g-II/IIIa inhibitors				X
Potassium chloride				X
Pralidoxime	X ⁴	X ⁴	X ⁴	X
Procainamide				X
Procaine				X
Proparacaine				X
Propofol				X ⁸
Proton pump inhibitors				X
Sodium bicarbonate				X
Steroid preparations				X
Thiamine			X	X
Thrombolytic agents				X
Topical hemostatic agents	X	X	X	X
Total Parenteral Nutrition				X
Tranexamic Acid (TXA)				X ¹¹
Tuberculosis skin test			X ⁶	X ⁶
Valproic acid				X
Vasopressin			X	X

Medications	EMR	EMT	AEMT	MEDIC
Vasopressor				X ¹⁶
Whole blood and components				X
Ziprasidone				X

¹ EMR and EMT use of epinephrine is limited to the treatment of anaphylaxis and may be administered only by auto injector, unless approved by EMS System Medical Director and OEMS.

² EMT use of beta-agonists and nitroglycerine is limited to patients who currently are prescribed the medication unless approved by the EMS System Medical Director and OEMS as part of the expanded scope. EMTs may administer these medications from EMS supplies.

³ EMR/EMT administration of diphenhydramine is limited to the oral route.

⁴ As a component of preparedness for domestic terrorism, EMS personnel, public safety officers, and other first responders recognized by the EMS system, may carry, self-administer, or administer to a patient atropine and/or pralidoxime, based on written protocols and medical direction. All personnel except for Paramedics must administer these medications by an auto injector.

⁵ Administration of oxygen does not require medical direction.

⁶ Administration of immunizations and TB skin tests are not limited to public health initiatives.

⁷ Ketamine use is restricted to programs that have been approved by the OEMS State Medical Director. It can be used as an induction or post intubation sedation agent in approved DAI programs. Use outside of DAI programs must meet all the requirements outlined in Medical Policy 2 'Ketamine Program Requirements'.

⁸ Propofol use is restricted to programs that have been approved by the OEMS State Medical Director. EMS Systems and SCTP's must submit a policy and education plan to the OEMS prior to approval. EMS personnel cannot initiate Propofol, it can only be used for interfacility transport where infusion has already been started at transferring facility. **EMS units cannot stock Propofol or CroFab. This medication must be provided by the transferring hospital.**

⁹ FR, EMR, and EMT administration of Naloxone is limited to the intra-nasal (IN), intra-muscular (IM), and auto-injector routes.

¹⁰ First Responders (FR) who administer Naloxone must do so under the medical oversight of the County EMS Medical Director, following protocols and procedures approved by the OEMS State Medical Director. FR administration must be monitored by the EMS Systems peer review program.

¹¹ For an EMS System to use Tranexamic Acid (TXA), they must submit for approval by the OEMS State Medical Director a signed letter from any Trauma Centers that would be the recipient of the patient that the destination Trauma Center agrees with its use and will give the 2nd required dose of Tranexamic Acid (TXA).

¹² All Paramedic systems must carry some form of anti-arrhythmic agent. This must either be amiodarone, lidocaine, **or** procainamide.

¹³ Paramedic systems must carry either a calcium channel blocker **or** beta-blocker.

¹⁴ All Paramedic systems must carry some form of injectable benzodiazepine.

¹⁵ AEMT systems must carry either acetaminophen **or** a non-steroidal anti-inflammatory.

¹⁶ All Paramedic systems must carry an approved vasopressor. This must either be dobutamine, dopamine, epinephrine, norepinephrine, phenylephrine, **or** vasopressin.

¹⁷ Paralytic agent use is restricted to Drug Assisted Intubation (DAI) programs approved by the OEMS State Medical Director. They require the submission of; signed NCCEP DAI policy by local medical director, unaltered NCCEP DAI protocols, training documentation, and process for peer review of cases. All DAI must have an EMS Airway Evaluation form completed and signed by local medical director in accordance with the NCCEP DAI policy. Systems utilizing must submit monthly airway forms and cases to the OEMS for review.

North Carolina Medical Board Approved Skills for Credentialed EMS Personnel

All items highlighted in “red” are required by NCCEP in all systems with EMS personnel credentialed at the specified level. Specialty Care (SCTP) required items are not listed here, as they can be found on the Specialized Ambulance Protocol Summary (SAPS) form.

Skills	EMR	EMT	AEMT	MEDIC
12-Lead ECG Acquisition & Transmission		X	X	X
12-Lead ECG Interpretation				X
15-Lead ECG Acquisition				X
Airway Adjuncts (NPA/OPA)	X	X	X	X
Arterial Access - Blood Draw				X
Arterial Line maintenance				X
Blind Insertion Airway Device (BIAD)	X ¹	X ¹	X	X
Capnography (Waveform)	X ⁶	X ⁶	X ⁶	X ⁶
Carbon Monoxide Measurement (non-invasive)	X	X	X	X
Cardiac Monitoring		X ⁴	X ⁴	X
Cardiac Pacing				X
Cardiopulmonary Resuscitation	X	X	X	X
Cardioversion				X
Carotid Massage				X
Central Venous Pressure Line Maintenance				X
Chest Compression-External Device	X	X	X	X
Chest Decompression-Needle				X
Chest Tube Maintenance				X
Childbirth	X	X	X	X
Cricothyrotomy-Needle				X
Cricothyrotomy-Surgical				X ⁵
Decontamination	X	X	X	X
Defibrillation-Automated	X	X	X	X
Defibrillation-Manual				X
Direct Laryngoscopy			X	X
Drug Assisted Intubation (DAI)				X ^{5,6}
Endotracheal Tube Introducer			X	X
Epidural Catheter Maintenance				X
Foreign Body Airway Obstruction	X	X	X	X
Gastric Intubation		X ³	X ³	X
Glucose Measurement	X	X	X	X
Hemostatic Agent	X	X	X	X
Injections – Subcutaneous and Intramuscular		X ²	X	X
Intra-Ventricular Catheter Maintenance				X
Intubation - Nasotracheal			X	X
Intubation - Orotracheal			X ⁶	X ^{6,7}
Intubation Confirmation - Capnometry (color)			X	X
Medication Administration	X ²	X ²	X ²	X ²
Nebulizer Inhalation Therapy		X	X	X
Non-Invasive Positive Pressure Ventilation	X ⁹	X	X	X
Orthostatic Blood Pressure	X	X	X	X
Oxygen Administration	X	X	X	X

Skills	EMR	EMT	AEMT	MEDIC
Patient Assessment	X	X	X	X
Pulse Oximetry	X	X	X	X
Reperfusion Checklist	X	X	X	X
Respirator Operation		X	X	X
Restraints		X	X	X
Specimen Collection		X	X	X
Spinal Motion Restriction	X	X	X	X
Splinting	X	X	X	X
Stroke Screen	X	X	X	X
Suction-Basic	X	X	X	X
Suction-Advanced			X ¹⁰	X ¹⁰
Swan-Ganz Catheter maintenance				X
Taser Probe Removal	X	X	X	X
Temperature Measurement	X	X	X	X
Tourniquet Application	X	X	X	X
Tracheostomy Tube Change			X	X
Urinary Catheterization				X
Venous Access-Blood Draw			X	X
Venous Access-Existing catheters				X
Venous Access-Femoral Line				X
Venous Access-Intraosseous			X	X
Venous Access-Peripheral			X	X
Ventilator Operation		X ⁸	X ⁸	X
Wound Care	X	X	X	X

¹ EMRs and EMTs using blind insertion airway devices must be functioning in EMS systems with medical direction and written treatment protocols.

² EMS personnel educated in approved programs, credentialed by the OEMS, and functioning under physician medical oversight may perform acts and administer intravenous fluids and medications as allowed by the North Carolina Medical Board pursuant to G.S. 143-514. The administration of oxygen does not require medical direction.

³ Gastric tube insertion may be performed only when utilized in conjunction with a blind insertion airway device.

⁴ EMT and AEMT may use the cardiac monitor for vital sign monitoring and EKG transmission.

⁵ Systems performing drug assisted intubation (DAI) must have the ability to perform surgical cricothyrotomy. Commercial cricothyrotomy or tracheostomy kits that create an airway comparable to a surgical cricothyrotomy are acceptable.

⁶ End-tidal (EtCO₂) monitoring is mandatory following placement of an endotracheal tube. EtCO₂ monitoring is mandatory following placement of a BIAD once available on scene.

⁷ Pediatric intubation is an optional skill/procedure.

⁸ Ventilator patients may be transported by EMT/AEMT when all of the following conditions are met:

-Patient is receiving home (or skilled nursing) ventilator therapy.

-The ventilator is portable and can continue to ventilate the patient during transport.

-The patient is accompanied by a non-EMS adult (from either the home or facility) who is knowledgeable, capable, and willing to maintain the ventilator during the EMS transport.

-While in transit, the patient is monitored using pulse oximetry.

⁹ Bag Valve Mask ONLY

¹⁰ For a patient currently being assisted by an airway adjunct such as a naso-tracheal tube, endotracheal tube, BIAD, tracheostomy tube or a cricothyrotomy tube.

-EMD personnel are responsible for:

- 1) Pre-arrival instructions to callers
- 2) Determining and dispatching appropriate EMS resources
- 3) All EMD skills must be performed in EMS systems with medical oversight and written EMS protocols

Appendix Table of Contents

1. Appendix A: Disposition Instruction
2. Appendix B: On Scene Physician Form
3. Appendix C: APGAR Score
4. Appendix D: LA Prehospital Stroke Screen
5. Appendix E: Pain Scale
6. Appendix F: Restraint Checklist
7. Appendix G: Approved Medical Abbreviations
8. Appendix H: Reperfusion Checklist
9. Appendix I: Difficult Airway Evaluation
10. Appendix J: Burns Resources
11. The NC Eye Bank EMS Referral Policy
12. North Carolina EMS Airway Evaluation Form
13. Example: MOST Form
14. Example: DNR Form
15. Recommended EMS Guidelines for Children and Youth with Special Health Care Needs (CYSHCN)



Disposition Instruction Form

Instructions

The EMS Patient Disposition Information (PDI) form has been designed to be used by EMS personnel to legally document a variety of situations. This duplicate form consists of a single page. The front of the page is used to describe the situation and the back lists a variety of specific patient instructions by complaint.

The form should be used to document any refusal of care by a patient (complete refusal or refusal of specific aspects of care) and to document the patient / guardian's understanding of medical instructions.

To understand the intent of this form, it is probably simplest to walk through several common patient encounter situations.

1. Complete refusal of EMS care or transport: The first box "Patient Refusal" should be marked. In the first section, the appropriate blocks for "paramedic recommendation" should also be marked. This section should be explained to the patient or guardian, who should understand that their refusal may result in complications up to and including death. The patient or guardian should be asked to sign the form, indicating that he/she understands the seriousness of the situation and the information provided. If the situation warrants, the paramedic should explain the risks of the refusal using the patient instructions section and the back of the form for assistance. If the instructions section is used, the appropriate blocks should also be checked.
2. Refusal of a specific procedure (IV therapy, for example): The first box "Patient Refusal" should be marked. In the first section, the specific refused procedure should be marked. The first section should be explained to the patient or guardian, who should understand the potential consequences of their refusal. The patient or guardian should be asked to sign the form, indicating that he/she understands the seriousness of the situation.
3. The box "Patient Instructions" and the appropriate blocks in that section should be marked. This section and the specific instructions (on the back) should all be carefully explained to the patient and/or guardian, who must understand them. The patient or guardian should be asked to sign the form, indicating that he/she understands the instructions and the seriousness of the situation.

In all situations, the top part of the form should be completed, and as much of the signature portion as necessary. It is preferable to have witnesses, particularly if the patient or guardian refuses to sign. The original form should be kept on file, while a duplicate copy should be provided for the patient or guardian.

Discharge Instructions

UNIVERSAL INSTRUCTIONS:

- YOU HAVE NOT RECEIVED A COMPLETE MEDICAL EVALUATION. SEE A PHYSICIAN AS SOON AS POSSIBLE.
- IF AT ANY TIME AFTER YOU HAVE TAKEN ANY MEDICATION, YOU HAVE TROUBLE BREATHING, START WHEEZING, GET HIVES OR A RASH, OR HAVE ANY UNEXPECTED REACTION, CALL 911 IMMEDIATELY.
- IF YOUR SYMPTOMS WORSEN AT ANY TIME, YOU SHOULD SEE YOUR DOCTOR, GO TO THE EMERGENCY DEPARTMENT OR CALL 911.

ABDOMINAL PAIN:

- Abdominal pain is also called belly pain. Many illnesses can cause abdominal pain and it is very difficult for EMS to identify the cause.
- Take your temperature every 4 hours.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- Your pain gets worse or is now only in 1 area
- You vomit (throw up) blood or find blood in your bowel movement
- You become dizzy or faint
- Your abdomen becomes distended or swollen
- You have a temperature over 100° F
- You have trouble passing urine
- You have trouble breathing

BACK PAIN:

- Apply heat to the painful area to help relieve pain. You may use a warm heating pad, whirlpool bath, or warm, moist towels for 10 to 20 minutes every hour.
- Stay in bed as much as possible the first 24 hours.
- Begin normal activities when you can do them without causing pain.
- When picking things up, bend at the hips and knees. Never bend from the waist only.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- You have shooting pains into your buttocks, groin, legs, or arms or the pain increases.
- You have trouble urinating or lose control of your stools or urine.
- You have numbness or weakness in your legs, feet, arms, or hands.

FEVER:

- Always take medications as directed. Tylenol and Ibuprofen can be taken at the same time.
- If you are taking antibiotics, take them until they are gone, not until you are feeling better.
- Drink extra liquids (1 glass of water, soft drink or gatorade per hour of fever for an adult)
- If the temperature is above 103° F, it can be brought down by a sponge bath with room temperature water. Do not use cold water, a fan, or an alcohol bath.
- Temperature should be taken every 4 hours.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- Temperature is greater than 101° F for 24 hours
- A child becomes less active or alert.
- The Temperature does not come down with Acetaminophen (Tylenol) or Ibuprofen with the appropriate dose.

HEAD INJURY:

- Immediately after a blow to the head, nausea, and vomiting may occur.
- Individuals who have sustained a head injury must be checked, and if necessary awakened, every 2 hours for the first 24 hours.
- Ice may be placed on the injured area to decrease pain and swelling.
- Only drink clear liquids such as juices, soft drinks, or water the first 12 hours after injury..
- Acetaminophen (Tylenol) or Ibuprofen only may be used for pain.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- The injured person has persistent vomiting, is not able to be awakened, has trouble walking or using an arm or leg, has a seizure, develops unequal pupils, has a clear or bloody fluid coming from the ears or nose, or has strange behavior.

INSECT BITE/STING:

- A bite or sting typically is a red lump which may have a hole in the center. You may have pain, swelling and a rash. Severe stings may cause a headache and an upset stomach (vomiting).
- Some individuals will have an allergic reaction to a bite or sting. Difficulty breathing or chest pain is an emergency requiring medical care.
- Elevation of the injured area and ice (applied to the area 10 to 20 minutes each hour) will decrease pain and swelling.
- Diphenhydramine (Benadryl) may be used as directed to control itching and hives.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- You develop any chest pain or difficulty breathing.
- The area becomes red, warm, tender, and swollen beyond the area of the bite or sting.
- You develop a temperature above 101° F.

RESPIRATORY DISTRESS:

- Respiratory Distress is also known as shortness of breath or difficulty breathing.
- Causes of Respiratory Distress include reactions to pollen, dust, animals, molds, foods, drugs, infections, smoke, and respiratory conditions such as Asthma and COPD. If possible avoid any causes which produce respiratory distress.
- If you have seen a physician for this problem, take all medication's as directed.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- Temperature is greater than 101° F.
- The cough, wheezing, or breathing difficulty becomes worse or does not improve even when taking medications.
- You have Chest Pain.
- Sputum (spit) changes from clear to yellow, green, grey, or becomes bloody.
- You are not able to perform normal activities.

EXTREMITY INJURY:

- Extremity Injuries may consist of cuts, scrapes, bruises, sprains, or broken bones (fractures).
- Apply ice on the injury for 15 to 20 minutes each hour for the first 1 to 2 days.
- Elevate the extremity above the heart as possible for the first 48 hours to decrease pain and swelling.
- Use the extremity as pain allows.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- Temperature is greater than 101° F.
- The bruising, swelling, or pain gets worse despite the treatment listed above.
- Any problems listed on the **Wound Care instructions** are noted.
- You are unable to move the extremity or if numbness or tingling is noted.
- You are not improved in 24 to 48 hours or you are not normal in 7 to 10 days.

VOMITING/DIARRHEA:

- Vomiting (throwing up) can be caused by many things. It is common in children, but should be watched closely.
- Diarrhea is most often caused by either a food reaction or infection.
- Dehydration is the most serious problem associated with vomiting or diarrhea.
- Drink clear liquids such as water, apple juice, soft drinks, or gatorade for the first 12 hours or until things improve. Adults should drink 8 to 12 glasses of fluids per day with diarrhea. Children should drink 1 cup of fluid for each loose bowel movement.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- Temperature is greater than 101° F.
- Vomiting or Diarrhea lasts longer than 24 hours, gets worse, or blood is noted.
- You cannot keep fluids down or no urination is noted in 8 hours.

WOUND CARE:

- Wounds include cuts, scrapes, bites, abrasions, or puncture wounds.
- If the wound begins to bleed, apply pressure over the wound with a clean bandage and elevate the wound above the heart for 5 to 10 minutes.
- Unless instructed otherwise, clean the wound twice daily with soapy water, and keep the wound dry. It is safe to take a shower but do not place the wound in bath or dish water.
- See a physician for a tetanus shot if it has been 10 years or more since your last one.

Call or see a physician, go to the emergency department, or call 911 immediately if:

- See the **Extremity Injury instructions**.
- Temperature is greater than 101° F.
- Bruising, swelling, or pain gets worse or bleeding is not controlled as directed above.
- Any signs of infection, such as redness, drainage of yellow fluid or pus, red streaks extending from the wound, or a bad smell is noted.



On-Scene Physician Form

This EMS service would like to thank you for your effort and assistance. Please be advised that the EMS Professionals are operating under strict protocols and guidelines established by their medical director and the State of North Carolina. As a licensed physician, you may assume medical care of the patient. In order to do so, you will need to:

1. Receive approval to assume the patient’s medical care from the EMS Agencies Online Medical Control physician.
2. Show proper identification including current North Carolina Medical Board Registration/ Licensure.
3. Accompany the patient to the hospital.
4. Carry out any interventions that do not conform to the EMS Agencies Protocols. EMS personnel cannot perform any interventions or administer medications that are not included in their protocols.
5. Sign all orders on the EMS Patient Care Report.
6. Assume all medico-legal responsibility for all patient care activities until the patient’s care is transferred to another physician at the destination hospital.
7. Complete the “Assumption of Medical Care” section of this form below.

Assumption of Medical Care

I, _____, MD; License #: _____,
(Please Print your Name Here)

have assumed authority and responsibility for the medical care and patient management for

(Insert Patient’s Name Here)

I understand that I must accompany the patient to the Emergency Department. I further understand that all EMS personnel must follow North Carolina EMS Rules and Regulations as well as local EMS System protocols.

_____, MD Date: ____/____/____ Time: ____AM/PM
(Physician Signature Here)

_____, EMS _____ Witness
(EMS Lead Crew Member Signature Here) (Witness Signature Here)



Apgar Score

The Apgar score should be obtained and recorded initially and at 5 minutes with the birth of delivery of any infant.

- Each of the 5 parameters should be scored and then totaled.
- The Minimum score is 0
- The Maximum score is 10

Sign	0	1	2
Heart Rate	Absent	<100 min.	>100 min.
Respiratory Effort	Absent	Weak Cry	Strong Cry
Muscle Tone	Limp	Some Flexion	Good Flexion
Reflex Irritability (when feet stimulated)	No Response	Some Motion	Cry
Color	Blue; Pale	Body Pink Extremities Blue	Pink



Los Angeles Prehospital Stroke Screen (LAPSS)

1. Patient Name: _____
(last name) (first name)

2. Information/History from: Patient Family Member Other

(name - if other than patient) (phone)

3. Last known time patient was at baseline or deficit free and awake:

(military time) (date)

SCREENING CRITERIA

	Yes	Unknown	No
4. Age > 45	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
5. History of seizures or epilepsy absent	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
6. Symptom duration less than 24 hours	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
7. At baseline, patient is not wheelchair bound or bedridden	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
8. Blood glucose between 60 and 400	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
9. Exam: LOOK FOR OBVIOUS ASYMMETRY			
	Normal	Right	Left
Facial smile/grimace	<input type="checkbox"/>	<input type="checkbox"/> Droop	<input type="checkbox"/> Droop
Hand grip	<input type="checkbox"/>	<input type="checkbox"/> Weak	<input type="checkbox"/> Weak
		<input type="checkbox"/> No grip	<input type="checkbox"/> No grip
Arm strength	<input type="checkbox"/>	<input type="checkbox"/> Drifts dn	<input type="checkbox"/> Drifts dn
		<input type="checkbox"/> Falls fast	<input type="checkbox"/> Falls fast

Based on exam, patient has only unilateral (not bilateral) weakness: YES NO

10. Items 4, 5, 6, 7, 8, 9 all YES's (or unknown) --- LAPSS screening criteria met:

YES NO


11. If LAPSS criteria for stroke are met, alert the receiving hospital of a possible stroke patient. If not, then return to the appropriate treatment protocol.

(Note: the patient may be experiencing a stroke even if the LAPSS criteria are not met.)

12. Time LAPSS Exam Performed: Military Time: _____

13. Form Completed by: _____

Pain Scale Forms




No Pain Moderate Pain Worst Possible Pain

0 1 2 3 4 5 6 7 8 9 10

**If you are having pain, tell your doctor or nurse.
Use these pain scales to describe your pain.**

No Hurt	Hurts Little Bit	Hurts Little More	Hurts Even More	Hurts Whole Lot	Hurts Worst
					

UNC Health Care Pain Management Committee
March 2001



Sin dolor Duele todavía más El peor dolor

0 1 2 3 4 5 6 7 8 9 10

**Si tiene dolor, dígaselo a su doctor o enfermera.
Use esta escala para describir su dolor.**

Sin Dolor	Duele un poquito	Duele un poquito más	Duele todavía más	Duele mucho	El peor dolor
					

UNC Health Care
Pain Management Committee
March 2001

From Hockenberry MJ, Wilson D, Winkelstein ML; Wong's Essentials of Pediatric Nursing, ed. 7, St. Louis, 2005, p. 1259. Used with permission. Copyright, Mosby.



Restraint Checklist

Patient's Name: _____

PCR Number: _____ Date: _____

It is recommended that a Restraint Checklist be completed with any restraint use.

1. Reason for restraint (check all that apply):

- Patient attempting to hurt self
- Patient attempting to hurt others
- Patient attempting to remove medically necessary devices

2. Attempted verbal reassurance / redirection?

- Yes
- No

3. Attempted environmental modification? (i.e. remove patient from stressful environment)

- Yes
- No

4. Received medical control order for restraints?

- Yes _____, MD
- No (Medical Control Physician Name Here)

5. Time and Type of restraint applied (check all that apply):

Date: ____/____/____ Time: ____AM/PM

Limb restraints:

- LUE
- RUE
- LLE
- RLE

Chemical Restraint:

- Yes
- No

If Yes: Drug Used: _____

Total Dose: _____

6. Vital signs and extremity neurovascular exam should be taken every 15 minutes.

7. Transport Position (Patient should NOT be in prone position)

- Supine position for transport
- Lateral recumbent position for transport

Signature: _____

(EMS Lead Crew Member)



Approved Medical Abbreviations

The following is a list of approved medical abbreviations. In general, the use of abbreviations should be limited to this list.

A&O x 3	- alert and oriented to person, place and time
A&O x 4	- alert and oriented to person, place, time and event
A-FIB	- atrial fibrillation
AAA	- abdominal aortic aneurysm
ABC	- airway, breathing, circulation
ABD	- abdomen (abdominal)
ACLS	- advanced cardiac life support
AKA	- above the knee amputation
ALS	- advanced life support
AMA	- against medical advice
AMS	- altered mental status
AMT	- amount
APPROX	- approximately
ASA	- aspirin
ASSOC	- associated
BG	- blood glucose
BILAT	- bilateral
BKA	- below the knee amputation
BLS	- basic life support
BM	- bowel movement
BP	- blood pressure
BS	- breath sounds
BVM	- bag-valve-mask
C-SECTION	- caesarean section
C-SPINE	- cervical spine
C/O	- complaint of (complains of)
CA	- cancer
CABG	- coronary artery bypass graft
CAD	- coronary artery disease
CATH	- catheter
CC	- chief complaint
CEPH	- cephalic
CHF	- congestive heart failure
CNS	- central nervous system
COPD	- chronic obstructive pulmonary disease
CP	- chest pain
CPR	- cardiopulmonary resuscitation
CSF	- cerebrospinal fluid
CT	- cat scan
CVA	- cerebrovascular accident (stroke)



Approved Medical Abbreviations

D5W	- 5% dextrose in water
DKA	- diabetic ketoacidosis
DNR	- do not resuscitate
DOA	- dead on arrival
DT	- delirium tremens
Dx	- diagnosis
ECG	- electrocardiogram
EEG	- electroencephelogram
ET	- endotracheal
ETOH	- ethanol (alcohol)
ETT	- endotracheal tube
EXT	- external (extension)
FB	- foreign body
FLEX	- flexion
Fx	- fracture
g	- gram(s)
GI	- gastrointestinal
GSW	- gunshot wound
gtts	- drops
GU	- gastrourinary
GYN	- gynecology (gynecological)
H/A	- headache
HEENT	- head, eyes, ears, nose, throat
HR	- heart rate (hour)
HTN	- hypertension
Hx	- history
ICP	- intracranial pressure
ICU	- intensive care unit
IM	- intramuscular
IV	- intravenous
JVD	- jugular vein distension
kg	- kilogram
KVO	- keep vein open



Approved Medical Abbreviations

L-SPINE	- lumbar spine
L/S-SPINE	- lumbar sacral spine
L&D	- labor and delivery
LAT	- lateral
lb	- pound
LLQ	- left lower quadrant
LMP	- last menstrual period
LOC	- level of consciousness (loss of consciousness)
LR	- lactated ringers
LUQ	- left upper quadrant
MAST	- military anti-shock trousers
mcg	- microgram(s)
MED	- medicine
mg	- milligram(s)
MI	- myocardial infarction (heart attack)
min	- minimum / minute
MS	- mental status
MS	- mental status change
MSO4	- morphine
MVC	- motor vehicle crash
N/V	- nausea/vomiting
N/V/D	- nausea/vomiting/diarrhea
NAD	- no apparant distress
NC	- nasal cannula
NEB	- nebulizer
NKDA	- no known drug allergies
NRB	- non-rebreather
NS	- normal saline
NSR	- normal sinus rhythm
OB/GYN	- obstetrics/gynecology
PALP	- palpation
PAC	- premature atrial contraction
PE	- pulmonary embolus
PEARL	- pupils equal and reactive to light
PMHx	- past medical history
PO	- orally
PRB	- partial rebreather
PRN	- as needed
PT	- patient
PVC	- premature ventricular contraction



Approved Medical Abbreviations

RLQ	- right lower quadrant
RUQ	- right upper quadrant
Rx	- medicine
RXN	- reaction
S/P	- status post
SOB	- shortness of breath
SQ	- subcutaneous
ST	- sinus tachycardia
SVT	- supraventricular tachycardia
Sx	- symptom
SZ	- seizure
T-SPINE	- thoracic spine
T	- temperature
TIA	- transient ischemic attack
TKO	- to keep open (refers to IV's - same as KVO)
Tx	- treatment
UOA	- upon our arrival
URI	- upper respiratory infection
UTI	- urinary tract infection
VF	- ventricular fibrillation
VS	- vital signs
VT	- ventricular tachycardia
WAP	- wandering atrial pacemaker
WNL	- within normal limits
YO (YOA)	- years old (years of age)
M or ♂	- male
F or ♀	- female
+	- positive
-	- negative
?	- questionable
Ψ	- psychiatric
~	- approximately
>	- greater than
<	- less than
=	- equal



Approved Medical Abbreviations

↑	- upper (increased)
ā	- before
p̄	- after
c̄	- with
s	- without
Δ	- change
L	- left
R	- right
↓	- lower (decreased)
1°	- primary
2°	- secondary



Reperfusion Checklist

The Reperfusion Checklist is an important component in the initial evaluation, treatment, and transport of patients suffering from an acute ST-elevation myocardial infarction (STEMI) or acute Stroke. Both of these conditions can be successfully treated using fibrinolysis (thrombolytics) if the patient arrives at the appropriate hospital within the therapeutic window of time.

This form should be completed for all acute STEMI and acute Stroke patients.

Patient's Name: _____

PCR Number: _____ Date: _____

1. Has the patient experienced chest discomfort for greater than 15 minutes and less than 12 hours?

Yes No

2. Has the patient developed a sudden neurologic deficit with a positive Los Angeles Prehospital Stroke Screen?

Yes No

3. Are there any contraindications to fibrinolysis?

If any of the following are checked "Yes", fibrinolysis MAY be contraindicated.

- Yes No Systolic Blood Pressure greater than 180 mm Hg
- Yes No Diastolic Blood Pressure greater than 110 mm Hg
- Yes No Right vs. Left Arm Systolic Blood Pressure difference of greater than 15 mm Hg
- Yes No History of structural Central Nervous System disease (tumors, masses, hemorrhage, etc.)
- Yes No Significant closed head or facial trauma within the previous 3 months
- Yes No Recent (within 6 weeks) major trauma, surgery (including laser eye surgery), gastrointestinal bleeding, or severe genital-urinary bleeding
- Yes No Bleeding or clotting problem or on blood thinners
- Yes No CPR performed greater than 10 minutes
- Yes No Currently Pregnant
- Yes No Serious Systemic Disease such as advanced/terminal cancer or severe liver or kidney failure.

4. (STEMI Patients Only) Does the patient have severe heart failure or cardiogenic shock?

These patients may benefit more from a percutaneous coronary intervention (PCI) capable hospital.

- Yes No Presence of pulmonary edema (rales greater than halfway up lung fields)
- Yes No Systemic hypoperfusion (cool and clammy)

If any contraindication is checked as "Yes" and an acute Stroke is suspected by exam or a STEMI is confirmed by ECG, activate the EMS Stroke Plan or EMS STEMI Plan for fibrinolytic ineligible patients. This may require the EMS Agency, an Air Medical Service, or a Specialty Care Transport Service to transport directly to an specialty center capable of interventional care within the therapeutic window of time.

Evaluating for the difficult airway

Between 1 – 3% of patients who require endotracheal intubation have airways that make intubation difficult. Recognizing those patients who may have a difficult airway allows the paramedic to proceed with caution and to keep as many options open as possible. It also allows the paramedic to prepare additional equipment (such as a cricothyrotomy kit) that may not ordinarily be part of a standard airway kit. The mnemonic LEMON is useful in evaluating patients for signs that may be consistent with a difficult airway and should raise the paramedic's index of suspicion.

Look externally


External indicators of either difficult intubation or difficult ventilation include: presence of a beard or moustache, abnormal facial shape, extreme cachexia, edentulous mouth, facial trauma, obesity, large front teeth or "buck teeth", high arching palate, receding mandible, short bull neck.

Evaluate 3-3-2 Rule

- 3 fingers between the patient's teeth (patient's mouth should open adequately to permit three fingers to be placed between the upper and lower teeth)
- 3 fingers between the tip of the jaw and the beginning of the neck (under the chin)
- 2 fingers between the thyroid notch and the floor of the mandible (top of the neck)

Mallampati

This scoring system is based on the work of Mallampati et al published in the Canadian Anaesthesia Society Journal in 1985. The system takes into account the anatomy of the mouth and the view of various anatomical structures when the patient opens his mouth as wide as possible. This test is performed with the patient in the sitting position, the head held in a neutral position, the mouth wide open, and the tongue protruding to the maximum. Inappropriate scoring may occur if the patient is in the supine position (instead of sitting), if the patient phonates or if the patient arches his or her tongue.



Class I (easy) = visualization of the soft palate, fauces, uvula, anterior and posterior pillars.

Class II = visualization of the soft palate, fauces and uvula.

Class III = visualization of the soft palate and the base of the uvula.

Class IV (difficult) = soft palate is not visible at all.

Obstruction?

Besides the obvious difficulty if the airway is obstructed with a foreign body, the paramedic should also consider other obstructers such as tumor, abscess, epiglottitis, or expanding hematoma.

Neck Mobility

Ask the patient to place their chin on their chest and to tilt their head backward as far as possible. Obviously, this will not be possible in the immobilized trauma patient.



Burns Resources

Fluid Formula

Formula for Fluid Resuscitation of the Burn Patient (Also known as the Parkland Formula)

Pts Wt kg x %TBSA x 4.0cc LR infused over 24 hours with half given in the first 8 hours.

(For the equation, the abbreviations are: PW x TBSA x 4.0 cc)

EMS focuses on the care given during the 1st hour or several hours following the event. Thus the formula as adapted for EMS and the first 8 hours is:

$$PW \times TBSA \times 4.0 \text{ cc, divide by 2}$$

to take this to the hourly rate, divide that solution by 8 and the equation becomes:

$$PW \times TBSA \times 4.0\text{cc} / 2 / 8 = \text{total to be infused for each of the first 8 hours.}$$

Another way to state the equation is to use:

$$PW \times TBSA \times 0.25\text{cc} = \text{total to be infused for each hour of the first 8 hours.}$$

Example. 80 kg patient with 50 %TBSA x 0.25 cc = 1000 cc/hr.


Remember:

Patient's Weight in kg (2.2 lbs = 1.0 kg) example: 220 lbs adult = 100 kg

% TSBA = Rule of Nine Total Body Surface Area

Factor for the 1st hr. and each hr. for the 1st 8 hrs. = 0.25

(Reminder, if two IV's are running, divide total amount to be infused each hr. by 2)



Wt (kg)	% TBSA	Factor	/Hr for 1st 8 Hrs of Care	60 gtt set, gtt/min	20 gtt set, gtt/min	15 gtt set, gtt/min	10 gtt set, gtt/min
10	10	0.25	25	25	8.3	6.3	4.2
10	20	0.25	50	50	16.7	12.5	8.3
10	30	0.25	75	75	25.0	18.8	12.5
10	40	0.25	100	100	33.3	25.0	16.7
10	50	0.25	125	125	41.7	31.3	20.8
20	10	0.25	50	50	16.7	12.5	8.3
20	20	0.25	100	100	33.3	25.0	16.7
20	30	0.25	150	150	50.0	37.5	25.0
20	40	0.25	200	200	66.7	50.0	33.3
20	50	0.25	250	250	83.3	62.5	41.7
30	10	0.25	75	75	25.0	18.8	12.5
30	20	0.25	150	150	50.0	37.5	25.0
30	30	0.25	225	225	75.0	56.3	37.5
30	40	0.25	300	300	100.0	75.0	50.0
30	50	0.25	375	375	125.0	93.8	62.5
40	10	0.25	100	100	33.3	25.0	16.7
40	20	0.25	200	200	66.7	50.0	33.3
40	30	0.25	300	300	100.0	75.0	50.0
40	40	0.25	400	400	133.3	100.0	66.7
40	50	0.25	500	500	166.7	125.0	83.3
50	10	0.25	125	125	41.7	31.3	20.8
50	20	0.25	250	250	83.3	62.5	41.7
50	30	0.25	375	375	125.0	93.8	62.5
50	40	0.25	500	500	166.7	125.0	83.3
50	50	0.25	625	625	208.3	156.3	104.2
60	10	0.25	150	150	50.0	37.5	25.0
60	20	0.25	300	300	100.0	75.0	50.0
60	30	0.25	450	450	150.0	112.5	75.0
60	40	0.25	600	600	200.0	150.0	100.0
60	50	0.25	750	750	250.0	187.5	125.0
70	10	0.25	175	175	58.3	43.8	29.2
70	20	0.25	350	350	116.7	87.5	58.3
70	30	0.25	525	525	175.0	131.3	87.5
70	40	0.25	700	700	233.3	175.0	116.7
70	50	0.25	875	875	291.7	218.8	145.8
80	10	0.25	200	200	66.7	50.0	33.3
80	20	0.25	400	400	133.3	100.0	66.7
80	30	0.25	600	600	200.0	150.0	100.0
80	40	0.25	800	800	266.7	200.0	133.3
80	50	0.25	1000	1000	333.3	250.0	166.7
90	10	0.25	225	225	75.0	56.3	37.5
90	20	0.25	450	450	150.0	112.5	75.0
90	30	0.25	675	675	225.0	168.8	112.5
90	40	0.25	900	900	300.0	225.0	150.0
90	50	0.25	1125	1125	375.0	281.3	187.5
100	10	0.25	250	250	83.3	62.5	41.7
100	20	0.25	500	500	166.7	125.0	83.3
100	30	0.25	750	750	250.0	187.5	125.0
100	40	0.25	1000	1000	333.3	250.0	166.7
100	50	0.25	1250	1250	416.7	312.5	208.3



Critical (Red)

>15% TBSA 2nd/3rd Degree Burn
 Burns with Multiple Trauma
 Burns with definitive airway compromise
 (When reasonable accessible, transport to a Burn Center)



Serious (Yellow)

5-15% TBSA 2nd/3rd Degree Burn
 Suspected Inhalation injury or requiring intubation for airway stabilization
 Hypotension
 GCS < 14
 (When reasonable accessible, transport to either a Level I Burn Center or a Trauma Center)



Minor (Green)

< 5% TBSA 2nd/3rd Degree Burn
 No inhalation injury, Not Intubated,
 Normotensive
 GCS > 14
 (Transport to the Local Hospital)

The NC Eye Bank EMS Referral Policy

Policy:

EMS will refer all appropriate field deaths to The North Carolina Eye Bank, using established criteria, in a timely and consistent manner.

Purpose:

- Enable the North Carolina Eye Bank to offer donation opportunities to families.
- Ensure the notification of The North Carolina Eye Bank for facilitating donation options.
- Honor donation wishes of registered donors at the time of death.

Procedure:

- EMS will call The North Carolina Eye Bank.
- EMS will use following criteria:
 1. Ages 2-75.
 2. Last Seen Alive Time <6 hours.
 3. No John/Jane Does. (unless pending investigation)
- Essential information that should be provided to The North Carolina Eye Bank is as follows:
 1. Caller name and title
 2. Patient demographics
 3. Last seen alive date and time/time of death
 4. Circumstances of death
 5. Next of kin name and contact information
 6. Where the body is going (ex: funeral home, hospital, M.E.)
- The North Carolina Eye Bank is responsible for approaching families about donation when appropriate.



The North Carolina
EYE BANK

24 hour Referral Hotline: 1-800-552-9956

North Carolina EMS Airway Evaluation Form

The NC EMS Airway Evaluation Form is required to be completed with all patients receiving Drug-Assisted Intubation in the Pre-hospital Environment.

FOR ORAL ROUTE:
Each Insertion of
Blade into Oropharynx = 1 Attempt

FOR NASAL ROUTE:
Pass of Tube Past the Nares =
1 Attempt

1. Patient Demographic Information

Date: ___/___/___ Dispatch Time: ___:___ am/pm

PCR # _____

EMS Agency Name: _____

Patient Age (yr): _____ Patient Sex: M F

2. Glasgow Coma Score (GCS) before intubation

Eye (1) (2) (3) (4)

Verbal (1) (2) (3) (4) (5)

Motor (1) (2) (3) (4) (5) (6)

3. Was ETI successful for the overall encounter?

Yes No Uncertain

4. Was intubation attempt due to Trauma?

Yes No

5. Level of training of each rescuer assisting with intubation

Rescuer A

State ID: _____

- Paramedic
- EMT-I
- Medic Student
- Nurse
- Phys. Assist
- MD/DO
- Other: _____

Rescuer B

State ID: _____

- Paramedic
- EMT-I
- Medic Student
- Nurse
- Phys. Assist
- MD/DO
- Other: _____

Rescuer C

State ID: _____

- Paramedic
- EMT-I
- Medic Student
- Nurse
- Phys. Assist
- MD/DO
- Other: _____

6. Indicate drugs given to facilitate intubation

- Atropine _____ mg
- Etomidate _____ mg
- Lidocaine _____ mg
- Midazolam _____ mg
- Rocuronium _____ mg
- Succinylcholine _____ mg
- Vecuronium _____ mg
- Other-Specify _____ - _____ mg
- Other-Specify _____ - _____ mg

7. Times and Vital Signs

	Time	Heart Rate	Resp. Rate	Blood Pressure	Pulse Oximetry	ECTO ₂
Pre-Airway Assessment Values	:			/		
Successful Airway Obtained	:					
Post-Airway Assessment Values	:			/		

8. Provide information for each laryngoscopy attempt.

Attempt	ETI Method	Rescuer	Successful?
1	<input type="checkbox"/> Direct <input type="checkbox"/> Nasal <input type="checkbox"/> Video	<input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C	<input type="checkbox"/> Yes <input type="checkbox"/> No
2	<input type="checkbox"/> Direct <input type="checkbox"/> Nasal <input type="checkbox"/> Video	<input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C	<input type="checkbox"/> Yes <input type="checkbox"/> No
3	<input type="checkbox"/> Direct <input type="checkbox"/> Nasal <input type="checkbox"/> Video	<input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C	<input type="checkbox"/> Yes <input type="checkbox"/> No
4	<input type="checkbox"/> Direct <input type="checkbox"/> Nasal <input type="checkbox"/> Video	<input type="checkbox"/> A <input type="checkbox"/> B <input type="checkbox"/> C	<input type="checkbox"/> Yes <input type="checkbox"/> No

9. Who verified placement of ET Tube?

- Rescuer performing intubation
- Another rescuer on the same team
- Receiving helicopter/EMS crew
- Receiving hospital team
- Other: _____

11. Endotracheal tube confirmation

	Auscultation	ETCO ₂	Breath Sounds	Absent Epigastric
Placement Confirmation	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>
Tube Size	Tube Depth			
Security Method				

10. If all attempts FAILED, indicate secondary airway technique used (Check all that apply)

- Bag-Valve-Mask (BVM) Combitube
- Open Cricothyroidotomy King LTD
- LMA Other _____

12. Were pulses maintained while under agencies care?

Yes No

13. Signature of Receiving Physician/Healthcare Provider (Confirming Destination/Transfer Tube Placement)

Yes No Uncertain

14. Signature of EMS Medical Director (Confirming Review of Completed Form)

Chart Review Done Remediation Required Approved

Date and Time: _____ : _____ am/pm

Date: _____

HIPAA PERMITS DISCLOSURE OF MOST TO OTHER HEALTH CARE PROFESSIONALS AS NECESSARY



Medical Orders
for Scope of Treatment (MOST)

This is a Physician Order Sheet based on the patient's medical condition and wishes. Any section not completed indicates full treatment for that section. **When the need occurs, first follow these orders, then contact physician.**

Patient's Last Name:	Effective Date of Form:
Patient's First Name, Middle Initial:	Patient's Date of Birth:

Section A
Check One Box Only

CARDIOPULMONARY RESUSCITATION (CPR): Patient has no pulse and is not breathing.
 Attempt Resuscitation (CPR) **Do Not Attempt Resuscitation (DNR/no CPR)**
 When not in cardiopulmonary arrest, follow orders in B, C, and D.

Section B
Check One Box Only

MEDICAL INTERVENTIONS: Patient has pulse and/or is breathing.
 Full Scope of Treatment: Use intubation, advanced airway interventions, mechanical ventilation, cardioversion as indicated, medical treatment, IV fluids, etc.; also provide comfort measures. **Transfer to hospital if indicated.**
 Limited Additional Interventions: Use medical treatment, IV fluids and cardiac monitoring as indicated. Do not use intubation or mechanical ventilation. May consider use of less invasive airway support such as BiPAP or CPAP. Also provide comfort measures. **Transfer to hospital if indicated. Avoid intensive care.**
 Comfort Measures: Keep clean, warm and dry. Use medication by any route, positioning, wound care and other measures to relieve pain and suffering. Use oxygen, suction and manual treatment of airway obstruction as needed for comfort. **Do not transfer to hospital unless comfort needs cannot be met in current location.**
 Other Instructions _____

Section C
Check One Box Only

ANTIBIOTICS
 Antibiotics if indicated
 Determine use or limitation of antibiotics when infection occurs
 No Antibiotics (use other measures to relieve symptoms)
 Other Instructions _____

Section D
Check One Box Only in Each Column

MEDICALLY ADMINISTERED FLUIDS AND NUTRITION: Offer oral fluids and nutrition if physically feasible.
 IV fluids if indicated **Feeding tube long-term if indicated**
 IV fluids for a defined trial period **Feeding tube for a defined trial period**
 No IV fluids (provide other measures to ensure comfort) **No feeding tube**
 Other Instructions _____

Section E
Check The Appropriate Box

DISCUSSED WITH AND AGREED TO BY:

<input type="checkbox"/> Patient	<input type="checkbox"/> Majority of patient's reasonably available parents and adult children
<input type="checkbox"/> Parent or guardian if patient is a minor	<input type="checkbox"/> Majority of patient's reasonably available adult siblings
<input type="checkbox"/> Health care agent	<input type="checkbox"/> An individual with an established relationship with the patient who is acting in good faith and can reliably convey the wishes of the patient
<input type="checkbox"/> Legal guardian of the patient	
<input type="checkbox"/> Attorney-in-fact with power to make health care decisions	
<input type="checkbox"/> Spouse	

Basis for order must be documented in medical record.

MD/DO, PA, or NP Name (Print):	MD/DO, PA, or NP Signature and Date (Required):	Phone #:
--------------------------------	---	----------

Signature of Patient, Parent of Minor, Guardian, Health Care Agent, Spouse, or Other Personal Representative
(Signature is required and must either be on this form or on file)

I agree that adequate information has been provided and significant thought has been given to life-prolonging measures. Treatment preferences have been expressed to the physician (MD/DO), physician assistant, or nurse practitioner. This document reflects those treatment preferences and indicates informed consent.

If signed by a patient representative, preferences expressed must reflect patient's wishes as best understood by that representative. Contact information for personal representative should be provided on the back of this form.

You are not required to sign this form to receive treatment.

Patient or Representative Name (print)	Patient or Representative Signature	Relationship (write "self" if patient)
--	-------------------------------------	--

SEND FORM WITH PATIENT/RESIDENT WHEN TRANSFERRED OR DISCHARGED

HIPAA PERMITS DISCLOSURE OF MOST TO OTHER HEALTH CARE PROFESSIONALS AS NECESSARY

Contact Information

Patient Representative:	Relationship:	Phone #: Cell Phone #:
Health Care Professional Preparing Form:	Preparer Title:	Preferred Phone #: Date Prepared:

Directions for Completing Form

Completing MOST

- MOST must be reviewed and prepared by a health care professional in consultation with the patient or patient representative.
- MOST is a medical order and must be signed and dated by a licensed physician (MD/DO), physician assistant, or nurse practitioner to be valid. **Be sure to document the basis for the order in the progress notes of the medical record.** Mode of communication (e.g., in person, by telephone, etc.) also should be documented.
- The signature of the patient or his/her representative is required; however, if the patient’s representative is not reasonably available to sign the original form, a copy of the completed form with the signature of the patient’s representative must be placed in the medical record and “on file” must be written in the appropriate signature field on the front of this form or in the review section below.
- Use of original form is required. **Be sure to send the original form with the patient.**
- MOST is part of advance care planning, which also may include a living will and health care power of attorney (HCPOA). If there is a HCPOA, living will, or other advance directive, a copy should be attached if available. **MOST may suspend any conflicting directions in a patient’s previously executed HCPOA, living will, or other advance directive.**
- **There is no requirement that a patient have a MOST.**
- MOST is recognized under N. C. G en. Stat. 90-21.17.

Reviewing MOST

Review of the MOST form is recommended when:

- The patient is admitted to and/or discharged from a health care facility; or
- There is a substantial change in the patient’s health status.

This MOST must be reviewed if:

- The patient’s treatment preferences change.

If MOST is revised or becomes invalid, draw a line through Sections A – E and write “VOID” in large letters.

Revocation of MOST

A patient with capacity or the patient’s representative (if the patient lacks capacity) can revoke the MOST at any time and request alternative treatment based on the known preferences of the patient or, if unknown, the patient’s best interests.

Review of MOST

Review Date	Reviewer and location of review	MD/DO, PA, or NP Signature (required)	Signature of patient or representative (preferred)	Outcome of Review
				<input type="checkbox"/> No Change <input type="checkbox"/> FORM VOIDED, new form completed <input type="checkbox"/> FORM VOIDED, no new form
				<input type="checkbox"/> No Change <input type="checkbox"/> FORM VOIDED, new form completed <input type="checkbox"/> FORM VOIDED, no new form
				<input type="checkbox"/> No Change <input type="checkbox"/> FORM VOIDED, new form completed <input type="checkbox"/> FORM VOIDED, no new form
				<input type="checkbox"/> No Change <input type="checkbox"/> FORM VOIDED, new form completed <input type="checkbox"/> FORM VOIDED, no new form
				<input type="checkbox"/> No Change <input type="checkbox"/> FORM VOIDED, new form completed <input type="checkbox"/> FORM VOIDED, no new form

SEND FORM WITH PATIENT/RESIDENT WHEN TRANSFERRED OR DISCHARGED

DO NOT ALTER THIS FORM!



Effective Date: _____

Expiration Date, if any _____

Check box if no expiration

DO NOT RESUSCITATE ORDER

Patient's full name _____

In the event of cardiac and/or pulmonary arrest of the patient, efforts at cardiopulmonary resuscitation of the patient SHOULD NOT be initiated. This order does not affect other medically indicated and comfort care.

I have documented the basis for this order and the consent required by the NC General Statute 90-21.17(b) in the patient's records.

Signature of Attending Physician/Physician Assistant/Nurse Practitioner

Printed Name of Attending Physician

Address

City, State, Zip

Telephone Number (office)

Telephone Number (emergency)

Do Not Copy

Do Not Alter



Recommended EMS Guidelines for Children and Youth with Special Health Care Needs (CYSHCN)*

Office of Emergency Medical Services
NC EMSC Advisory Committee
2009

Table of Contents

I. Important Considerations for Providing Care for Children and Youth with Special Health Care Needs.....	2
II. Emergencies involving Tracheostomies	3
III. Emergencies involving Indwelling Central Lines	5
IV. Emergencies involving Gastrostomy Tubes.....	6
V. Emergencies involving Ventilators	7
VI. Emergencies involving Urinary Drainage Catheters	9
VII. Emergencies involving Hemodialysis Lines	10
VIII. Emergencies involving Peritoneal Dialysis Catheters	11
IX. Care of the following equipment:	
a. Tracheostomy Tubes	3
b. Central Lines	5
c. Feeding Tubes	6
d. Ventilators.....	7
e. Apnea Monitors.....	13
f. VP Shunts	13
g. Internal Pacemakers	14
h. Vagal Nerve Stimulators	15
i. Colostomy Bags	15
Bibliography and Acknowledgements	16

*** These guidelines are meant to provide general guidance only and are not meant to supercede state and local medical protocols. Perform procedures in line with current scope of practice and consult local medical control when necessary.**

I. Important Considerations for Providing Care to Children and Youth with Special Health Care Needs

- Treat ABCs first
 - Treat the child, not the equipment
 - If the emergency is secondary to the child's equipment, use your own equipment
- Remember to always speak with the family since they are the experts on their child. Find out the child's baseline vital signs, medications, allergies, and other medical information, which may not be typical.
 - CYSHCN have many allergies. Ask about LATEX allergy, especially for patients with spina bifida. Make sure latex-free equipment is stocked ahead of time.
 - Ask for help from the parents and home health staff. They are generally trained to troubleshoot equipment and respond to emergencies. Practice TEAM (Trust Every Available Member)
- Physical handicaps do not necessarily imply mental deficits. Remember to communicate with the child. Assess and communicate with the child based on his/her developmental age, not chronological age.
- Ask for the "*go bag*" which generally has the child's spare equipment and supplies and bring this with you during transport. Also, this may have equipment you need on scene.
- Do not use excessive force to straighten or manipulate contracted extremities. The patient may be osteopenic and prone to fractures. (Some CYSHCN will not be able to straighten extremities beyond a nominal degree)
 - A slow, careful transfer with two or more people is preferable.
- Know which children in a given geographic area have special needs
 - Ask if they have a brief emergency medical information form, card, or notebook.
 - KIDBASE form
 - Look for MedicAlert® jewelry
- Transfer the child if possible to their medical "home" hospital.

II. Respiratory Distress in the Child with a Tracheostomy Tube

1. Secure the scene
2. ABCs
 - a. Open the airway using a head tilt/chin lift
3. Assess the tracheostomy tube
 - a. Is the tube in place?
 - b. Has the obturator (stylet) been removed?
 - c. In a double lumen trach tube, is the inner cannula in place?
 - d. Has a decannulation plug or speaking valve been removed?
4. Breathing
 - a. Assess rate, auscultation, inspection, effort and adequacy of chest rise
 - b. ALS: Check pulse oximetry, EtCO₂
 - c. Respiratory distress (retractions, altered mental status, hypoxia, etc.)?
 - i. Mucous or debris obstructing the tracheostomy tube is very common
 - d. Attempt to suction the trach tube.
 - i. Ask if family has a suction catheter. If so, use theirs, as it will be appropriately sized.
 - ii. If no suction catheter, ask family and use the size they generally use.
 - iii. If they do not know size, estimate the suction catheter size by doubling the inner diameter of trach and rounding down to an available catheter size, or use the largest size that will easily pass.
 - iv. Determine suction depth: ask family, or length of spare trach, or no more than 3-6 cm. **Suction with 100 mm Hg and instill 2-3 cc of saline before suctioning if secretions are thick.** Do not suction for more than 10 seconds and attempt to preoxygenate before getting started. **NEVER FORCE THE CATHETER.**
 - v. For a double lumen trach: take inner cannula out to suction then replace before assisted ventilations.
 - e. If unable to pass suction catheter the tracheostomy tube should be changed.
 - i. *Direct Technique:*
 - Allow the family to help change the tracheostomy tube.
 - For a double lumen trach, remove the inner cannula before insertion, and then, once inserted, replace inner cannula before confirming placement with BVM.
 - If the old trach has a cuff, deflate the cuff before removal.
 - Remove the old trach by pulling outward and toward the patient's feet.
 - Gently insert the new tracheostomy tube in the anatomical direction: curve downward and the tube aimed toward the

patient's feet... Make sure the obturator is in place for insertion then remove once trach is placed.

ii. *Facilitated Technique:*

- Allow the family to help change the tracheostomy tube.
 - If the old trach has a cuff, deflate the cuff before removal.
 - Remove the old trach by pulling outward and toward the patient's feet.
 - Before placing the new trach, slide a catheter through the trach.
 - Pass the suction catheter into the stoma and gently advance 3-6 cm
 - Advance the new trach over the suction catheter, using the catheter as a guide.
 - Once the trach is in place, remove the suction catheter.
- iii. If the new trach tube does not pass, attempt again with a smaller tube.
- iv. Confirm placement by assessment of breath sounds and adequacy of chest expansion.
- f. If a new tracheostomy tube cannot be placed, one is not available or it does not advance easily, place a similar or smaller internal-diameter size endotracheal tube (preferably cuffed) and advance only as far as a tracheostomy tube would have been advanced.
- g. If a new trach or ETT cannot be placed through the stoma, attempt orotracheal intubation if possible.
- i. If orotracheal intubation is unsuccessful, use mask-to-mouth bag ventilation with stoma occluded.
 - ii. If ventilation still inadequate, attempt infant mask-to-stoma bag ventilation.
- h. For severe respiratory distress despite suctioning, persistent hypoxia, or respiratory arrest, begin assisted ventilations through trach tube with appropriate ventilation bag with 100% high flow oxygen.

5. Reassess frequently. Monitor pulse oximetry and EtCO₂
6. If the patient has a trach and bronchospasm present: follow Wheezing protocol.
7. Assess circulation and follow appropriate protocols.
8. Keep warm. Expose only if necessary.
9. Contact Medical Control as necessary.
10. Remember: DOPE
- a. D-displaced, dislodged or damaged
 - b. O-obstructed (mucus, food, blood, secretions)
 - c. P-pulmonary problems

- d. E-equipment failure (bent tubing, ventilator malfunction, depleted oxygen supply)

III. Emergencies Involving Indwelling Central Lines

General Information:

- Types of central venous catheter:
 - Tunneled catheter-Broviac or Hickman
 - Implanted catheter-Mediport
 - Peripheral inserted catheter-PICC
 - These catheters are used to deliver nutritional substances or special medications directly into a central vein.
 - Most emergencies with lines include: blockage of the line, complete or partial accidental removal, or complete or partial laceration of the line
 - Children with indwelling catheters are always at risk for blood stream and catheter infections. Always use strict sterile technique when dressing or accessing the catheter.
1. Ask parents/caretakers about child's underlying condition: may be experiencing complications from underlying medical condition
 2. Obtain a complete medical history for the patient, including a history of the present illness and the past medical history
 3. Whenever assessing a child who has a central intravenous catheter, check the site where the tube is placed to see if it appears clean and well maintained.
 4. Identify location of central line:
 - a. Check for blockage of the line
 - b. Check for accidental removal or laceration of the line
 5. ***If line is blocked***, do not attempt to force catheter open
 6. ***If line is lacerated***, clamp proximal to laceration utilizing a padded clamp and do not use.
 7. ***If line is out or partially out:***
 - a. Do not push the line back in
 - b. Apply direct pressure to skin site
 - c. Stop any infusions*
 - d. Always bring line with you to the hospital
 8. Estimate blood loss and assess for signs and symptoms of an air embolism (tachypnea, chest pain, shortness of breath, or loss of consciousness) or blood clots. If an air embolism is suspected, clamp the central line with the clamp on the tube itself, place the child on the left side in a head down position, and administer high flow oxygen.

9. If the indwelling catheter is not damaged, is functioning, and does not have a continuous infusion already running, it may be used for fluid and medication administration.
 - a. Allow caregiver or home health personnel to access implanted catheters.
 - b. Use strict sterile technique when accessing an indwelling catheter.
 - c. In the event of a cardiac arrest, the indwelling central catheter is the preferred route of medication administration.

***There are some infusions that may be detrimental to stop, even briefly. Ask the caregiver if it is all right to stop or change the infusion first. Contact Medical Control for additional instructions.**

Summary:

D	Displaced, dislodged, or damaged	Stop infusing and do not use Direct pressure if bleeding from site Clamp or tie tubing if bleeding from catheter
O	Obstructed (blood clot, medication)	If line does not flush easily it needs to be replaced
P	Pulmonary Embolism	Clamp catheter and lie patient on left side with head down
E	Equipment failure (bent tubing, infusion pump malfunction)	If tube flushes easily the problem is probably with the pump

IV: Emergencies in Children with Gastrostomy Tubes and/or Feeding Tubes

Definitions:

- *Non-surgical feeding tubes:* Used for short term use:
 - Nasogastric tube (NGT): runs through the nose to the stomach
 - Nasojejunal Tube (NJT): runs through the nose into the small intestine
 - Orogastric tube (OGT): runs through the mouth into the stomach
- *Surgical Feeding Tubes:*
 - Gastrostomy Tube (GT): passes through the abdomen into the stomach
 - Jejunostomy Tube (JT): passes through the abdomen into the small intestine

Complications with gastrostomy tubes include: obstruction or dislodgement

1. When examining a child with a surgically implanted feeding tube, check for irritation and bleeding at the site where the tube enters the skin.
 - a. Treat minor bleeding with direct pressure and sterile dressings
 - b. A leaking feeding tube may cause skin irritation.
 - i. If there are any signs of infection at the entry site, the child should be transported for further medical attention.
 - ii. Cover the site with a sterile dressing and assess the abdomen.

2. Obstruction is usually not an emergency but the child requires transport. If the child is dependent on the feeds then the tube will need correction immediately.
3. Dislodgement is not life threatening but the tube should be replaced as soon as possible.
 - a. Keep the child flat on his/her back to prevent gastric fluid from leaking
 - b. If a new gastrostomy tube is available and stoma is open, attempt to reinsert the new tube.
 - c. If any resistance is met when inserting the gastrostomy tube STOP and cover the site with a clean dressing and assess the abdomen
 - d. If the new tube passes easily, secure with sterile dressing and tape BUT DO NOT REINFLATE BALLOON.
 - e. If caregiver is trained to replace gastrostomy tube, assist in placing new tube. If the new gastrostomy tube is successfully placed, DISCUSS TRANSPORT OPTIONS WITH CAREGIVER AND MEDICAL DIRECTOR
 - f. If no new gastrostomy tube is available, a foley catheter (same size or one size smaller) may be used and inserted...please follow same recommendations as above.
 - g. If tube does not pass easily: *Do not attempt to replace the tube; it is not as easy as it seems and there may be other complications.* Bring the dislodged tube with the child to the hospital.
 - h. Remember to cover the site with a clean dressing and control any bleeding with direct pressure
4. If there is formula infusing through the feeding tube, determine the nature of the fluids and the time that the fluids were started and stopped.
5. Assess for dehydration and/or hypoglycemia. Treat as necessary.
6. For non-surgical tubes (nasal or oral), assess for respiratory symptoms which may be a sign of placement in the respiratory tract.
 - a. If respiratory distress severe, remove tube carefully and treat respiratory symptoms.
7. Non-emergent transport to the nearest facility capable of replacing the tube.
8. If the parent has extra replacement tubes, bring these to the hospital.

V. Emergencies in Children on Ventilators

General Information:

- Children on mechanical ventilators may have a sudden or gradual deterioration, cardiac arrest, increased oxygen demand, increased respiratory rate, retractions, or change in mental status. This may be related to malfunction of the ventilator or due to worsening in their underlying disease.
- Common reasons for chronic mechanical ventilation in children include chronic respiratory failure and neurologic disease causing impaired airway control or respiratory effort.

- Some children requiring chronic mechanical ventilation never have a “normal” respiratory exam. Parents and other caregivers can provide information about the child’s baseline exam.

1. Pulse oximetry and End-tidal CO₂
2. If there is no increased respiratory distress, normal pulse oximetry, normal End-tidal CO₂, and normal mental status, the child should be transported on ventilator on current settings.
3. If there is respiratory distress, desaturation below baseline levels, or altered mental status:
 - a) Examine the child quickly for possible causes of distress which can be easily corrected: detached oxygen source, dislodged or obstructed tracheostomy tube, detached ventilator circuit.
 - b) Look at the ventilator and determine alarm code (i.e. apnea, low respiratory rate, low minute ventilation, high pressure, etc.) (See “Ventilator Troubleshooting” below)
 - Do not delay treatment while assessing the ventilator. Treat the patient, not the machine.
 - c) Remove the child from the ventilator and manually bag with a secure oxygen source
 - d) Look for normal chest rise, breath sounds on both sides, and improvement in oxygen saturation.
 - e) If the chest rise is shallow, adjust the patient’s airway position, check to see that the bag-valve device is securely connected to the tracheostomy tube, and use higher pressure if necessary.
 - f) Assess and treat problems with tracheostomy according to protocol.
4. Obtain relevant history of the present illness, past medical history and interventions taken to correct the emergency before EMS arrival.
5. Obtain any medical information forms that the caregivers may have for emergency medical providers.
6. Transport the child to the appropriate medical facility. Bring the ventilator to the hospital.
7. Some caregivers carry a “go bag” for their children with extra supplies. Bring this with the child if available.

Ventilator Troubleshooting

<i>Alarm</i>	<i>Possible Causes</i>	<i>Interventions</i>
Low pressure/apnea	Loose or disconnected circuit Leak in circuit Leak around tracheostomy site	Ensure all circuits are connected Check tracheostomy balloon Ensure tracheostomy well seated
Low power	Internal battery depleted	Plug the ventilator into a power outlet
High Pressure	Plugged or obstructed airway	Clear obstruction Suction tracheostomy

	Coughing/bronchospasm	Administer bronchodilator
Setting Error	Settings incorrectly adjusted	Manually ventilate patient Transport ventilator and patient
Power Switchover	Unit switched from AC to internal battery	Press “Alarm silent” button after ensuring battery is powering ventilator

Remember if the problem can not be remedied, EMS provider should remove the child from the ventilator, ventilate the child with a BVM, and take the ventilator with them to the hospital so a more qualified person can troubleshoot.

VI. Emergencies in Children with Urinary Drainage Catheters

General Information:

- Types of Urinary catheters:
 - Foleys: From urethra to bladder
 - Nephrostomy: From skin directly into kidney
 - Suprapubic: From skin directly into bladder
 - Ureterostomy: From skin into ureter
- These catheters are used to drain urine.
- Most emergencies with catheters include: Blockage, bleeding or dislodgement.

1. Ask caretakers about child’s underlying condition: may be experiencing complications from underlying medical condition
2. Obtain a complete medical history for the patient, including a history of the present illness and the past medical history
3. Most skin catheters will make the skin slightly erythematous and encrusted

If catheter is blocked:

- a. Flush once with 5cc of saline.
- b. Do not flush more than once.
- c. If catheter works well after flushing, discuss with caregiver and medical director.
- d. If catheter remains blocked, transport to appropriate medical facility.

If catheter is lacerated:

- a. Do not remove.
- b. Tape in place to avoid dislodgement
- c. Allow to continue to drain
- d. Transport to appropriate medical facility

If catheter is partially out:

- a. Do not push the line back in
- b. Secure to skin to avoid complete dislodgement
- c. Transport to appropriate medical facility

If catheter is completely out:

- a. Cover opening with sterile gauze
- b. Transport to appropriate medical facility

If blood is seen in catheter:

- a. Allow catheter to drain
- b. Secure to skin to prevent dislodgement
- c. Transport to appropriate medical facility

VII. Emergencies in Children with Hemodialysis Lines

General Information:

- Types of hemodialysis catheters:
 - Hemodialysis catheters (External tubing from a large artery to the skin)
 - Hemodialysis grafts (Gortex tubing under skin to artery and vein)
 - These catheters are used to filter/clean the blood in patients with renal failure.
 - Most emergencies with lines include: Infection of the line, bleeding from the line, and complete or partial dislodgement from trauma
 - Children with indwelling catheters are always at risk for blood and catheter infections. Always use strict sterile technique when dressing or accessing the catheter.
1. Ask caretakers about child's underlying condition: may be experiencing complications from underlying medical condition.
 2. Obtain a complete medical history for the patient, including a history of the present illness and the past medical history.
 3. Whenever assessing a child who has a hemodialysis catheter, check the site where the tube is placed to see if it appears clean and well maintained.
 4. Identify location of DIALYSIS LINE:
 - a. Check for accidental removal or laceration of the line
 5. ***If line is blocked*** DO NOT MANIPULATE
 6. ***If line is lacerated***, clamp proximal to laceration utilizing a padded clamp and do not use.
 7. ***If line is out or partially out:***
 - Do not push the line back in
 - Apply direct pressure to skin site
 - Stop any infusions*
 - Always bring line with you to the hospital

8. Estimate blood loss and assess for signs and symptoms of an air embolism (tachypnea, chest pain, shortness of breath, or loss of consciousness) or blood clots. If an air embolism is suspected, clamp the central line with the clamp on the tube itself, place the child on the left side in a head down position, and administer high flow oxygen.

9. If the indwelling catheter is not damaged, **UNLIKE CENTRAL LINES, do not use for IV access.** Infection and sepsis are frequent in large bore dialysis catheters compared to Broviacs.

10. In a life threatening emergency these large lines are excellent for IV access and can be used.

***There are some infusions that may be detrimental to stop, even briefly. Ask the caregiver if it is all right to stop or change the infusion first. Contact Medical Control for additional instructions.**

Transport all patients with hemodialysis lines to the appropriate medical facility.

VIII. Emergencies in Children with Peritoneal Dialysis Catheters

General Information:

- Peritoneal dialysis catheters run from the skin into the peritoneum.
- Dialysis is done by using the peritoneal lining as the dialysis membrane.
- Fluid is placed into the peritoneum and left for hours or overnight
- It is then drained removing extra electrolytes, acid, etc. from the patient.
- Dialysis is usually done at home by the patient or a nurse.
- Most emergencies with catheters include: Infection of the abdomen (peritonitis), infection of the catheter entry site, fracture of the catheter, bleeding from the catheter, and complete or partial dislodgement.
- Children with indwelling catheters are at risk for catheter infections. Always use strict sterile technique when dressing or accessing the catheter.
- These are not vascular lines and **can not** be used for IV access.

There are two major complications of peritoneal catheters: Infection and Outflow obstruction.

Exit Site Infections:

Drainage with blood and/or pus from the exit site

Associated with redness, tenderness, overgrown granulation tissue and swelling

Peritonitis:

Staphylococcus aureus

Caused by auto-inoculation by touch or contamination with respiratory secretions

Symptoms:

Abdominal pain
Abdominal tenderness
Abdominal distention
Cloudy peritoneal dialysis fluid
Fever
Nausea and vomiting

1. Ask Caretakers about child's underlying condition: may be experiencing complications from underlying medical condition

2. Obtain a complete medical history for the patient, including whether there is fluid presently in the abdomen or if it is drained.

3. Whenever assessing a child who has a peritoneal dialysis catheter, check the site where the tube is placed to see if it appears clean and well maintained.

4. ***If catheter is blocked*** DO NOT MANIPULATE.

5. ***If catheter is fractured***, clamp proximal to fractured utilizing a padded clamp and do not use.

6. ***If catheter is out or partially out:***

- Do not push back in
- Apply direct pressure to skin site if bleeding.
- Stop any infusions
- Always bring catheter with you to the hospital

7. ***If the catheter is leaking clear fluid:***

- Cover with sterile gauze.

Transport all patients with peritoneal dialysis catheters to the appropriate medical facility.

IX. Care of the Following Equipment:

- a. Tracheostomy tubes (included in protocol I.)
- b. Central lines (included in protocol III.)
- c. Feeding tubes (included in protocol IV.)
- d. Ventilators (included in protocol V.)
- e. Apnea monitors
- f. VP shunts
- g. Internal pacemakers
- h. Vagal nerve stimulators
- i. Colostomy bags

e. Apnea Monitors

ABCs

Pulse oximetry

If the patient is not breathing, open airway and begin bag-valve ventilation with 100% oxygen

Check the pulse: if no pulse, start chest compressions

Assess circulation and perfusion

Ask the caregiver for baseline vital signs

Look at the apnea monitor and determine the alarm code (i.e. heart rate, apnea etc.)

Check the electrodes or monitor chest belt and ensure proper placement

Make sure the monitor is powered and is not low on batteries

If the child has respiratory distress or cardiac arrest, call for ALS support and follow the appropriate algorithm and transport to the nearest appropriate facility.

Bring any of the child's emergency medical records and supplies or "go bag" with the patient to assist in the care of the child.

Bring the apnea monitor to the hospital with the child, so that it may be evaluated and stored information can be downloaded for analysis.

f. VP shunts

A cerebral spinal fluid shunt (CSF shunt) is a catheter that is inserted into the ventricles within the brain and then threaded under the skin from the skull to the right atrium (VA shunt) or the peritoneum of the abdomen (VP shunt). It drains excess CSF that would otherwise build up in the brain.

The child with a CSF shunt is vulnerable to brain infections. The shunt can develop an obstruction, and if this occurs it can result in any of the following signs & symptoms:

*Have a heightened awareness of the following:**

- Altered mental status
- Irritability
- Listlessness
- Increased sleep
- High-pitched cry
- Nausea and vomiting
- Fever
- Headaches
- Blurred vision
- Difficulty walking
- Apnea
- Bradycardia or other arrhythmias
- Seizures
- Redness along the shunt track
- Rapid worsening of mental status

Prehospital personnel should... Provide appropriate initial intervention and transport:

- Establish responsiveness
- Assess the patient's airway and breathing: ABCs
- Maintain a patent airway
- Provide high flow oxygen, positive pressure with bag-valve-mask mask if necessary
- Check pulse, if no pulse, begin chest compressions
- Assess circulation and perfusion
- Ask caregiver for the child's baseline vital signs
- Assess for signs and symptoms of shunt obstruction or infection*
- Obtain a complete history of present illness and past medical history
- Rapid transport to the appropriate facility

g. Internal Pacemakers

Pacemakers are implanted medical devices that regulate the heart rate.

For the child with an internal pacemaker, the following questions need to be asked:

- What type of heart problem does the child have?
- What is the child's baseline rhythm and rate?
- What type of pacemaker does the child have?
- Is the child dependent on the pacemaker?
- How long has the child had the pacemaker? (Generally 3-5yr battery life)

An internal cardiac defibrillator (ICD) or automatic implantable cardiac defibrillator (AICD) is an electronic device implanted under the skin. It monitors the heart rhythm and can slow down or stop excessively fast rates that originate in the ventricles.

For the child with an internal defibrillator:

- What type of heart problem does the child have?

What is the child's baseline rhythm and rate?

What heart rate causes the defibrillator to fire?

How many shocks has the child felt?

Has the child experienced any of the following?

Felt more than 3 shocks in a row

Unusual symptoms like dizziness or palpitations after a shock

Sensation of dizziness, lightheadedness or palpitations, for a period of time without any shocks

When was the defibrillator implanted? (3-5yr battery life)

EMS Care Tips

The internal pacemaker can easily be felt near the clavicle or in a small child in the abdomen.

Never place defibrillator paddles, or pacing patches directly over the internal pacemaker or defibrillator generator.

Remember the battery life is 3-5 years

Common Problem: Failure

1. Assess heart rate and perfusion
2. Treat for shock
3. Follow ABCs
4. Transport

h. Vagal Nerve Stimulators (VNS)

What is a vagal nerve stimulator? Device that is surgically implanted in the patient's chest, under the skin with the electrodes to the vagus nerve on the left side of the neck. This device produces electrical energy which works to dissipate seizures.

Ask the following questions:

Any recent trauma to the left side of neck or chest over the device?

Has the patient noticed anything different regarding the device?

When was the VNS implanted?

When was the VNS last checked?

What are the current settings?

Is the child having seizures when the device is functioning properly?

If seizures are still present, is the magnet being used?

Have you noticed any change in your child's seizures recently?

Increased intensity?

Increase in frequency?

i. Colostomies and Ileostomies

Colostomy or ileostomy: a portion of the large or small intestine is attached to the abdominal wall and an external bag is in place to collect the digestive waste.

Assess carefully for signs or symptoms of dehydration and/or shock, particularly if there has been any history of diarrhea or decreased oral intake.

Check the ostomy site for signs of infection or irritation:

Signs of infection include: red, warm, tender skin spreading away from the site

Ask the child or parents if the area is more tender than usual.

If any concerns, transport for further evaluation.

If the ostomy bag breaks, the parent or caregiver can usually help and replace it.

If another bag is not available, circle the ostomy with moist gauze and attach any available bag that can serve as a substitute until a proper replacement bag is obtained.

Bibliography and Acknowledgments:

1. Singh T., Wright JL., Adirim TA. Children with Special Health Care Needs: A Template for Prehospital Protocol Development. *Prehospital Emergency Care* July/Sept 2003;7(3):336-351.
2. EMSC Partnership for Children/National Association of EMS Physicians Model Pediatric Protocols: 2003 Revision. Pediatric Committee, National Association of EMS Physicians. *Prehospital Emergency Care* October/December 2004;8(4):343-365.
3. Adirim TA., Smith E., Singh T. Scope: Special Children's Outreach and Prehospital Education Jones and Bartlett Publishers 2006
4. Dieckmann RA. PEPP: Pediatric Education for Prehospital Professionals, 2nd edition. Jones and Bartlett Publishers Copyright 2006 by American Academy of Pediatrics.
5. State of Ohio EMS Board: Emergency Medical Services Pediatric Guidelines & Procedures Manual 2003
6. Illinois Emergency Medical Services for Children: Position Statement Pediatric Prehospital Protocols (Collaboration program between the Illinois Department of Public Health & Loyola University Medical Center) www.luhs.org/emsc March 18, 1999.
7. Children with Special Health Care Needs Provider Manual. Georgia Department of Human Resources, Division of Public Health.
8. Pediatric Special Health Care Needs Protocols: Children's National Medical Center for Prehospital Pediatrics, Division of Emergency Medicine and Trauma Services, Washington, D.C., November 2002.
9. New Your State's EMS-C: Children with Special Health Care Needs www.health.state.ny.us/nysdoh/ems/pdf/referencecard.pdf March 2003.

This document was compiled by the members of the North Carolina Emergency Medical Services for Children Advisory Committee. The main contributors were Donna Moro-Sutherland, MD (Chair of the EMSC Research and Education Committee) and Ben Alexander, MD (Past Chair of the EMSC Advisory Committee).

We want to thank all members of the committee who reviewed this document and took the time to forward their comments and suggestions which were incorporated into the document.